

08/11/00

Press Mail Label No. EL440514415US

8-14-00

A

**UTILITY PATENT APPLICATION TRANSMITTAL
(Small Entity)***(Only for new nonprovisional applications under 37 CFR 1.53(b))*Docket No.
42,982-C3-CPA-C (1095)Total Pages in this Submission
7**TO THE ASSISTANT COMMISSIONER FOR PATENTS****Box Patent Application
Washington, D.C. 20231**

Transmitted herewith for filing under 35 U.S.C. 111(a) and 37 C.F.R. 1.53(b) is a new utility patent application for an invention entitled:

THERAPEUTIC GUANIDINES

and invented by:

Stanley Goldin, et al.JCS957 U.S. PRO
09/637512
08/11/00If a **CONTINUATION APPLICATION**, check appropriate box and supply the requisite information:☒ **Continuation** ☐ **Divisional** ☐ **Continuation-in-part (CIP)** of prior application No.: 08/482,984

Which is a:

☒ **Continuation** ☐ **Divisional** ☐ **Continuation-in-part (CIP)** of prior application No.: PCT/US/95/01536

Which is a:

☐ **Continuation** ☐ **Divisional** ☒ **Continuation-in-part (CIP)** of prior application No.: 08/191,793

Enclosed are:

Application Elements

1. ☒ Filing fee as calculated and transmitted as described below
2. ☒ Specification having 145 pages and including the following:
 - a. ☒ Descriptive Title of the Invention
 - b. ☐ Cross References to Related Applications *(if applicable)*
 - c. ☐ Statement Regarding Federally-sponsored Research/Development *(if applicable)*
 - d. ☐ Reference to Microfiche Appendix *(if applicable)*
 - e. ☒ Background of the Invention
 - f. ☒ Brief Summary of the Invention
 - g. ☐ Brief Description of the Drawings *(if drawings filed)*
 - h. ☒ Detailed Description
 - i. ☒ Claim(s) as Classified Below
 - j. ☒ Abstract of the Disclosure

UTILITY PATENT APPLICATION TRANSMITTAL (Small Entity)

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Docket No.
42,982-C3-CPA-C (1095)

Total Pages in this Submission
7

Application Elements (Continued)

3. ☐ Drawing(s) (when necessary as prescribed by 35 USC 113)
- a. ☐ Formal b. ☐ Informal Number of Sheets _____
4. ☒ Oath or Declaration
- a. ☐ Newly executed (original or copy) ☐ Unexecuted
- b. ☒ Copy from a prior application (37 CFR 1.63(d)) (for continuation/divisional application only)
- c. ☒ With Power of Attorney ☐ Without Power of Attorney
- d. ☐ DELETION OF INVENTOR(S)
Signed statement attached deleting inventor(s) named in the prior application,
see 37 C.F.R. 1.63(d)(2) and 1.33(b).
5. ☒ Incorporation By Reference (usable if Box 4b is checked)
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
6. ☐ Computer Program in Microfiche
7. ☐ Genetic Sequence Submission (if applicable, all must be included)
- a. ☐ Paper Copy
- b. ☐ Computer Readable Copy
- c. ☐ Statement Verifying Identical Paper and Computer Readable Copy

Accompanying Application Parts

8. ☒ Assignment Papers (cover sheet & documents)
9. ☐ 37 CFR 3.73(b) Statement (when there is an assignee)
10. ☐ English Translation Document (if applicable)
11. ☒ Information Disclosure Statement/PTO-1449 ☐ Copies of IDS Citations
12. ☒ Preliminary Amendment
13. ☒ Acknowledgment postcard
14. ☒ Certificate of Mailing
- ☐ First Class ☒ Express Mail (Specify Label No.): EL440514415US

UTILITY PATENT APPLICATION TRANSMITTAL (Small Entity)

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Docket No.
42,982-C3-CPA-C (1095)

Total Pages in this Submission
7

Accompanying Application Parts (Continued)

15. ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)

16. ☒ Small Entity Statement(s) - Specify Number of Statements Submitted: One

17. ☐ Additional Enclosures (please identify below):

Fee Calculation and Transmittal

CLAIMS AS FILED

For	#Filed	#Allowed	#Extra	Rate	Fee
Total Claims	56	- 20 =	36	x \$9.00	\$324.00
Indep. Claims	9	- 3 =	6	x \$39.00	\$234.00
Multiple Dependent Claims (check if applicable) <input checked="" type="checkbox"/>					\$130.00
BASIC FEE					\$345.00
OTHER FEE (specify purpose)					\$0.00
TOTAL FILING FEE					\$1,033.00

- ☒ A check in the amount of **\$1,033.00** to cover the filing fee is enclosed.
- ☒ The Commissioner is hereby authorized to charge and credit Deposit Account No. **04-1105** as described below. A duplicate copy of this sheet is enclosed.
- ☐ Charge the amount of _____ as filing fee.
- ☒ Credit any overpayment.
- ☒ Charge any additional filing fees required under 37 C.F.R. 1.16 and 1.17.
- ☐ Charge the issue fee set in 37 C.F.R. 1.18 at the mailing of the Notice of Allowance, pursuant to 37 C.F.R. 1.311(b).

Dated: August 11, 2000


Signature

Jennifer K. Holmes (Reg. No. P-46,778)
Dike, Bronstein, Roberts & Cushman
Intellectual Property Practice Group
EDWARDS & ANGELL, LLP
130 Water Street
Boston, MA 02109
Tel: 617/523-3400

cc:

PATENT

Attorney's Docket No.

42982-C3

Applicant or Patentee:

S. Goldin et al.

Serial or Patent No.: 08 / 482,984

Filed or Issued: June 7, 1995

For: THERAPEUTIC GUANIDINES

COPY

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS (37 CFR 1.9(f) and 1.27(c))—SMALL BUSINESS CONCERN

I hereby declare that I am

- ☐ the owner of the small business concern identified below:
☒ an official of the small business concern empowered to act on behalf of the concern identified below:

NAME OF CONCERN Cambridge NeuroScience, Inc.

ADDRESS OF CONCERN One Kendall Square, Building 700
Cambridge, Massachusetts 02139

I hereby declare that the above identified small business concern qualifies as a small business concern as defined in 13 CFR 121.3-18, and reproduced in 37 CFR 1.9(d), for purposes of paying reduced fees under Section 41(a) and (b) of Title 35, United States Code, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third-party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed, to and remain with the small business concern identified above with regard to the invention, entitled

THERAPEUTIC GUANIDINES

by inventor(s) Stanley M. Goldin, James B. Fischer, Andrew Gannett Knapp, N. Laxma Reddy, David Berlove, Graham J. Durant, Subbarao Katragadda, Lain-Yen Hu,
Sharad Magar, Wenhong Fan, Elizabeth Yost and Jun Qing Guo
described in

- ☐ the specification filed herewith.
☒ application serial no. 08 / 482,984, filed June 7, 1995.
☐ patent no. _____, issued _____.

If the rights held by the above identified small business concern are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR 1.9(d) or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

*NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27).

08/06/95 14:30:00

COPY

NAME _____
ADDRESS _____

☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

NAME _____
ADDRESS _____

☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small business entity is no longer appropriate. (37 CFR 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING Gregory B. Butler, Ph.D., Esq.

TITLE OF PERSON OTHER THAN OWNER Senior Director, Legal Affairs and Chief

ADDRESS OF PERSON SIGNING One Kendall Square, Building 700 Patent Counsel
Cambridge, Massachusetts 02139

SIGNATURE Gregory B. Butler Date 11-21-95

Express Mail Label No. EL440514415US
Docket No. 42,982-C3-CPA-C (1095)

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICANT: S. Goldin, et al.
U.S.S.N.: Unassigned EXAMINER: Unassigned
FILED: Herewith GROUP: Unassigned
FOR: THERAPEUTIC GUANIDINES

CERTIFICATE OF EXPRESS MAILING

I hereby certify that this paper (along with any paper referred to as being attached or enclosed) is being deposited with the United States Postal Service on August 11, 2000, in an envelope as "Express Mail Post Office To Addressee", Label Number EL440514415US, addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

By


Donna M. Tomaso

ASSISTANT COMMISSIONER FOR PATENTS
WASHINGTON, D.C. 20231

Sir/Madam:

PRELIMINARY AMENDMENT

Please preliminarily amend the above-identified Continuation Application as follows:

IN THE SPECIFICATION:


On page 1, below the title, please insert --This Application is a Continuation of USSN 08/482,984, filed June 7, 1995, which is a Continuation of PCT/US95/01536, filed February 3, 1995, which is a Continuation-in-part of USSN 08/191,793, filed

S. Goldin, et al.
USSN: Unassigned
Filed: Herewith
Page 3

It is respectfully submitted that the subject application is in a condition for allowance. Early and favorable action is requested.

Applicants believe that additional fees are not required for consideration of the within Preliminary Amendment. However, if for any reason a fee is required, a fee paid is inadequate or credit is owed for any excess fee paid, you are hereby authorized and requested to charge Deposit Account No. **04-1105**.

Respectfully submitted,


Jennifer K. Holmes
Reg. No. P-46,778

Date: August 11, 2000

Dike, Bronstein, Roberts & Cushman
Intellectual Property Practice Group
EDWARDS & ANGELL, LLP
130 Water Street
Boston, MA 02109
Tel: (617) 523-3400

141485

U.S. PATENT & TRADEMARK OFFICE

THERAPEUTIC GUANIDINES

BACKGROUND OF THE INVENTION

This application is a continuation-in-part of copending International Application PCT/US95/01536 which designated the U.S., which in turn is a continuation of U.S. application serial number 08/191,793, filed February 3, 1994, both said application incorporated herein by reference.

1. Field of the Invention

The present invention pertains to certain substituted guanidines, and methods of treatment and pharmaceutical compositions that utilize or comprise one or more such guanidines.

10 2. Background

Neurons of the mature central nervous system ("CNS") are highly specialized and in general do not replace themselves. Consequently, death or degeneration of cells in the nervous system can have far more serious consequences than cell death or degeneration in other organs. Abnormal neuronal death can be rapid and widespread as in traumatic brain injury, or can occur over many years among very specific populations of neurons as in chronic neurodegenerative diseases.

Substantial evidence now points to pernicious overactivity of normal neurotransmitter systems as a contributory mechanism in many instances of pathological neuronal degeneration. In particular, overstimulation of neuronal receptors for L-glutamate, the brain's most prevalent excitatory amino acid ("EAA") neurotransmitter, has been recognized as a causal or exacerbating factor in several acute neurological disorders, and has been proposed to underlie a number of chronic neurodegenerative diseases as well [Choi, D.W., *Neuron.*, 1:623 (1988); Choi, D.W., *Cerebrov. and Brain Metab. Rev.*, 2:105 (1990); Albers, G.W., et al., *Ann. Neurol.*, 25:398 (1989)]. Indeed, it is believed that glutamate neurotoxicity is involved in acute injury to the nervous system as observed with seizure, hypoxia, hypoglycemia, and trauma, as well as in chronic degenerative diseases such as Huntington's disease, olivopontocerebellar atrophy associated with glutamate dehydrogenase deficiency and decreased glutamate catabolism, amyotrophic lateral sclerosis/Parkinsonium-dementia, Parkinson's disease,

and Alzheimer's disease [Choi, D.W., *Neuron*, 1:623-634 (1988); Choi, D.W., *Cereb. Brain Met., Rev.* 2:105-147 (1990); Courtier et al., *Lancet*, 341:265-268 (1993); Appel, S.H., *Trends Neurosci.*, 16:3-5 (1993)].

5 In the mammalian brain, glutamate interacts with three major classes of receptors, i.e., N-methyl-D-aspartate ("NMDA") receptors, non-NMDA receptors and metabotropic receptors [Watkins, J.D., et al., *Trends Neurosci.*, 10:265 (1987); and Seeburg, *TIPS*, 141:297 (1993)]. While triggering distinctive postsynaptic responses, all three classes of glutamate
10 receptors can act to increase the intracellular concentration of free Ca^{2+} in nerve cells [A.B. MacDermott, *Nature* 321:519 (1986)]. Thus, binding of glutamate to the NMDA receptor opens a cation-selective channel that is markedly permeable to Ca^{2+} , leading to a large and rapid increase in intracellular Ca^{2+} . A subclass of non-NMDA receptors has been found to be
15 linked to a Ca-permeable cation channel [Sommer, B., and Seeburg, P.H., *Trends Pharmacol. Sci.* 13:291-296 (1992)]. Although non-NMDA receptors are in most other instances linked to cation channels that largely exclude calcium, they can indirectly promote Ca^{2+} entry into neurons by depolarizing the cell membrane, which in turn opens voltage-activated Ca^{2+} -
20 channels. The so-called "metabotropic receptor", on the other hand, is not associated with an ion channel but can promote the release of Ca^{2+} from intracellular stores via the second-messenger inositol triphosphate.

 Irrespective of the triggering mechanism, prolonged elevation of
25 cytosolic Ca^{2+} is believed to be a key event in the initiation of neuronal destruction. Adverse consequences of elevated intracellular Ca^{2+} include derangement of mitochondrial respiration, activation of Ca^{2+} -dependent proteases, lipases and endonucleases, free radical formation and lipid peroxidation of the cell membrane [Choi, D.W., *Neuron*, 1:623-624 (1988)].

5 The NMDA subtype of excitatory amino acid receptors is strongly involved in nerve cell death which occurs following brain or spinal chord ischemia. Upon the occurrence of ischemic brain insults such as stroke, heart attack or traumatic brain injury, an excessive release of endogenous glutamate occurs, resulting in the over-stimulation of NMDA receptors. Associated with the NMDA receptor is an ion channel. The recognition site, i.e., the NMDA receptor, is external to the ion channel. When glutamate interacts with the NMDA receptor, it causes the ion channel to open, thereby permitting a flow of cations across the cell membrane, e.g., Ca^{2+} and Na^{+} into the cell and K^{+} out of the cell. It is believed that this flux of ions, especially the influx of Ca^{2+} ions, caused by the interaction of glutamate with the NMDA receptor, plays an important role in nerve cell death [see, e.g., Rothman, S.M. and Olney, J.W., *Trends in Neurosci.*, 10(7):299-302 (1987)]. Additionally, excessive excitation of neurons occurs in epileptic seizures and it has been shown that over-activation of NMDA receptors contributes to the pathophysiology of epilepsy [(Porter, R.J., *Epilepsia*, 30(Suppl. 1):S29-S34 (1989); and Rogawski, M.A., et al., *Pharmacol. Rev.*, 42:224-286 (1990)].

20 Non-NMDA receptors constitute a broad category of postsynaptic receptor sites which, as is the case for NMDA receptors, are directly linked to ion channels. Specifically, the receptor sites are physically part of specific ion channel proteins. Non-NMDA receptors have been broadly characterized into two major subclasses based on compounds selective therefor: kainate receptors and AMPA/quisqualate receptors [see J.C. Watkins et al., *Trends Neurosci.*, 10:265 (1987)]. AMPA is an abbreviation for α -amino-3-hydroxyl-5-methyl-4-isoazole propionic acid. These subclasses may be categorized as "non-NMDA" receptors.

30 Compared to NMDA receptors, non-NMDA receptors have received less pharmacological scrutiny -- the existing antagonists are all competitive -

- and *in vivo* research in this area has been hampered by the lack of drugs that cross the blood-brain barrier. Nonetheless, *in vivo* studies have clearly demonstrate that non-NMDA receptor agonists can also be as excitotoxic, although longer exposures can be required. In addition, evidence from
5 animal studies and from human epidemiological studies suggests that excitotoxicity mediated by non-NMDA receptors may be clinically important in certain pathologies. [see M.D. Ginsberg et al., *Stroke*, 20:1627 (1989)].

One such disorder is global cerebral ischemia or hypoxia, as occurs
10 following cardiac arrest, drowning, and carbon monoxide poisoning. Transient, severe interruption of the cerebral blood supply and/or interruption of the delivery of oxygen to the brain of animals causes a syndrome of selective neuronal necrosis, in which degeneration occurs among special populations of vulnerable neurons (including neocortical
15 layers 3, 5 and 6, pyramidal cells in hippocampal zones CA1 and CA3, and small and medium sized striatal neurons). The time course of this degeneration is also regionally variable, and can range from a few hours (striatum) to several days (hippocampus).

20 NMDA antagonists generally have not proven highly effective in animal models of global ischemia; indeed, it has been suggested that positive results obtained using NMDA antagonists may largely be the artifactual result of induction of hypothermia rather than due to direct inhibition of NMDA receptor-linked Ca entry into brain neurons [Buchan, A.
25 et al., *J. Neurosci.*, 11 (1991) 1049-1056]. In contrast, the competitive non-NMDA receptor antagonist 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo(F)quinoxaline ("NBQX") is dramatically effective in preventing delayed neuronal degeneration following transient forebrain ischemia in both gerbils and rats [M.J. Sheardown et al., *Science*, 247:571-574 (1990)].

At present, there is a critical need for effective treatments which limit the extent of nerve cell death following a stroke or traumatic brain injury. Recent advances in the understanding of the mechanisms underlying this nerve cell death have led to the hope that a drug treatment can be developed. Research and development efforts in this area have focussed on blocking the actions of glutamate that are mediated by the NMDA receptor-channel complex. Two approaches have been developed: competitive NMDA receptor antagonists [Choi D.W., *Cerebrov. Brain Metab. Rev.* 1:165-211 (1990); Watkins, J.C. and Olverman, H.J., *Trends Neurosci.*, 10:265-272 (1987)] and blockers of the ion channel of the NMDA receptor-channel complex [Meldrum, B., *Cerebrovascular Brain Metab. Rev.* 2:27-57 (1987); Choi, D.W., *Cerebrovascular Brain Metab. Rev.* 2:105-147 (1987); and Kemp, J.A. et al., *Trends Neurosci.*, 10:265-272 (1987)]. However, some toxicity with certain of the aforementioned agents has been reported [Olney, J.W. et al., *Science*, 244:1360-1362 (1989); Koek, W. and Colpaert, J., et al., *J. Pharmacol. Exp. Ther.*, 252:349-357 (1990)].

Blockers of neurotransmitter release, in particular blockers of the release of glutamate, have also received some attention as potential neuroprotective agents [see Meldrum, B., *Cerebrovascular and Brain Metab., Rev.* 2: 27-57 (1990); Dolphin, A.C. *Nature*, 316:148-150 (1985)]; Evans, M.C. et al., *Neurosci. Lett.*, 83:287-292 (1987); Ault, B. and Wang, C.M., *Br. J. Pharmacol.*, 87:695-703 (1986); Kaneko, T., et al., *Arzneim-Forsch./Drug Res.*, 39:445-450 (1989); Malgouris, C., et al., *J. Neurosci.*, 9:3720-3727 (1989); Jimonet, P. et al. *BioOrgan. and Med. Chem. Lett.*, 983-988 (1993); Wahl, F. et al., *Eur. J. Pharmacol.*, 230:209-214 (1993); Koek, J.W. and Colpaert, F.C., *J. Pharmacol. Exp. Ther.*, 252:349-357 (1990); Kaneko, T. et al., *Arzneim.-Forsch./Drug Res.*, 39:445-450 (1989)]. Certain compounds said to inhibit glutamate release also have been reported to show anticonvulsant activity [Malgouris, C., et al., *J. Neurosci.*, 9: 3720-

3727 (1989); Miller, A.A., et al., in *New Anticonvulsant Drugs*, Meldrum, B.S. and Porter R.J. (eds), London: John Libbey, 165-177 (1986)].

Calcium antagonists acting at L-type Ca channels such as nimodipine
5 have been reported to act both as cerebral vasodilators [Wong, M.C.W. et al., *Stroke*, 24:31-36 (1989)], and to block calcium entry into neurons [Scriabine, A., *Adv. Neurosurg.*, pp. 173-179 (1990)]. Modest improvement in the outcome of stroke has been observed in clinical trials [Gelmers, H.J. et al., *N. Eng. J. Med.*, 318:203-207 (1988)]. While there
10 are significant cardiovascular side effects, nimodipine appears less toxic in other respects than certain NMDA antagonists.


Antagonists of voltage-gated Na channels can exhibit neuroprotective properties. [Graham, S.H., et al., *J. Cereb. Blood Flow and Metab.*, 13:88-
15 97 (1993), Meldrum, B.S., et al., *Brain Res.*, 593:1-6 and Stys, P.K., et al., *J. Neurosci.*, 12: 430-439 (1992)]. In stroke, sustained hypoxia in the "core region" results from occlusion of the blood supply by a clot. As hypoxia develops, ATP depletion leads to an inability of the Na, K-ATPase to maintain the ion gradients which generate the normal membrane potential of
20 resting nerve cells. As the cell depolarizes and reaches the threshold for action potential firing, Na channels are activated. Stys et al. [Stys, et al., *J. Neurosci.*, 12: 430-439 (1992)] recently reported the development of Na channel hyperactivity in anoxia of central white matter and demonstrate *in vitro* the neuroprotective effect of the Na channel blockers tetrodotoxin
25 (TTX) and saxitoxin (STX).

SUMMARY OF THE INVENTION

The present invention provides therapeutically useful substituted guanidine compounds, including compounds that modulate, particularly
30 inhibit, the release of a neurotransmitter such as glutamate, and methods of treatment comprising such compounds. Preferred compounds of the

invention modulate, particularly inhibit, neurotransmitter (e.g., glutamate) release from ischemic neuronal cells, especially mammalian cells such as human neuronal cells. The compounds of the invention are useful for a number of therapeutic applications, including treatment of those diseases that result from modulation of a particular neurotransmitter system and that can be counteracted by one or more of the substituted guanidines of the invention which act either on the same or another class of neurotransmitters, and treatment of a variety of disorders of the nervous system and cardiovascular system, and of endocrine function.

In a first aspect, the present invention provides N,N-disubstituted guanidines of Formula I:




wherein:

R and R¹ are each independently substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least about 5 ring atoms, substituted or unsubstituted aralkyl having at least about 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having from 1 to 3 rings, 3 to 8 ring members in each

ring and from 1 to 3 hetero atoms, with at least one of R and R¹ being carbocyclic aryl, aralkyl, a heteroaromatic group or a heterocyclic group;


R² and R³ each being independently selected from the group consisting of hydrogen, substituted and unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted and unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted and unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted and unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted and unsubstituted alkylsulfonyl having from 1 to about 20 carbon atoms, and substituted and unsubstituted aminoalkyl; and pharmaceutically acceptable salts thereof.

A preferred group of compounds of Formula I are N,N-disubstituted compounds of the following Formula IA:



20 wherein R and R¹ are as defined above for Formula I, and pharmaceutically acceptable salts thereof.

A further preferred group of compounds of Formula I are compounds of the following Formula IB:




30 wherein R and R¹ are as defined above for Formula I, and R² and R³ each being independently selected from the group consisting of hydrogen,

substituted and substituted alkyl having from 1 to about 20 carbon atoms, substituted and unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted and unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted and unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted and unsubstituted alkylsulfonyl having from 1 to about 20 carbon atoms, and substituted and unsubstituted aminoalkyl, with at least one of R² and R³ being other than hydrogen; and pharmaceutically acceptable salts thereof.

Preferred compounds of Formulas I, IA or IB include those compounds where at least one, or more preferably both, of R and R¹ is substituted or unsubstituted carbocyclic aryl or substituted or unsubstituted aralkyl or substituted or unsubstituted alkaryl. Preferred compounds of Formulas I, IA and IB include those compounds having substituents with 1 to about 6 carbon atoms, particularly R² and/or R³ groups that have 1 to 6 carbon atoms. Particularly preferred R² and R³ substituents of compounds of Formulas I, IA or IB include unsubstituted alkyl and heteroalkyl such as alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl and aminoalkyl. Preferred R and R¹ groups include substituted and unsubstituted acenaphthyl, phenyl, biphenyl, naphthyl, fluorenyl and benzyl, particularly alkyl-substituted and alkoxy-substituted phenyl and benzyl. Particularly preferred R and R¹ groups include straight and branched chain C₁₋₈-alkyl substituted phenyl and benzyl such as *tert*-butylphenyl, *tert*-butylbenzyl, *sec*-butylphenyl, *sec*-butylbenzyl, *n*-butylphenyl, *n*-butylbenzyl, *iso*-butylphenyl, *iso*-butylbenzyl, pentylphenyl, pentylbenzyl, hexylphenyl, hexylbenzyl and the like; straight and branched chain C₁₋₈-alkoxy (including haloalkoxy, i.e. alkoxy substituted by F, Cl, Br and/or I) substituted phenyl and benzyl such as butoxyphenyl, butoxybenzyl, pentoxyphenyl, pentoxybenzyl, hexoxyphenyl, hexoxybenzyl, trifluoromethoxyphenyl, trifluorobenzyl, fluoro and the like; alkaryl (including alkoxyaryl) substituted phenyl and benzyl, particularly substituted and unsubstituted benzyl and benzyloxy (especially -OCH₂C₆H₅). Cycloalkyl and

aryl (particularly carbocyclic aryl) such as substituted phenyl, benzyl and naphthyl are also preferred R and R¹ groups such as biphenyl, phenylbenzyl (i.e. -CH₂C₆H₄C₆H₅), cyclohexylphenyl, cyclohexylbenzyl and the like. Halo (i.e., F, Cl, Br and/or I) substituted R and R¹ groups are also preferred including halo-substituted phenyl, naphthyl and benzyl.

In another aspect, the invention provides compounds of the following Formula II:



wherein

15 R is selected from the group of fluorenyl, phenanthracenyl, anthracenyl and fluoranthenyl;


R¹ is substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfonyl having from 1 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least about 5 ring atoms, substituted or unsubstituted aralkyl having at least about 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having 1 to 3 rings, 3 to 8 ring members in each ring and 1 to 3 heteroatoms;

25 R² and R³ are each independently hydrogen or a group as defined for R¹ above; and pharmaceutically acceptable salts thereof.

30

Preferred compounds of Formula II include N,N'-disubstituted compounds, i.e. where R² and R³ are each hydrogen, as well as tri- and tetra-substituted compounds where one or both of R² and R³ are other than hydrogen. Preferred R¹ groups include cycloalkyl, particularly adamantyl, and carbocyclic aryl, particularly substituted or unsubstituted phenyl, naphthyl or acenaphthyl, more preferably substituted or unsubstituted phenyl or naphthyl, such as alkyl or alkoxy substituted phenyl or naphthyl. Alkyl such as methyl, ethyl or propyl is a preferred R² or R³ group.

In a further aspect, the invention provides compounds of the following Formula III:



wherein R and R¹ are each independently substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aryloxy having from 6 to about 20 carbon atoms, substituted or unsubstituted aralkoxy having from 6 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfonyl having 1 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least 5 ring atoms, substituted or unsubstituted aralkyl having at least 5 ring atoms, or a substituted or unsubstituted heteroaromatic or

R² and R³ are each independently hydrogen or a group as defined for R and R¹ above, and preferably are each substituted or unsubstituted alkyl, alkenyl, alkynyl, alkoxy, aminoalkyl, alkylthio or alkylsulfinyl; or R¹ and R³ together form a ring having 5 or more ring members;

X and X' are each independently a chemical bond (i.e., a bond between the guanidine nitrogen and R or R¹), substituted or unsubstituted alkylene having from 1 to about 8 carbon atoms, substituted or unsubstituted alkenylene having from 2 to about 8 carbon atoms, or substituted or unsubstituted alkynylene having from 2 to about 8 carbon atoms, substituted or unsubstituted heteroalkylene having from 1 to about 8 carbon atoms, substituted or unsubstituted heteroalkenylene having 2 to about 8 carbon atoms, and substituted or unsubstituted heteroalkynylene having from 2 to about 8 carbon atoms, with at least one X and X' being other than a bond; and pharmaceutically acceptable salts thereof.


25



wherein the groups R, R¹, R² and R³ are as defined above for Formula III, and the value n is equal to 1, 2 or 3; and pharmaceutically acceptable salts thereof.

Preferred compounds of Formula III include those where R and R¹ together form a ring having 5 or more ring atoms, either with the guanidine nitrogen as the sole hetero atom or with one or more other N, O or S atoms as ring members, typically just one other N, O or S ring atom in addition to the guanidine N. Generally preferred is where R and R¹ together form a ring having 5-7 ring atoms, e.g., forming the following substituted or unsubstituted rings: morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, thiomorpholinyl, pyrrolidinyl, piperidinyl and tetrahydroquinolinyl. Preferred substituents of such rings include e.g. C₁₋₈alkyl, C₁₋₈alkoxy and substituted and unsubstituted alkaryl, particularly substituted and unsubstituted benzyl.

Particularly preferred such compounds of Formula III are those of the following Formula IIIB:



20 wherein R and R² of said formula are each independently substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aryloxy having from 6 to about 20 carbon atoms, substituted or unsubstituted aralkoxy having from 6 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfonyl having 1 to about 20 carbon atoms, substituted or unsubstituted

carbocyclic aryl having at least 5 ring atoms, substituted or unsubstituted aralkyl having at least 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having 1 to 3 rings, 3 to 8 ring members in each ring and 1 to 3 heteroatoms;


5 n is 1, 2, or 3, and preferably is 1 or 2, more is preferably 1; W is a carbon atom, or N, O or S; m is an integer of from 0 to 5, and preferably is 1, 2 or 3, more preferably 0, 1 or 2.

 X is substituted or unsubstituted alkylene having from 1 to about 8 carbon atoms, substituted or unsubstituted alkenylene having from 2 to
10 about 8 carbon atoms, or substituted or unsubstituted alkynylene having from 2 to about 8 carbon atoms, substituted or unsubstituted heteroalkylene having from 1 to about 8 carbon atoms, substituted or unsubstituted heteroalkenylene having 2 to about 8 carbon atoms, and substituted or unsubstituted heteroalkynylene having from 2 to about 8
15 carbon atoms, and preferably X is substituted or unsubstituted alkylene, particularly alkylene having 1 to 2 carbon atoms;

 each Y substituent is independently halogen, substituted or unsubstituted alkyl having 1 to about 10 carbon atoms, substituted or unsubstituted alkenyl having 2 to about 10 carbon atoms, unsubstituted
20 alkynyl having 2 to about 10 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 10 carbon atoms, substituted or unsubstituted alkylthio having 1 to about 10 carbon atoms, substituted or unsubstituted aminoalkyl having from 1 to about 10 carbon atoms, or substituted or unsubstituted carbocyclic aryl having about 6 or more ring
25 members; and pharmaceutically acceptable salts thereof.

 Particularly preferred compounds of Formula IIIB are those where R and R₂ are each independently aryl, particularly substituted or unsubstituted carbocyclic aryl such as substituted or unsubstituted phenyl, substituted or
30 unsubstituted naphthyl or substituted or unsubstituted acenaphthyl.

Especially preferred compounds of Formula IIIB are those where R and R₂ are each substituted or substituted phenyl, such as *sec*-butylphenyl or *tert*-butylphenyl, particularly para-*sec*-butylphenyl or para-*tert*-butylphenyl, n is 1 and X is alkylene of one or two carbons. Particularly preferred are compounds of the following Formula IIIBB:



wherein W is a carbon atom, or N, O or S;

15 each Y, each Y' and each Y'' is each independently halogen, substituted or unsubstituted alkyl having 1 to about 10 carbon atoms, substituted or unsubstituted alkenyl having 2 to about 10 carbon atoms, unsubstituted alkynyl having 2 to about 10 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 10 carbon atoms, substituted or unsubstituted alkylthio having 1 to about 10 carbon atoms, substituted or unsubstituted aminoalkyl having from 1 to about 10 carbon atoms, or substituted or unsubstituted carbocyclic aryl having about 6 or more ring members; n is 1 or 2, and each m, m' and m'' is independently an integer of from 0 to 5, and preferably is each m, m' and m'' is independently 0, 1, 2 or 3, more preferably 0, 1 or 2; and pharmaceutically acceptable salts thereof. Generally preferred compounds of Formulas IIIB and IIIBB are those where a Y group is bonded to the W ring member, particularly where W and Y together form a substituted carbon atom or N atom such as a C₁₋₈alkyl or C₁₋₈alkoxy substituted carbon or nitrogen ring atom. As will be of course understood by those skilled in the art, where m, m' or m'' is 0, the corresponding ring would be "fully" hydrogen-substituted. Specifically preferred compounds of Formula IIIBB include N-(4-butoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-piperidiny)guanidine; N-(4-butoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-benzylpiperidiny)guanidine; N-(4-butoxyphenyl)-N-(4-*tert*-

20

25

30

butylbenzyl)-N'-(4-morpholinyl)guanidine; and N-(4-butoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(3,5-dimethyl-4-morpholinyl)guanidine.


5 A further group of preferred compounds of Formula III are defined the same as Formulas IIIB and IIIBB above, but where two Y substituents are taken together to form an aryl or alicyclic fused ring. Generally preferred is where the fused ring is a heterocyclic or carbocyclic aryl, particularly phenyl, naphthyl, 1,2,3,6-tetrahydroquinoliny, thiomorpholinyl, pyrrolidinyl, piperazinyl and the like, or a cycloalkyl such as cyclohexyl. A specifically
10 preferred compound is N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(1,2,3,4-tetrahydroisoquinoliny)guanidine.

Preferred compounds of Formulas III, IIIA and IIIB include those where R and/or R¹ is substituted or unsubstituted carbocyclic aryl,
15 particularly substituted or unsubstituted phenyl, substituted or unsubstituted naphthyl or substituted or unsubstituted acenaphthyl. Particularly preferred are compounds of Formula III and IIIA where R, R¹ and R² are each substituted or unsubstituted carbocyclic aryl, particularly substituted or unsubstituted phenyl, substituted or unsubstituted naphthyl or substituted
20 or unsubstituted acenaphthyl. Especially preferred are compounds of Formula IIIA where R, R¹ and R² are each such a substituted or unsubstituted carbocyclic aryl, R₃ is hydrogen or C₁₋₄ alkyl such as methyl or ethyl, and n is 1 or 2. Of those especially preferred compounds one or more of R, R¹ and R² is preferably substituted or unsubstituted phenyl e.g.
25 C₁₋₈alkyl-substituted phenyl such as *sec*-butylphenyl *tert*-butylphenyl and the like, halo-substituted phenyl, C₁₋₈alkoxy-substituted phenyl such as butoxyphenyl or pentoxyphenyl or carbocyclic alkaryloxy-substituted phenyl such benzyloxyphenyl.

30 Preferred compounds of Formula III and IIIA also include N,N'-disubstituted compounds, i.e. where R² and R³ are each hydrogen, as well as tri- and tetra-substituted compounds where one or both of R² and R³ are

other than hydrogen. Preferred R^2 and R^3 groups include alkyl such as methyl, ethyl or propyl, and substituted alkyl, particularly haloalkyl such as C_1 - C_6 or C_1 - C_4 alkyl substituted by one or more F, Cl or Br. Alkylene and heteroalkylene are preferred X or X' groups, including those heteroalkylene groups containing 1 or 2 N, O, or S atoms as chain members. Particularly preferred X and X' groups of compounds of Formulas III, IIIA and IIIB include $-CH_2-$, $-CH_2CH_2-$, $-CH(CH_3)CH_2-$ and $-CH_2CH(CH_3)-$. Preferred compounds of Formulas III and IIIA include those where each R and R^1 group of a compound is bonded to the same carbon atom of the X or X' chain.

In a further aspect, compounds of the following Formula IV are provided:



wherein

each R is independently halo, hydroxy, amino, nitro, substituted or unsubstituted alkyl having from 3 to about 10 carbon atoms and preferably from 4 to about 10 carbon atoms, substituted or unsubstituted alkoxy, substituted or unsubstituted aryloxy, substituted or unsubstituted aralkoxy, substituted or unsubstituted aminoalkyl, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfinyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted alkenyl having 3 to about 10 carbon atoms, or substituted or unsubstituted alkynyl having 3 to about 10 carbon atoms;

n is an integer of from 1 to 5, preferably from 1 to 3;


R^1 is substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkoxy having from 1

to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having 1 to about 20 carbon atoms, substituted or unsubstituted aryloxy having from 6 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfonyl having 1 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least 5 ring atoms, substituted or unsubstituted aralkyl having at least 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having 1 to 3 rings, 3 to 8 ring members in each ring and 1 to 3 heteroatoms;

R^2 and R^3 are each independently hydrogen or a group as defined for R^1 above; or R^2 and R^3 are taken together to form a substituted or unsubstituted alkylene linkage of from 2 to about 6 carbon atoms; and pharmaceutically acceptable salts thereof.

Preferred compounds of Formula IV include those where one or more R substituents is a branched group such as *sec*-butyl or *tert*-butyl, or where one or more R substituent is substituted or unsubstituted aralkoxy, particularly substituted or unsubstituted benzyloxy. The value n is preferably 1, 2 or 3. Para-substitution and meta-substitution of the phenyl group by R substituent(s) is preferred. Preferred compounds of Formula IV include N,N'-disubstituted compounds, i.e. where R^2 and R^3 are each hydrogen, as well as tri- and tetra-substituted compounds where one or both of R^2 and R^3 are other than hydrogen. Alkyl such as methyl, ethyl or propyl is a preferred R^2 or R^3 group. Substituted or unsubstituted aryl such as substituted or unsubstituted phenyl or benzyl are preferred R^1 groups.


A particularly preferred group of compounds of Formula IV are those of the following Formula IV(A):



IV(A)

wherein each R' and R" are each independently selected from the same group as defined for R of Formula IV above; R² and R³ are defined the same as in Formula IV above; and n' and n" are each an integer of 1 to 5, and preferably are each 1 or 2; and pharmaceutically acceptable salts thereof. It is preferred that n' and n" are each 1, and R' and R" are each a meta or para substituent, more preferably each being a para substituent. It is further preferred that R' and R" are each branched substituents such as *tert*-butyl, *sec*-butyl, *iso*-butyl, *iso*-pentyl and the like. Preferably R² and R³ are hydrogen or C₁₋₄alkyl such as methyl or ethyl. Specifically preferred compounds of Formula IV(A) include N,N'-bis(3-*sec*-butylphenyl)guanidine and N,N'-bis(4-*tert*-butylphenyl)guanidine. In another aspect, the invention provides compounds of said Formula IV(A), particularly for use for the methods of treatment disclosed herein, but where excluded from said Formula IV(A) are the compounds of N,N'-bis(4-neopentylphenyl)guanidine, N,N'-bis(4-*t*-butylphenyl)guanidine, N,N'-bis(4-*n*-butylphenyl)guanidine, and N,N'-bis(4-cyclohexylphenyl)guanidine.

Preferred compounds of Formula IV, particularly for use in methods of treatment of the invention, also include those compounds wherein R² and R³ are taken together to form a substituted or unsubstituted alkylene linkage of from 3 to about 6 carbon atoms, particularly compounds with an alkylene linkage of 3 carbon atoms as represented of the following Formula IV(B):




wherein R, R' and n are each the same as defined above for Formula IV, and pharmaceutically acceptable salts thereof. Preferred R' groups of compounds of Formula IV(B) include substituted and unsubstituted alkaryl, particularly substituted and unsubstituted benzyl, and substituted and unsubstituted carbocyclic aryl, particularly substituted and unsubstituted

30

phenyl, substituted and unsubstituted naphthyl and substituted and unsubstituted acenaphthyl. Preferred substituents of such substituted phenyl, naphthyl or acenaphthyl R¹ groups include C₁₋₈alkyl, C₁₋₈alkoxy, C₁₋₈(mono- or dialkyl)amine, alkoxyaryland carbocyclic aryloxy, such as sec-butyl, tert-butyl, hexyl, butyloxy, phenoxy, benzyloxy, and the like.

Specifically, particularly for use in methods of treatment of the invention, are compounds of the above Formula IV(B) where R¹ is a substituted or unsubstituted phenyl, as represented by the following Formula IV(BB):



where each R and R' are each independently halo, hydroxy, amino, nitro, substituted or unsubstituted alkyl having from 3 to about 10 carbon atoms and preferably from 4 to about 10 carbon atoms, substituted or unsubstituted alkoxy, substituted or unsubstituted aryloxy, substituted or unsubstituted aralkoxy, substituted or unsubstituted aminoalkyl, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfinyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted alkenyl having 3 to about 10 carbon atoms, or substituted or unsubstituted alkynyl having 3 to about 10 carbon atoms; and

each n and n' is independently an integer of from 0 to 5, preferably from 1 to 5, more preferably 1 to 2 or 3. Particularly preferred is where each n and n' is 1. Preferred compounds of Formula IV(BB) include those where one or more R or R' substituents is a branched group such as a branched alkyl group e.g. *sec*-butyl or *tert*-butyl, or where one or more R substituent is substituted or unsubstituted aralkoxy, particularly substituted or unsubstituted benzyloxy. Para-substitution and meta-substitution of the

phenyl group by R and R' substituent(s) is typically preferred. Particularly preferred is where n and n' are each 1, and R and R' are each para substituents. Preferred R and R' substituents include alkoxy and alkoxyaryl, such as butoxy, pentoxy, hexoxy, substituted and unsubstituted benzyloxy and the like. Specifically preferred compounds of Formula IV(BB) include


5 N,N'-bis-(alkylphenyl)-2-iminopyrimidazolidine including N,N'-bis-(butylphenyl)-2-iminopyrimidazolidine, N,N'-bis-(pentylphenyl)-2-iminopyrimidazolidine, and N,N'-bis-(hexylphenyl)-2-iminopyrimidazolidine.

10 The invention also includes compounds of Formulas IV(B) or IV(BB) where two R substituents or two R' substituents together form a ring fused to the phenyl ring. Preferred fused rings have 5 to about 7 or 8 ring members and may be a carbocyclic aryl or saturated carbon ring, or a heteroaromatic or heteroalicyclic ring having 1 or 2 N, O or S atoms.

15 Exemplary fused rings include e.g. tetralinyl, indane, or a saturated six carbon ring to form a tetralinyl fused ring.

In another aspect, acenaphthyl-substituted guanidines of the following Formula V are provided:

20



25

wherein:

R is substituted or unsubstituted heteroaromatic containing 1-3 rings, 3 to 8 ring members in each ring and 1-3 heteroatoms;

R¹ and R² are each independently hydrogen, substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or

30 unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms,

substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfonyl having from 1 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least 5 ring atoms, substituted or unsubstituted aralkyl having at least 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having 1 to 3 rings, 3 to 8 ring members in each ring and 1 to 3 heteroatoms; and pharmaceutically acceptable salts thereof.

Preferred R groups of Formula V include 1,2,3,4,-tetrahydroquinoliny, indoliny, piperonyl, benz[cd]indoliny and [benz[cd]indo-2[1H]-one. The above depicted acenaphthyl group preferably is substituted by the guanidine nitrogen at the 3-position or 5-position. Preferred compounds of Formula V include N,N'-disubstituted compounds, i.e. where R¹ and R² are each hydrogen, as well as tri- and tetra-substituted compounds where one or both of R¹ and R² are other than hydrogen. Alkyl such as methyl, ethyl or propyl is a preferred R¹ and R² group of compounds of Formula V.

At least some compounds of the invention may exist as any one of a number of tautomeric forms. Each of these tautomeric forms are within the scope of the invention, including as defined by the formulas specified herein.

The present invention includes methods for treatment and/or prophylaxis of neurological conditions such as epilepsy, neurodegenerative conditions and/or nerve cell death resulting from, e.g., hypoxia, hypoglycemia, brain or spinal chord ischemia, brain or spinal chord trauma, stroke, heart attack, drowning or carbon monoxide poisoning. In this

regard, compound of the invention are particularly useful to administer to mammals, particularly humans, susceptible or suffering from stroke or heart attack. Compounds of the invention also are useful to treat and/or prevent various neurodegenerative diseases such as Parkinson's disease,

5 Huntington's disease, Amyotrophic Lateral Sclerosis, Alzheimer's disease, Down's Syndrome, Korsakoff's disease, olivopontocerebellar atrophy, HIV-induced dementia and blindness or multi-infarct dementia. Compounds of the invention also may be used to treat anxiety, e.g. by administration to subjects susceptible to generalized anxiety disorder. Compounds of the
10 invention will have particular utility for treatment of global cerebral ischemia as may occur following cardiac arrest, drowning and carbon monoxide poisoning. Compounds of the invention also may be used to treat other disorders of the nervous system, disorders of the cardiovascular system such as hypertension, cardiac arrhythmias or angina pectoris, endocrine
15 disorders such as acromegaly and diabetes insipidus, as well as use for treatment of chronic pain and as a local anesthetic. Compounds of the invention will have further utility for the treatment of those diseases in which the pathophysiology of the disorder involves excessive otherwise inappropriate (e.g., hypersecretory) cellular secretion, e.g., secretion of an
20 endogenous substance such as a catecholamine, a hormone or a growth factor. Exemplary diseases are specifically discussed *infra*. Compounds of the invention also will have utility for the treatment of those diseases in which the pathophysiology of the disorder involves excessive or otherwise inappropriate (e.g., hypersecretory) cellular secretion, e.g., secretion of an
25 endogenous substance such as a catecholamine, a hormone or a growth factor. The methods of treatment of the invention (which includes prophylactic therapy) generally comprise administration of a therapeutically effective amount of one or more compounds of the invention to an animal, including a mammal, particularly a human.

Further provided are diagnostic methods comprising use of the compounds of the invention. More specifically, a compound of the invention can be suitably labelled such as by radiolabelling a compound with ^{125}I , tritium, ^{32}P , ^{99}Tc , or the like, preferably ^{125}I . The labelled compound
5 can be administered to a subject such as a human and the subject imaged for a disease or disorder involving ion-channel activity such as stroke.

The invention also provides pharmaceutical compositions that comprise one or more compounds of the invention and a suitable carrier.
10

DETAILED DESCRIPTION OF THE INVENTION

It has been found that compounds of the invention have the ability to modulate, i.e. inhibit or potentiate the release of neurotransmitter(s), or to decrease or lengthen the time course of release of neurotransmitter(s), from neuronal tissue. It has thus been found that the compounds will have utility
15 to treat or prevent those pathophysiologic conditions which result from excessive or inadequate release of neurotransmitters. It is thought that substituted guanidines of the invention mediate the inhibition of neurotransmitter release by blocking presynaptic calcium channels and/or
20 sodium channels. Accordingly, the invention provides methods for blockage of voltage-activated calcium channels or sodium channels of neuronal cells, particularly mammalian cells such as human neuronal cells, comprising the administration to the cells of an effective amount of a compound of the invention, particularly such administration to a mammal in need of such
25 treatment. By such blockage of calcium channels and/or sodium channels of neuronal cells, conditions associated with excessive endogenous neurotransmitter release can be treated.

More particularly, some disorders such as neuronal damage in stroke
30 may be alleviated by inhibiting the release of excitatory amino acids such as glutamate. Some disorders such as depression may be alleviated by

inhibiting the release of inhibitory neurotransmitters such as gamma-aminobutyric acid. Further and without wishing to be bound by theory, inhibiting the release of an excitatory neurotransmitter such as glutamate by administration of a compound of the invention may indirectly potentiate the release or subsequent actions of an inhibitory transmitter such as gamma-aminobutyric acid, and thus the compound of the invention may treat disorders known to be alleviated by more direct potentiation of inhibitory neurotransmission, e.g., anxiety or insomnia.

10 Compounds of the invention may be considered effective inhibitors of neurotransmitter release if the compound causes at least about a 50% inhibition of neurotransmitter release, such as release of glutamate, at a concentration of about 100 μ M according to the protocol disclosed in Example 146 *infra*. More preferably the compound will cause at least about
15 a 50% inhibition of release of a neurotransmitter, such as glutamate, at a concentration of about 30 μ m according to the protocol disclosed in Example 146 *infra*. As used herein, the phrase "high inhibition of glutamate release" indicates that the specified compound(s) will cause at least 50% inhibition of glutamate release at a concentration of about 100 μ M
20 according to the protocol disclosed in Example 146 *infra*.

 Compounds of the invention may modulate release of neurotransmitters that include glutamate, dopamine, GABA (δ -amino butyric acid), norepinephrine, glycine, aspartate, serotonin, acetylcholine, adenosine
25 triphosphate and epinephrine, particularly glutamate. Compounds of the invention also may modulate release of peptides such as tachykinins, including substance P and substance k, enkephalins, luteinizing hormone-releasing hormone (LHRH) or derivatives thereof (see Harrington's Principles of Internal Medicine, 1705 (McGraw Hill 1987)), bombesin, cholecystokinin,
30 neuropeptide Y, dynorphin, gastrin-releasing hormone (GRH), neurotensin, somatostatin, and vasoactive intestinal peptide (VIP).

Specifically, compounds of the invention show significant ability to block depolarization-stimulated, calcium-dependent glutamate release from brian synaptosomes, when tested by the rapid superfusion method described in Example 146, *infra*. Said method identifies compounds that
5 act to block voltage-activated presynaptic Ca channels in nerve terminals, but may also identify compounds that block glutamate release by interfering with other processes involved in the control of Ca-dependent glutamate release. Compounds of the invention have demonstrated significant (\geq 50%) attenuation of glutamate release at concentrations of 10 μ M or less,
10 and many are effective at concentrations at or below 3 μ M. See, e.g., Table I of Example 146, *infra*. Moreover, these compounds of the invention exhibit relatively low binding affinity for the ion channel of the NMDA subclass of glutamate receptors and the di-tolyl guanidine (DTG) binding site associated with the sigma receptor. This indicates that compounds of the
15 invention have a clearly distinct therapeutic mechanism of action relative to that of known compounds which exhibit high affinity for either the ion channel of the NMDA subclass of glutamate receptors, and/or the DTG binding site of sigma receptors.

20 More particularly, a number of preferred compounds of the invention, including compounds of Formula IV and IV(A), will exhibit high inhibition of glutamate release (as that phrase is defined to mean above), but relatively low affinity for the PCP receptor, specifically an IC_{50} of 5 to 100 μ M or more in a typical PCP receptor assay as described in U.S. Patent No.
25 4906779 (see columns 10-11). A number of preferred compounds of the invention also will exhibit high inhibition of glutamate release, but relatively low affinity for the sigma receptor in a typical sigma receptor binding assay such as the assay disclosed in Weber et al., Proc. Natl. Acad. Sci. (USA), 83:8784-8788 (1986), specifically an IC_{50} of 10-200 μ M or more in the
30 DTG sigma binding assay disclosed in Weber et al., Proc. Natl. Acad. Sci. (USA), 83:8784-8788 (1986). As used herein, the phrase "low affinity to

the NMDA receptor" is intended to mean the specified compound(s) exhibits an IC₅₀ of 5 to 100 μ M or greater in said PCP receptor assay described in U.S. Patent No. 4906779; and the term "low affinity to the sigma receptor" is intended to mean the specified compound(s) exhibits an IC₅₀ of 10-200
5 μ M or greater in the DTG sigma binding assay disclosed in Weber et al., Proc. Natl. Acad. Sci. (USA), 83:8784-8788 (1986).

The preferred mechanism which underlies the ability of compounds of the invention to block neurotransmitter release *in vivo* is blockade of
10 voltage-activated Na channels and/or Ca channels which regulate neurotransmitter release [McBurney et al., *J. Neurotrauma*, 9, Suppl. 2:S531-S543 (1992); Kattragadda, S. et al., *Abs. Soc. for NeuroSci. Abs.*, 18:436 (1992)]. Examples 146-148 below describe the ability of compounds of the invention to block said voltage-activated Na and Ca
15 channels, providing further indication that compounds of the invention will effectively block release of a variety of neurotransmitters upon administration of the compounds to a mammal including a human.

Secretion of a variety of substances from non-neuronal secretory cells
20 occurs by a process of Ca-dependent exocytosis closely resembling the mechanism of Ca-dependent neurotransmitter release [Rubin, R.P., *Pharmacol. Rev.*, 22:389-428 (1970)]. Examples of this include release of norepinephrine from adrenal chromaffin cells [Landsberg, L. et al., *Harrington's Principles of Internal Medicine*, 11th Ed., eds., New York, McGraw-Hill, pp.358-370 (1987); Neher, E. et al., *Neuron*, 10:21-30
25 (1993)], secretion of peptide hormones from the pituitary [Tse et al., *Science*, 260:82-84 (1993); Chang, J.P. et al., *Endocrinology*, 123:87-94 (1988)], secretion of digestive enzymes from pancreatic acinar cells [Muallem, S., *Ann. Rev. Physiol.*, 51:83-105 (1989)], and secretion of
30 insulin from pancreatic beta cells [Larner, J., *The Pharmacological Basis of Therapeutics*, 7th Ed., eds., New York, MacMillan, pp. 1490-1503 (1985)].

The voltage-activated Ca channels that play a major role in governing said processes resemble those that govern neurotransmitter release, in terms of structure, pharmacology, and mechanism [Bean, B.P., *Ann. Rev. Physiol.* 51:367-384 (1989); Hess, P., *Ann. Rev. Neurosci.* 13:337-56 (1989);

5 Cohen, C.J. et al., *J. Physiol.*, 387:195-225 (1987)]. Accordingly the ability of compounds of the invention to block neurotransmitter release as described in Example 146, *infra* and the ability of said compounds to block the activity of voltage-activated Ca channels described in Examples 147 and 148, *infra*, constitutes strong evidence that compounds of the invention will

10 be effective inhibitors of exocytotic secretion of a variety of substances from a variety of non-neuronal cells. In particular, Example 148 demonstrates the ability of compounds of the invention to block Ca channels in GH4C1 clonal pituitary cells. Said pituitary cells secrete prolactin and growth hormone [Tashjian, A.H. *Meth. Enzymol.*, 58:527-535,

15 Acad. Press, N.Y. (1979)]. Block of voltage-activated Ca channels of pituitary GH4C1 cells inhibits secretion of said peptides [Tan, K.N. et al., *J. Biol. Chem.*, 259:418-426 (1984)]. Therefore, the results disclosed in Example 148 indicate compounds of the invention can function as effective antisecretory agents.

20

Example 147 *infra*, describes the ability of a subset of the compounds of the invention to block the presynaptic voltage-activated Ca channels which regulate release of neurotransmitters from mammalian brain synaptosomes. Said presynaptic Ca channels are structurally and

25 functionally related, and in some instances are identical, to Ca channels found in the cell bodies of neurons in the brain, elsewhere in the central nervous system, and in the peripheral nervous system [Zhang, J.F. et al., *Neuropharmacol.*, 32:1075-1088 (1993); Snutch, T.P. et al., *Curr. Opin. Neurobiol.*, 2:247-253 (1992)]. Therefore, the results disclosed in Example

30 147 indicate the general ability of compounds of the invention to block neuronal Ca channels.

Example 149 *infra*, describes the ability of a subset of the compounds of the invention to block Type II (alias R_{II}) voltage-activated Na channels, which are identical or closely related to the presynaptic and axonal voltage-activated Na channels which also govern release of neurotransmitters from mammalian brain synaptosomes [Westenbrook et al. *Neuron*, 3:695-704 (1989); Catterall, W.A., *Physiol. Rev.*, 72:S15-S48 (1992)]. Said presynaptic and axonal Type II Na channels are structurally and functionally similar, and in some instances are identical, to Na channels found in the cell bodies of neurons in the brain; elsewhere in the central nervous system; the peripheral nervous system; the cardiac conduction system comprising the Purkinje cell network and the atrial bundle branches; and cardiac, skeletal, and smooth muscle cells [Catterall, W.A., *ibid.*; Trimmer, J.S. et al., *Ann. Rev. Physiol.*, 51:401-418 (1989)].

Example 148 *infra* describes the ability of a subset of the compounds of the invention to block L-type voltage-activated Ca channels. L-type Ca channels regulate release of certain peptide neurotransmitters from mammalian brain nerve terminals [Rane et al., *Pflugers Arch.*, 409:361-366], and the secretion of a variety of substances such as peptide hormones of the pituitary [DeRiemer, S. A. et al., *Exp. Brain Res. Suppl.* 14:139-154 (1986)]. Said L-type Ca channels are structurally and functionally related, and in some instances are identical, to L-type Ca channels found in the cell bodies of neurons in the brain; elsewhere in the central nervous system; the peripheral nervous system; the cardiac conduction system comprising the Purkinje cell network and the atrial bundle branches; and cardiac, skeletal, and smooth muscle cells [Bean, B.P., *Ann. Rev. Physiol.*, 51:367-384 (1989); Snutch, T.P. et al., *Curr. Opin. Neurobiol.* 2:247-253 (1992)]. Therefore, the results disclosed in Example 148 indicate the ability of compounds of the invention to block said channels in the aforementioned tissues.

Compounds of the invention may be considered effective blockers of said voltage-activated Na and Ca channels if the compound causes at least a 50% reduction of the flow of cations through said channels at concentrations of about 100 μ M according to the protocols disclosed in

5 Examples 146-148, *infra*. More preferably the compound will cause at least a 50% inhibition of cation flow through said channels at a concentration of about 10 μ M according to the protocols of said Examples 146-148, *infra*.

10 Compounds of the invention have also demonstrated anticonvulsant activity in an in vivo as disclosed in Example 150, *infra*.

Suitable halogen groups of compounds of the invention (including compounds of Formulas I, IA, IB, II, III, IIIA, IIIB, IIIBB, IV, IV(A), IV(B), IV(BB) or V as specified above) are F, Cl, Br and I. Preferred alkyl groups
15 include those having 1 to about 12 carbon atoms, more preferably 1 to about 10 carbon atoms such as methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *t*-butyl, *iso*-butyl, *sec*-butyl, pentyl, hexyl, heptyl, etc. Preferred alkenyl and alkynyl groups include those groups having one or more unsaturated linkages, preferably one or two unsaturated linkages and from 2 to about 12
20 carbon atoms, more preferably 2 to about 8 carbon atoms. Each of the terms alkyl, alkenyl and alkynyl as used herein refer to both cyclic and noncyclic groups, although typically straight or branched chain noncyclic groups are generally more preferred. Preferred alkoxy groups of compounds of the invention include groups having one or more oxygen linkages and
25 from 1 to about 12 carbon atoms, more preferably 1 to about 8 carbon atoms, still more preferably 1 to about 6 carbons. Straight and branched chain butoxy, pentoxy, and hexoxy are particularly preferred. Preferred aryloxy groups have 6 to about 20 carbon atoms or from 6 to about 12 carbon atoms and include an oxygen atom. Substituted or unsubstituted
30 phenoxy and naphthoxy are preferred aryloxy groups. Preferred aralkoxy groups have from 6 to about 20 carbon atoms and include an alkoxy group

as specified above that contains one or more aryl substituents, particularly one or more carbocyclic aryl substituents. Typically an oxygen will be the terminal group of the substituent. Substituted or unsubstituted benzyloxy (i.e., $C_6H_5CH_2O-$) are preferred aralkoxy groups. Preferred thioalkyl groups include groups having one or more thioether linkages and from 1 to about 12 carbon atoms, more preferably 1 to about 8 carbon atoms, still more preferably 1 to about 6 carbons. Preferred aminoalkyl groups include those groups having one or more primary, secondary and/or tertiary amine groups, and from 1 to about 12 carbon atoms, more preferably 1 to about 8 carbon atoms, still more preferably 1 to about 6 carbons. Secondary and tertiary amine groups are generally more preferred than primary amine moieties. Preferred alkylsulfinyl groups have one or more sulfinyl (SO) groups, more typically one sulfinyl group, and from 1 to about 12 carbon atoms, more preferably 1 to about 6 carbons, and even more preferably 1-3 carbon atoms. Preferred alkylsulfonyl groups have one or more sulfono (SO_2) groups, more typically one sulfono group, and from 1 to about 12 carbon atoms, more preferably 1 to about 6 carbons, and even more preferably 1-3 carbon atoms. Preferred alkenylene and alkynylene X and X' groups of compounds of Formulas III and IIIA have one or two carbon-carbon multiple bonds. Preferred heteroalkylene, heteralkenylene and heteroalkynylene X and X' groups of compounds of Formulas III and IIIA contain 1 to about 3 hetero atoms consisting of N, O and/or S atoms, where one or more of said hetero atoms is a chain member of the X or X' group, and more preferably contain about 1-3 carbon atoms in addition to said hetero atoms. Suitable heteroaromatic and heteroalicyclic groups of compounds of the invention contain one or more N, O or S atoms and include, e.g., quinolinyl including 8-quinolinyl, indolinyl including 5-indolinyl, furyl, thienyl, pyrrolyl, thiazolyl, pyridyl, pyrimidinyl, pyridazinyl, oxazolyl and phthalimido groups all of which may be optionally independently substituted at one or more available positions and/or fused to a benzene ring; and substituted or unsubstituted tetrahydrofuranyl, tetrahydropyranyl, piperidinyl, piperazinyl, morpholino,

pyrrolidinyl groups, pyrazinyl, coumarinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, benzothiazolyl, benzotriazolyl, and bezimidazolyl. Preferred carbocyclic aryl groups include those having about 6 to about 20 carbons, more preferably about 1 to 3 separate or fused rings and from 6 to about 18 carbon atoms such as phenyl, naphthyl, acenaphthyl, phenanthryl, anthracyl and fluorene groups.

Said substituted moieties of compounds of the invention may be substituted at one or more available positions by one or more suitable groups such as, e.g., halogen such as F, Cl, Br, or I; cyano; hydroxyl; nitro; azido; carboxy; carbocyclic aryl; alkyl groups including alkyl groups having from 1 to about 12 carbon atoms or from 1 to about 6 carbon atoms; alkenyl and alkynyl groups including groups having one or more unsaturated linkages and from 2 to about 12 carbon atoms or from 2 to about 6 carbon atoms; alkoxy groups such as those groups having one or more oxygen linkages and from 1 to about 12 carbon atoms or from 1 to about 6 carbon atoms; thioalkyl groups such as those groups having one or more thioether linkages and from 1 to about 12 carbon atoms or from 1 to about 6 carbon atoms; aminoalkyl groups such as groups having one or more N atoms and from 1 to about 12 or 1 to about 6 carbon atoms; alkylsulfinyl such as those groups having one or more sulfinyl groups and from 1 to about 12 carbon atoms or from 1 to about 6 carbon atoms; alkylsulfonyl such as those groups having one or more sulfono groups and from 1 to about 12 carbon atoms or from 1 to about 6 carbon atoms.

25

Specifically preferred substituted groups include carboxylic acyl groups, preferably having from 1 to about 12 or 1 to about 6 carbon atoms such as acetyl, propanoyl, *iso*-propanoyl, butanoyl, *sec*-butanoyl, pentanoyl and hexanoyl groups. Also preferred substituted moieties are alkaryl groups which include single and multiple ring compounds, including multiple ring compounds that contain separate and/or fused aryl groups, e.g., above-

30

mentioned aryl groups substituted by one or more C₃-C₁₀ alkyl groups such as phenylpropyl, phenylbutyl, phenylpentyl and phenylhexyl groups as well as the branched chain isomers thereof such as *tert*-butylphenyl, *sec*-butylphenyl, etc. Haloalkyl and haloalkoxy are also preferred, particularly

5 fluoroalkyl and fluoroalkoxy such as trifluoromethyl and trifluoroalkoxy.

Aroyl groups are also preferred substituted groups such as carbonyl substituted by phenyl, naphthyl, acenaphthyl, phenanthryl, and anthracyl groups and carboxylic acyl groups substituted by one or more aryl groups, e.g., diphenylacetoxo and fluorene-carboxy groups. Aralkanoyl groups are

10 also preferred and include carbonyl substituted by the aralkyl groups described above. Aralkoxy groups are also preferred substituted groups and include alkoxy groups substituted by phenyl, naphthyl, acenaphthyl, phenanthryl, and anthracyl groups. Preferred substituted aryl groups include the above described aryl groups substituted by halo, hydroxy, alkyl, alkenyl,

15 alkynyl, alkoxy, amino, aminoalkyl, thioalkyl and the like.

Particularly preferred R and R¹ substituent groups of compounds of Formula I, as defined above, include substituted and unsubstituted carbocyclic aryl such as phenyl, butylphenyl including *tert*-butylphenyl,

20 cyclohexylphenyl, butoxyphenyl, trifluoromethoxyphenyl, halophenyl, methylthiophenyl, acenaphthyl, naphthyl including substituted naphthyl such as methoxy-1-naphthyl, and the like; and aralkyl groups including substituted and unsubstituted benzyl groups such as *tert*-butylbenzyl, *isopropyl*benzyl, halobenzyl, trifluoromethoxybenzyl, cinnamylmethylene,

25 naphthylmethylene, benzyloxy and the like.

Specifically preferred compounds of Formula I include:

N-(4-*sec*-butylphenyl)-N-benzylguanidine;

N-(5-acenaphthyl)-N-benzylguanidine;

30 N-(3-acenaphthyl)-N-benzylguanidine;

N-(5-acenaphthyl)-N-(4-*isopropyl*benzyl)guanidine;

- N-(3-acenaphthyl)-N-(4-*isopropylbenzyl*)guanidine;
N-(4-cyclohexylphenyl)-N-(4-*isopropylbenzyl*)guanidine;
N-(4-cyclohexylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(2-fluorenyl)-N-(4-*tert*-butylbenzyl)guanidine;
5 N-(4-*sec*-butylphenyl)-N-(cinnamylmethylene)guanidine;
N-(4-*n*-butoxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-biphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(5-indanyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-trifluoromethoxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
10 N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(5-acenaphthyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-acenaphthyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(methoxy-1-naphthyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(1-naphthyl)-N-(4-*tert*-butylbenzyl)guanidine;
15 N-(3-iodophenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(4-chloro-1-naphthyl)-N-(4-*tert*-benzyl)guanidine;
N-(4-*tert*-butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(4-iodophenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(1-naphthylmethyl)-N-(4-*tert*-butylbenzyl)guanidine;
20 N-(5-acenaphthyl)-N-(3-phenoxybenzyl)guanidine;
N-(3-trifluoromethylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-methylthiophenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(5-acenaphthyl)-N-(3-iodobenzyl)guanidine;
N-(5-acenaphthyl)-N-(cinnamyl)guanidine;
25 N-(5-acenaphthyl)-N-(4-iodobenzyl)guanidine;
N-(5-acenaphthyl)-N-(4-trifluoromethoxybenzyl)guanidine;
and pharmaceutically acceptable salts thereof.

Specifically preferred compounds of Formula II include:

- 30 N,N'-bis(2-fluorenyl)guanidine;
N,N'-bis(2-fluorenyl)-N-methylguanidine;

- N,N'-bis(2-fluorenyl)-N,N'-dimethylguanidine;
N,N'-bis(anthracenyl)guanidine;
N,N'-bis(anthracenyl)-N-methylguanidine;
N,N'-bis(anthracenyl)-N,N'-dimethylguanidine;
5 N,N'-bis(phenanthracenyl)guanidine;
N,N'-bis(phenanthracenyl)-N-methylguanidine;
N,N'-bis(phenanthracenyl)-N,N'-dimethylguanidine;
N,N'-bis(fluoranthenyl)guanidine;
N,N'-bis(fluoroanthenyl)-N-methylguanidine;
10 N,N'-bis(fluoroanthenyl)-N,N'-dimethylguanidine;
N-(anthracenyl)-N'-(1-adamantyl)guanidine;
N-(anthracenyl)-N'-(1-adamantyl)-N-methylguanidine;
N-(anthracenyl)-N'-(1-adamantyl)-N'-methylguanidine;
N-(anthracenyl)-N'-(1-adamantyl)-N,N'-dimethylguanidine;
15 N-(anthracenyl)-N'-(2-adamantyl)guanidine;
N-(anthracenyl)-N'-(2-adamantyl)-N-methylguanidine;
N-(anthracenyl)-N'-(2-adamantyl)-N'-methylguanidine;
N-(anthracenyl)-N'-(2-adamantyl)-N,N'-dimethylguanidine;
N-(phenanthracenyl)-N'-(1-adamantyl)guanidine;
20 N-(phenanthracenyl)-N'-(1-adamantyl)-N-methylguanidine;
N-(phenanthracenyl)-N'-(1-adamantyl)-N'-methylguanidine;
N-(phenanthracenyl)-N'-(1-adamantyl)-N,N'-dimethylguanidine;
N-(phenanthracenyl)-N'-(2-adamantyl)guanidine;
N-(phenanthracenyl)-N'-(2-adamantyl)-N-methylguanidine;
25 N-(phenanthracenyl)-N'-(2-adamantyl)-N'-methylguanidine;
N-(phenanthracenyl)-N'-(2-adamantyl)-N,N'-dimethylguanidine;
N-(fluorenyl)-N'-(1-adamantyl)guanidine;
N-(fluorenyl)-N'-(1-adamantyl)-N-methylguanidine;
N-(fluorenyl)-N'-(1-adamantyl)-N'-methylguanidine;
30 N-(fluorenyl)-N'-(1-adamantyl)-N,N'-dimethylguanidine;
N-(fluorenyl)-N'-(2-adamantyl)guanidine;

N-(fluorenyl)-N'-(2-adamantyl)-N-methylguanidine;
N-(fluorenyl)-N'-(2-adamantyl)-N'-methylguanidine;
N-(fluorenyl)-N'-(2-adamantyl)-N,N'-dimethylguanidine;
N-(fluorenyl)-N'-(methoxynaphthyl)guanidine;
5 N-(fluorenyl)-N'-(methoxynaphthyl)-N-methylguanidine;
N-(fluorenyl)-N'-(methoxynaphthyl)-N'-methylguanidine;
N-(fluorenyl)-N'-(methoxynaphthyl)-N,N'-dimethylguanidine;
and pharmaceutically acceptable salts of said compounds.

- 10 Particularly preferred R substituent groups of compounds of Formulas III and IIIA, as those formulas are defined above, include phenyl and naphthyl. Preferred values of n, n' and m of compounds of Formulas III and IIIA are 1 and 2. The R groups are suitably the same where n is greater than one. Carbocyclic aryl groups of compounds of Formulas III and IIIA are preferred R¹ groups, particularly substituted and unsubstituted phenyl,
15 naphthyl and acenaphthyl. Preferred X groups of compounds of Formulas III and IIIA include alkenylene groups substituted by halogen of F, Cl, Br and I; aryloxy such as substituted and unsubstituted phenoxy; aryl such as substituted and unsubstituted phenyl including phenyl substituted by C₁-C₆
20 alkyl, C₁-C₆ alkoxy, halogen such as F, Cl, Br or I.

Specifically preferred compounds of Formulas III and IIIA include:

- N-5-acenaphthyl-N'-benzhydrylguanidine;
N-5-acenaphthyl-N'-benzhydryl-N-methylguanidine;
25 N-5-acenaphthyl-N'-benzhydryl-N'-methylguanidine;
N-5-acenaphthyl-N'-benzhydryl-N,N'-dimethylguanidine;
N-3-acenaphthyl-N'-benzhydrylguanidine;
N-3-acenaphthyl-N'-benzhydryl-N-methylguanidine;
N-3-acenaphthyl-N'-benzhydryl-N'-methylguanidine;
30 N-3-acenaphthyl-N'-benzhydryl-N,N'-dimethylguanidine;
N-(5-acenaphthyl)-N'-[(1-naphthyl)-methyl]guanidine;

- N-(5-acenaphthyl)-N'-[(1-naphthyl)-methyl]-N-methylguanidine;
N-(5-acenaphthyl)-N'-[(1-naphthyl)-methyl]-N'-methylguanidine;
N-(5-acenaphthyl)-N'-[(1-naphthyl)-methyl]-N,N'-dimethylguanidine;
N-(5-acenaphthyl)-N'-(1-methyl-2-phenoxyethyl)guanidine;
5 N-(5-acenaphthyl)-N'-(1-methyl-2-phenoxyethyl)-N-methylguanidine;
N-(5-acenaphthyl)-N'-(1-methyl-2-phenoxyethyl)-N'-methylguanidine;
N-(5-acenaphthyl)-N'-(1-methyl-2-phenoxyethyl)-N,N'-dimethylguanidine;
N-(5-acenaphthyl)-N'-(1-methyl-2-(4-chlorophenyl)ethyl)guanidine;
N-(5-acenaphthyl)-N'-(1-methyl-2-(4-chlorophenyl)ethyl)-N-methylguanidine;
10 N-(5-acenaphthyl)-N'-(1-methyl-2-(4-chlorophenyl)ethyl)-N'-methylguanidine;
N-(5-acenaphthyl)-N'-(1-methyl-2-(4-chlorophenyl)ethyl)-N,N'-
dimethylguanidine;
N-(5-acenaphthyl)-N'-(1,2-diphenylethyl)guanidine;
N-(5-acenaphthyl)-N'-(1,2-diphenylethyl)-N-methylguanidine;
15 N-(5-acenaphthyl)-N'-(1,2-diphenylethyl)-N'-methylguanidine;
N-(5-acenaphthyl)-N'-(1,2-diphenylethyl)-N,N'-dimethylguanidine;
N-(5-acenaphthyl)-N'-(3-phenylpropyl)guanidine;
N-(5-acenaphthyl)-N'-(3-phenylpropyl)-N-methylguanidine;
N-(5-acenaphthyl)-N'-(2-methyl-2-phenylethyl)-N'-methylguanidine;
20 N,N'-(*sec*-butylphenyl)-N'-(2-phenoxyethyl)guanidine;
N,N'-(*sec*-butylphenyl)-N'-(2-phenoxyethyl)-N-methylguanidine;
N,N'-(*sec*-butylphenyl)-N'-(2-phenoxyethyl)-N'-methylguanidine;
N,N'-(*sec*-butylphenyl)-N'-(2-phenoxyethyl)-N,N'-dimethylguanidine;
N-(5-acenaphthyl)-N'-((4-*tert*-butylphenyl)-(4-*sec*-butylphenyl)-
25 methyl)guanidine;
N-(5-acenaphthyl)-N'-((4-*tert*-butylphenyl)-(4-*sec*-butylphenyl)-methyl)-N-
methylguanidine;
N-(5-acenaphthyl)-N'-((4-*tert*-butylphenyl)-(4-*sec*-butylphenyl)-methyl)-N'-
methylguanidine;
30 N-(5-acenaphthyl)-N'-((4-*tert*-butylphenyl)-(4-*sec*-butylphenyl)-methyl)-N,N'-
dimethylguanidine;

N-(4-butoxyphenyl)-N,N'-bis(4-*tert*-butylbenzyl)guanidine;
N-(4-butoxyphenyl)-N,N'-bis(4-*tert*-butylbenzyl)-N-methylguanidine;
N-(4-butoxyphenyl)-N,N'-bis(4-*tert*-butylbenzyl)-N'-methylguanidine;
N-(4-butoxyphenyl)-N,N'-bis(4-*tert*-butylbenzyl)-N,N'-dimethylguanidine;
5 and pharmaceutically acceptable salts of said compounds.

Particularly preferred R substituent groups of compounds of Formula IV, as that formula is defined above, include halo, alkyl, alkoxy, benzyloxy, aminoalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkenyl and alkynyl. Further
10 preferred is where R¹ is carbocyclic aryl, particularly substituted or unsubstituted phenyl, naphthyl or acenaphthyl. Alkyl including methyl, ethyl and propyl are preferred R² or R³ groups.

Specifically preferred compounds of Formula IV include:

15 N,N'-di-(4-*sec*-butylphenyl)guanidine;
N,N'-di-(4-*sec*-butylphenyl)-N-methylguanidine;
N,N'-di-(4-*sec*-butylphenyl)-N,N'-dimethylguanidine;
N-(2-naphthyl)-N'-(4-*isopropyl*phenyl)guanidine;
N-(2-naphthyl)-N'-(4-*isopropyl*phenyl)-N-methylguanidine;
20 N-(2-naphthyl)-N'-(4-*isopropyl*phenyl)-N'-methylguanidine;
N-(2-naphthyl)-N'-(4-*isopropyl*phenyl)-N,N'-dimethylguanidine;
N,N'-bis(4-*tert*-butylphenyl)guanidine;
N,N'-bis(4-*tert*-butylphenyl)-N-methylguanidine;
N,N'-bis(4-*tert*-butylphenyl)-N'-methylguanidine;
25 N,N'-bis(4-*tert*-butylphenyl)-N,N'-dimethylguanidine;
N-(4-*sec*-butylphenyl)-N'-(2,3,4-trichlorophenyl)guanidine;
N-(4-*sec*-butylphenyl)-N'-(2,3,4-trichlorophenyl)-N-methylguanidine;
N-(4-*sec*-butylphenyl)-N'-(2,3,4-trichlorophenyl)-N'-methylguanidine;
N-(4-*sec*-butylphenyl)-N'-(2,3,4-trichlorophenyl)-N,N'-dimethylguanidine;
30 N-(4-methoxy-1-naphthyl)-N'-(2,3,4-trichlorophenyl)guanidine;
N-(4-methoxy-1-naphthyl)-N'-(2,3,4-trichlorophenyl)-N-methylguanidine;

- N-(4-methoxy-1-naphthyl)-N'-(2,3,4-trichlorophenyl)-N'-methylguanidine;
N-(4-methoxy-1-naphthyl)-N'-(2,3,4-trichlorophenyl)-N,N'-dimethylguanidine;
N,N'-bis-(4-*sec*-butylphenyl)-2-iminopyrimidazolidine;
N,N'-bis(3-biphenyl)guanidine;
- 5 N,N'-bis(3-biphenyl)-N-methylguanidine;
N,N'-bis(3-biphenyl)-N'-methylguanidine;
N,N'-bis(3-biphenyl)-N,N'-dimethylguanidine;
N,N'-di-(3-*tert*-butylphenyl)guanidine;
N,N'-di-(3-*tert*-butylphenyl)-N-methylguanidine;
- 10 N,N'-di-(3-*tert*-butylphenyl)-N'-methylguanidine;
N,N'-di-(3-*tert*-butylphenyl)-N,N'-dimethylguanidine;
N,N'-bis-(4-methoxy-1-naphthyl)guanidine;
N,N'-bis-(4-methoxy-1-naphthyl)-N-methylguanidine;
N,N'-bis-(4-methoxy-1-naphthyl)-N'-methylguanidine;
- 15 N,N'-bis-(4-methoxy-1-naphthyl)-N,N'-dimethylguanidine;
N,N'-bis-(3-*sec*-butylphenyl)guanidine;
N,N'-bis-(3-*sec*-butylphenyl)-N-methylguanidine;
N,N'-bis-(3-*sec*-butylphenyl)-N'-methylguanidine;
N,N'-bis-(3-*sec*-butylphenyl)-N,N'-methylguanidine;
- 20 N,N'-bis(4-*n*-butylphenyl)guanidine;
N,N'-bis(4-*n*-butylphenyl)-N-methylguanidine;
N,N'-bis(4-*n*-butylphenyl)-N'-methylguanidine;
N,N'-bis(4-*n*-butylphenyl)-N,N'-dimethylguanidine;
N,N'-(*sec*-butylphenyl)-N'-(*n*-pentyl)guanidine;
- 25 N,N'-bis(3-benzyloxyphenyl)guanidine;
N,N'-bis(3-benzyloxyphenyl)-N-methylguanidine;
N,N'-bis(3-benzyloxyphenyl)-N,N'-dimethylguanidine;
N,N'-bis(4-benzyloxyphenyl)guanidine;
N,N'-bis(4-benzyloxyphenyl)-N-methylguanidine;
- 30 N,N'-bis(4-benzyloxyphenyl)-N,N'-dimethylguanidine;
N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)guanidine;

- N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N-methylguanidine;
N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-methylguanidine;
N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N,N'-dimethylguanidine;
N,N'-bis-(4-*tert*-butylphenyl)-2-iminopyrimidazolidine;
5 N,N'-bis-(4-pentylphenyl)-2-iminopyrimidazolidine;
N,N'-bis-(4-hexylphenyl)-2-iminopyrimidazolidine;
N,N'-bis-(naphthyl)-2-iminopyrimidazolidine;
N,N'-bis-(5-acenaphthyl)-2-iminopyrimidazolidine;
N,N'-bis-(tetralinyl)-2-iminopyrimidazolidine;
10 and pharmaceutically acceptable salts of said compounds.

Specifically preferred compounds of Formula V, as defined above, include:

- N-(5-acenaphthyl)-N'-(1,2,3,4-tetrahydroquinoliny)guanidine;
15 N-(5-acenaphthyl)-N'-(1,2,3,4-tetrahydroquinoliny)-N-methylguanidine;
N-(5-acenaphthyl)-N'-(1,2,3,4-tetrahydroquinoliny)-N'-methylguanidine;
N-(5-acenaphthyl)-N'-(1,2,3,4-tetrahydroquinoliny)-N,N'-dimethylguanidine;
N-(3-acenaphthyl)-N'-(indoliny)guanidine;
N-(3-acenaphthyl)-N'-(indoliny)-N-methylguanidine;
20 N-(3-acenaphthyl)-N'-(indoliny)-N'-methylguanidine;
N-(3-acenaphthyl)-N'-(indoliny)-N,N'-methylguanidine;
N-(5-acenaphthyl)-N'-(piperonyl)guanidine;
N-(5-acenaphthyl)-N'-(piperonyl)-N-methylguanidine;
N-(5-acenaphthyl)-N'-(piperonyl)-N'-methylguanidine;
25 N-(5-acenaphthyl)-N'-(piperonyl)-N,N'-dimethylguanidine;
and pharmaceutically acceptable salts of such compounds.

The invention also includes the following compounds, particularly for use in the methods of treatment disclosed herein:

- 30 N-(2-naphthyl)-N'-(2-adamantyl)guanidine;
N-(2-naphthyl)-N'-(2-adamantyl)-N-methylguanidine;

N-(2-naphthyl)-N'-(2-adamantyl)-N'-methylguanidine;
N-(2-naphthyl)-N'-(2-adamantyl)-N,N'-dimethylguanidine;
N,N'-bis-(5-indanyl)-guanidine;
N,N'-bis(6-benz[cd]indoliny-2[1H]-one)guanidine;

5 and pharmaceutically acceptable salts thereof.

Other specifically preferred compounds of the invention, including of the Formulas I-V above, include the following and are particularly preferred for use in the methods of treatment disclosed herein:

- 10 N-(3-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-*tert*-butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-pentoxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(5-acenaphthyl)-N-(4-benzyloxybenzyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-benzyloxybenzyl)guanidine;
15 N-(4-benzyloxyphenyl)-N-(4-benzyloxybenzyl)guanidine;
N-(5-acenaphthyl)-N-(3-benzyloxybenzyl)guanidine;
N-(4-isopropylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(4-benzyloxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(4-hexylphenyl)-N-(4-*tert*-hexylbenzyl)guanidine;
20 N-(4-*sec*-butylphenyl)-N-(4-*t*-butylbenzyl)-N'-pyrrolidinylguanidine;
N-(4-*sec*-butylphenyl)-N-(4-*t*-butylbenzyl)-N'-(4-thiomorpholinyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-piperidinylguanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-morpholinyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-
25 propylpiperidinyl)guanidine;
N-(4-butoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-piperidinyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-
benzylpiperidinyl)guanidine;
N-(4-benzyloxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-morpholinyl)guanidine;
30 N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(1,2,3,4-
tetrahydroisoquinolinyl)guanidine;

- N-(3-butoxy-4-methoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-morpholinyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(3,5-dimethyl-4-morpholinyl)guanidine;
- 5 N-(4-*tert*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-*sec*-butylphenyl)-N'-(methyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-*sec*-butylphenyl)-N'-(methyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(phenyl)guanidine;
- 10 N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-chlorophenyl)guanidine;
N-(4-butoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(phenyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(phenyl)-N'-methylguanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(3,4-dichlorophenyl)guanidine;
- 15 N-(4-hexylphenyl)-N-(4-*tert*-hexylbenzyl)-N'-phenylguanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-benzyloxyphenyl)guanidine;
N,N'-bis-(4-*tert*-butylphenyl)-N,N'-dimethylguanidine
N-(4-benzyloxyphenyl)-N'-(4-*tert*-butylphenyl)guanidine;
- 20 N,N'-bis-(3-(1'-methyl-2'-phenyl)ethyl)guanidine;
N-methyl-N-4-benzyloxyphenyl-N'-(4-*tert*-butylphenyl)guanidine;
N,N'-bis-(4-hexylphenyl)guanidine;
N-(3-(1-(4'-ethoxy)benzyl)phenethyl)-N'-(4-*tert*-butylphenyl)guanidine;
N-(4-benzyloxyphenyl)-N'-methyl-N-(4-*tert*-butylphenyl)guanidine;
- 25 N-(3-(4-*tert*-butylbenzyloxy)phenyl)-N'-(4-*tert*-butylphenyl)guanidine;
N-(3-(1'-benzylbutyl)phenyl)-N'-(4-*tert*-butylphenyl)guanidine;
N,N'-bis-(4-butylphenyl)-N-methylguanidine;
N,N'-bis-(4-*tert*-butylphenyl)-N,N'-dimethylguanidine;
N-(3-naphthaloxyphenyl)-N'-(4-*tert*-butylphenyl)guanidine;
- 30 N-(4-benzyloxyphenyl)-N'-(4-butylphenyl)guanidine;
N,N'-bis-(4-butylphenyl)-N-butylguanidine;

- N-3-(benzyloxymethyl)phenyl-N'-(4-*tert*-butylphenyl)guanidine;
N-(3,4-bis-butyloxyphenyl)-N'-(4-*tert*-butylphenyl)guanidine;
N-(3-benzyloxy)phenyl-N'-(4-*tert*-butylphenyl)guanidine;
N,N'-bis-(3-butoxy-4-methoxy)phenylguanidine;
5 N-(4-benzyloxyphenyl)-N-methyl-N'-(4-butylphenyl)guanidine;
N-(4-benzyloxyphenyl)-N'-methyl-N'-(4-butylphenyl)guanidine;
N,N'-bis-(6-tetralinyl)guanidine;
N-(6-tetralinyl)-N'-(4-*tert*-butylphenyl)guanidine;
N-(5-acenaphthyl)-N'-(6-benzothiazolyl)guanidine;
10 N-(5-acenaphthyl)-N'-(6-N-benzylindolinyl)guanidine;
N-(5-acenaphthyl)-N'-(4-benzo-2,1,3-thiadiazole)guanidine;
N-(5-acenaphthyl)-N'-[4-(6-methyl-benzothiazole)phenyl]guanidine;
N-(5-acenaphthyl)-N'-(1-benz[cd]indolinyl)guanidine;
N-(5-acenaphthyl)-N'-(6-benz[cd]indo-2[1H]-one)guanidine;
15 N-(4-butoxyphenyl)-N'-(4-chlorophenylethyl)guanidine;
N-(4-benzyloxyphenyl)-N,N'-diphenylguanidine;
N-(4-benzyloxyphenyl)-N'-benzyl-N'-phenylguanidine;
N-(3-benzyloxyphenyl)-N'-(4-thiobenzylphenyl)guanidine;
N,N'-bis(4-(phenylthio)phenyl)guanidine;
20 N,N'-bis(3-(phenylthio)phenyl)guanidine;
N-(5-acenaphthyl)-N'-(2-phenylethyl)guanidine;
N-(5-acenaphthyl)-N'-(3-butoxypropyl)guanidine;
N,N'-bis(2,2-diphenylethyl)guanidine;
N-(4-butoxyphenyl)-N'-(4-chlorophenylethyl)guanidine;
25 N-(4-butoxyphenyl)-N-(4-chlorobenzhydryl)guanidine;
(5-acenaphthyl)-N'-(phenethyl)-N'-benzylguanidine;
N-4-benzyloxyphenyl)-N'-(3-benzyloxyphenyl)-N'-(4-chlorobenzyl)guanidine;
N,N'-bis(4-benzyloxyphenyl)-N'-methylguanidine;
N-(4-benzyloxyphenyl)-N'-(3-benzyloxyphenyl)-N'-(4-chlorobenzyl)guanidine;
30 N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-phenylguanidine;
N-(4-*sec*-butylphenyl)-N'-(4-isopropoxyphenyl)-N'-phenylguanidine;

- N-(4-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-phenylguanidine;
N,N'-bis(3-octyloxyphenyl)guanidine;
N,N'-bis(4-butoxyphenyl)guanidine;
N,N'-bis(4-phenoxyphenyl)guanidine;
- 5 N-(3-benzyloxyphenyl)-N'-(4-phenoxyphenyl)guanidine;
N-(3-benzyloxyphenyl)-N'-(4-phenylazophenyl)guanidine;
N,N'-bis(3-benzyloxyphenyl)-N'-methylguanidine;
N-(4-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-methylguanidine;
N-(4-butoxyphenyl)-N'-(4-isopropoxyphenyl)guanidine;
- 10 N,N'-bis(4-(1-hydroxybutyl)phenyl)guanidine;
N-(4-butoxyphenyl)-N'-(3-methoxyphenyl)-N'-phenylguanidine;
N-(4-secbutylphenyl)-N'-phenyl-N'-(4-(2-isopropoxy)phenyl)guanidine;
N-(4-n-butoxyphenyl)-N'-(2-(4-chlorophenyl)ethyl)guanidine;
and pharmaceutically acceptable salts thereof.

15


- Compounds of the invention can be prepared by reaction of an amine, typically an amine salt such as an amine hydrochloride, with a preformed alkyl or aryl cyanamide (see S.R. Safer, et al., *J. Org. Chem.*, **13**:924 (1948)) or the corresponding N-substituted alkyl or aryl cyanamide. See
- 20 also G.J. Durant, et al., *J. Med. Chem.*, **28**:1414 (1985); C.A. Maryanoff, et al., *J. Org. Chem.*, **51**:1882 (1986); M.P. Kavanaugh, et al., *Proc. Natl. Acad. Sci. USA*, **85**:2844-2848 (1988); E. Weber, et al., *Proc. Natl. Acad. Sci. USA*, **83**:8784-8788 (1986); H.W.J. Cressman, *Org. Syn. Coll.*, **3**:608-609 (1955); International Applications WO 91/12797 and
- 25 PCT/US92/01050.

- More particularly, synthesis of N,N-disubstituted compounds of Formula I can be achieved by condensation of a disubstituted amine with cyanamide. For example, a disubstituted amine is prepared having the
- 30 desired substituents R and R¹ (as those substituents are defined in Formula I above), e.g., by condensation of a primary amine of the formula R-NH₂

with a compound of the formula R^1 wherein L is a leaving group and R and R^1 are as defined above for Formula I. Suitable reaction conditions can be readily determined based on the constituents employed. For example, a benzylhalide is suitably added to an arylamine at reduced temperature in the presence of a tertiary amine such as triethylamine and, after addition completion, the mixture is stirred at room temperature for about 15 hours. The resulting secondary amine can be purified by conventional means such as chromatography and then reacted with a suitable acid such as methanesulfonic acid to form the amine salt. The amine salt is reacted with a large molar excess of cyanamide in a suitable solvent such as methanol for a time and temperature sufficient to form the N,N-disubstituted guanidine.


Synthesis of symmetrical N,N'-disubstituted guanidines of the invention can be typically achieved by directly reacting two equivalents of the amine with one equivalent of cyanogen bromide in suitable solvent such as ethanol as depicted in "Method A" of Scheme 1 below. Unsymmetrical N,N'-disubstituted guanidines can be prepared by reacting a substituted cyanamide such as aryl cyanamides with the appropriate amine hydrohalide salts in suitable solvent such as refluxing chlorobenzene or toluene as depicted in "Method B" of Scheme 1 below. The requisite cyanamides can be synthesized from the corresponding amines by treatment with cyanogen bromide in suitable solvent such as ether.

SCHEME I



N,N,N'-tri-substituted or N,N,N',N'-tetra-substituted guanidines of the invention can be synthesized (Scheme II below, Method C and D respectively) by reacting a substituted cyanamide such as a N-alkyl-N-arylcyanoamide with an appropriate amine hydrohalide salt in a suitable solvent such as refluxing chlorobenzene or toluene. The starting cyanamides can be synthesized, e.g., by an alkylation of an arylcyanoamide with sodium hydride/alkyl halide in suitable solvent such as tetrahydrofuran.

SCHEME II



Appropriate substituted amine and cyanamide reagents are available or can be prepared by recognized procedures. A nitro acenaphthyl derivative having one or more additional ring substituents can be prepared as described by M.D. Varney, et al., *J. Med. Chem.*, **35**:671 (1992). Such a substituted nitro acenaphthyl derivative can be reduced to the corresponding amine by hydrogenation, and the amine reacted with BrCN as discussed above. For preparation of other acenaphthyl derivatives having an amine or amine precursor group and one or more additional ring substituents, see V.N. Komissarov, *Zh. Org. Khim.*, **26**(5): 1106-10 (1990); L. Skulski, et al., *Pol. J. Chem.*, **55**(9): 1809-24 (1981); A.F. Pozharskii, *Isobret. Prom. Obraztsy, Tovarnye Znaki*, (3), 96-7 (1982); J.P. Li, et al., US 78-890736 (1978); N.S. Vorozhtsov, *Zh. Org. Khim.*, **8**(2): 353-7 (1972); J. Wolinski et al., *Rocz. Chem.*, **44**(9): 1721-31 (1970); A.P. Karishin, et al., *Zh. Obshch. Khim.*, **39**(9): 2098-101 (1969); and V.V. Mezheritskii, et al., *Zh. Org. Khim.*, **27**(10): 2198-204 (1991).

Compounds of Formula IV where R² and R³ taken together form a substituted or unsubstituted alkylene linkage of from 2 to about 6 carbon atoms can be prepared as exemplified by the procedure disclosed in Example 8 which follows. Thus, as clearly understood by those skilled in the synthesis arts, an appropriate N-substituted diaminealkylene is suitably reacted with cyanogen bromide in an appropriate solvent such as an alcohol and the reaction conducted at a temperature and for a time sufficient to carry out the reaction. The cyclic reaction product can be suitably purified by conventional techniques if desired, such as by chromatography.

As discussed above, the substituted guanidines of the invention are useful for a number of therapeutic applications, including treatment of those diseases that result from modulation of a particular neurotransmitter system that can be counteracted by one or more of the substituted guanidines of the invention. As mentioned above, modulation of neurotransmitter release involves either the inhibition of neurotransmitter release, the potentiation of neurotransmitter release, or the increase or decrease of the time course of neurotransmitter release from neuronal tissue. Neurotransmitters which may be modulated by compounds of the invention include, but are not limited to those neurotransmitters identified above. One of ordinary skill in the art can select those compounds which are effective or particularly effective modulators of neurotransmitter release using the procedures disclosed herein, or in the literature such as PCT/US92/01050, with no more than routine experimentation. For example, compounds for the prevention of neuronal death in brain ischemia can be evaluated *in vivo* in one or more variations of the rat middle cerebral artery occlusion model. Such models are generally considered to be particularly predictive of neuroprotective efficacy in stroke [Ginsberg, et al., *Stroke*, 20:1627-1642 (1989)]. Efficacy of compounds of the invention also may be assessed in the 4-vessel occlusion model of global ischemia [Pulsinelli, et al., *Stroke*:19:913-941 (1988)].

In particular, the invention provides methods for treatment and/or prophylaxis of neurological conditions such as epilepsy, neurodegenerative conditions and/or nerve cell death resulting from, e.g., hypoxia, hypoglycemia, brain or spinal chord ischemia, brain or spinal chord trauma, stroke, heart attack or drowning. Typical candidates for treatment include heart attack, stroke, brain or spinal cord injury patients, patients undergoing major surgery where brain ischemia is a potential complication and patients such as divers suffering from decompression sickness due to gas emboli in the blood stream.

The invention also provides methods to treat and/or prevent various neurodegenerative diseases of a subject such as an animal, particularly a human, by administering a therapeutically effective dose of one or more compounds of the invention. Typical neurodegenerative diseases that can be treated and/or prevented include Parkinson's disease, Huntington's disease, Amyotrophic Lateral Sclerosis, Alzheimer's disease, Down's Syndrome, Korsakoff's disease, olivopontocerebellar atrophy, HIV-induced dementia and blindness, multi-infarct dementia or diabetic neuropathy. As disclosed by Dreyer et al., Science, 248:364-367 (1990), gp120 neurotoxicity is associated with increased levels of Ca^{2+} which are apparently mediated by Ca channels and blocked by dihydropyridine Ca channel antagonists. Though again not wishing to be bound by theory, compounds of the invention should have utility in treating HIV-induced dementia and blindness by means of preventing the release of excessive glutamate.

As noted above the invention provides methods of treating Korsakoff's disease, a chronic alcoholism-induced condition, comprising administering to a subject including a mammal, particularly a human, one or more compounds of the invention in an amount effective to treat the disease. Pretreatment of animals with the NMDA antagonist MK-801

(Merck Index, monograph 3392, 11th ed., 1989) markedly attenuates the extent of cell loss, hemorrhages and amino acid changes in a rat model of Korsakoff's disease. See P.J. Langlais, et al., *Soc. Neurosci. Abstr.*, 14:774 (1988). Therefore, compounds of the invention have utility for the
5 attenuation of cell loss, hemorrhages and amino acid changes associated with Korsakoff's disease.

At least some compounds of the invention will have utility in treating or preventing conditions treatable by the blockage of voltage-activated
10 sodium ion-channels. Accordingly, the invention provides methods for blockage of voltage-activated sodium channels of neuronal cells, particularly mammalian cells such as human neuronal cells, comprising the administration to the cells of an effective amount of a compound of the invention, particularly by such administration to a mammal in need of such
15 treatment. Conditions that can be treated by blockage of sodium channels will include, e.g., epilepsy. The invention also provides methods of for blockage of sodium channels of mammalian smooth or skeletal muscle cells, comprising administering to such cells an effective amount of one or more a compounds of the invention. Such methods will also be useful, e.g., for
20 therapy of a mammal such as a human having or susceptible to paramyotonia or hyperkalemic periodic paralysis [See Cannon, S.C. et al., *Neuron*, 10:317-326 (1993)].

Moreover, some compounds of the invention will block both sodium
25 channels as well as presynaptic calcium channels. This dual action may be particularly desirable for neuroprotective therapies [Kucharczyk, J. et al., *Radiology*, 179:221-227 (1991)].

It has been reported that NMDA antagonists which do not cross the
30 blood/brain barrier may be used to alleviate certain undesirable side effects of cancer chemotherapy, e.g. nausea and emesis [A. Fink-Jensen et al.,

Neurosci. Lett., 137(2):173 (1992)]. See also Price, M.T., et al., *Soc. NeuroSci. Abstr.*, 16:377, abstr. 161.16 (1990). These actions of NMDA antagonists are presumably mediated by blockade of the postsynaptic activity of glutamate released from neurons of the peripheral nervous system. Again without wishing to be bound by theory, this indicates that compounds which block the release of glutamate will be useful for this therapeutic indication. Compounds of the invention, particularly those compounds that are charged such as in the form of a pharmaceutically acceptable salt, and those compounds that are otherwise hydrophilic such as compounds that comprise one or more polar functionalities e.g. carboxy, amino, hydroxy and the like, may have comparatively limited ability to cross the blood brain barrier. It is thus believed that compounds of the invention, especially charged or otherwise hydrophilic compounds of the invention with limited blood brain barrier permeability, will be clinically useful to ameliorate the side effects associated with chemotherapy, particularly cancer chemotherapy, that may be experienced by a mammal, particularly a human receiving such chemotherapy. The compound of the invention would be typically administered to the subject in coordination with the chemotherapy regime.

Compounds of the invention may be used in therapy in conjunction with other medicaments. For example, for treatment of a stroke victim, one or more compounds of the invention may be suitably administered together with a pharmaceutical targeted for interaction in the blood clotting mechanism such as streptokinase, TPA and urokinase. See VonKummer, R. et al., *Stroke*, 23:646-652 (1992); Sereghy, T. et al., *Stroke*, 24:1702-1708 (1993).

Compounds of the invention will be useful for treatment of secretory disorders, particularly in view of the demonstrated ability of the compounds to block Ca channels which are identical or closely related to those which

regulate secretion. The invention thus includes methods for blocking voltage-activated calcium channels of mammalian secretory cells which comprises administering to such cells a blockage-effective amount of a compound of the invention. The invention further provides methods for
5 treatment of a disease in which the pathophysiology of the disorder involves inappropriate or excessive cellular secretion of a catecholamine, a growth factor or precursor thereof (including those growth factors specifically discussed *infra*), a hormone or precursor thereof (including those hormones specifically discussed *infra*) or a member of the neuregulin family of proteins
10 including glial growth factors, the heregulins and the neu differentiation factors. Such a method will be particularly useful for treating a mammal such as a human suffering from or susceptible to hypersecretory disorders discussed below.

15 More particularly, compounds of the invention could be used in treatment of a hypersecretory disorder such as pheochromocytoma, which is a disorder resulting from the presence of a tumor of the chromaffin cells in the adrenal medulla [Bravo, E.L. and Gifford, R.W. (1984) *New Eng. J. Med.* **311**: 1298-1300]. This disorder is characterized by the
20 hypersecretion of catecholamines, resulting in hypertension which may be paroxysmal and associated with attacks of palpitation, headache, nausea, breathing difficulty, and anxiety. Compound(s) of the invention also could be used in treatment of pancreatitis, which is an inflammation of the pancreas leading to hypersecretion of hormones and enzymes from the
25 acinar cells of the pancreas, among them hormones such as vasoactive intestinal peptide (VIP) and insulin; digestive enzymes and their inactive precursors, among them lipases and proteases, deoxyribonucleases, ribonucleases, and amylase [Greenberger, N.J. et al. *Harrison's Principles of Internal Medicine*, 11th Ed., New York, McGraw-Hill, pp.1372-1380
30 (1987)]. In severe cases of pancreatitis, autodigestion of the pancreas by the hypersecretion and subsequent activation of said digestive enzymes can

be fatal [Greenberger, N.J. et al., *ibid*]. For these and other indications mediated by hypersecretory activity outside the central nervous system, compounds of the invention, particularly those which are charged and/or hydrophilic or otherwise have limited blood/brain barrier permeability should be clinically useful upon systemic and/or local administration.

Certain hypersecretory disorders may result from abnormal activity of cells within the central nervous system, among them cells of the pituitary gland, also termed the hypophysis, located at the base of the brain.

Secretion of hormones and related substances from cells of the adenohipophysis is regulated by releasing factors, primarily those secreted by the hypothalamus [Cooper, P.E. et al., *Diseases of the Nervous System: Clinical Neurobiology*, eds, Saunders, Philadelphia, pp. 567-583 (1992)]. Substances secreted by the adenohipophysis include growth hormone, prolactin, thyroid stimulating hormone (TSH), and adrenocorticotrophic hormone (ACTH). Hypersecretion of these substances from the pituitary can lead to a variety of disorders of growth (e.g. acromegaly due to hypersecretion of growth hormone) and metabolism (e.g. secondary hyperthyroidism triggered by hypersecretion of TSH, and Cushing's disease, which results from excessive secretion by the pituitary of precursor peptides containing ACTH) [see Cooper, P.E. et al. *ibid.*]. These disorders are often due to benign tumors of the pituitary secretory cells. Compounds of the invention, particularly those that are relatively hydrophobic and/or by some means penetrate the blood/brain barrier, should have utility for treatment of such disorders by suitable administration to a subject, particularly a human. In some instances pharmacotherapy with compounds of the invention may obviate the necessity and attendant risk of neurosurgery performed for the purpose of removing such benign tumors. As referred to above, hydrophobic compounds of the invention would include those compounds that do not comprise highly polar moieties such as carboxy and the like.

Compounds of the invention also may be used in treatment of disorders involving hypersecretion of substances produced by the hypothalamus such as diabetes insipidus, which may be caused by hypersensitivity to, or excessive release of, AVP. AVP is a peptide synthesized in and released from neurons of the supraoptic and paraventricular nuclei of the hypothalamus [see Copper, P.E. et al., *Diseases of the Nervous System: Clinical Neurobiology*, eds, Saunders, Philadelphia, pp. 567-583 (1992)]. A current means of treatment of diabetes insipidus is surgical destruction of most of the cells in the supraoptic nucleus. Pharmacotherapy with compounds of the invention could in at least some instance obviate the need for such neurosurgery.

The pituitary has been indicated to secrete growth factors. Evidence shows a family of proteins, termed glial growth factors ("GGF"s), to be a group of such growth factors secreted by the pituitary. GGF's are mitogenic for myelin-forming Schwann cells, and as such may play an important role in development and regeneration of the nervous system [Marchionni et al., *Nature*, **362**:312-318 (1993)]. Bovine pituitary glands have been identified as an enriched source of GGF's, and a GGF of M_r 31,000 has been purified from bovine pituitary [Lemke, G.E. et al., *J. Neurosci.*, **4**:75-83 (1984)]. Multiple molecular forms of GGF may be secreted from the pituitary, either in active form or as precursors. GGF's from bovine pituitary extracts can be resolved into at least 3 activities with different molecular masses: GGF-I (34,000), GGFII (59,000), and GGFIII (45,000) [Goodearl et al., *J. Biol Chem.*, **268**:18095-18102 (1993)]. GGF's are structurally related to members of a family of proteins which are known to activate the p185^{erbB2} receptor kinase, including the heregulins [Holmes, W.E. et al., *Science*, **256**:1205-1210 (1992)], and neu differentiation factor [Wen, D. et al., *Cell*, **69**:559-572 (1992)].

While the precise role of GGF's and the aforementioned related proteins in the development, maintenance, and/or repair of the nervous system and muscle has yet to be elucidated, and the mechanism of secretion of GGF's and related molecules has yet to be defined, existing evidence indicates that pathophysiological circumstances may arise in which it may be desirable to regulate the secretion of GGF's and related proteins from the pituitary and/or other secretory sites. One such circumstance may be diseases that involve the deterioration of nerve, for example diabetic neuropathy [Duchen, L.W. (1983) in *Autonomic Failure: A Textbook of Clinical Disorders of the Nervous System*, Bannister, R., ed., N.Y., Oxford Univ. Press; Foster, D.W. (1987) in *Harrison's Principles of Internal Medicine, 11th Ed.*, New York, McGraw-Hill, pp. 1788-1795], or the deterioration of muscle, among them muscular dystrophies [Brooke, M.H. (1985) *A Clinician's View of Neuromuscular Disease*, 2nd ed., Baltimore, Williams and Wilkins; Huges, S. M., and Blau, H.M. (1990) *Nature* **345**: 350-352]. Compounds of the invention should have therapeutic utility in treating such disorders involving the deterioration of nerve or muscle. Without wishing to be bound by theory, it is believed that compounds of the invention will have utility in treating such disorders by modulating the exocytosis from the cells in which they are synthesized of GGF's, other members of the neuregulin family, and other factors secreted into the blood which are involved in development, maintenance, or repair of nerve and/or muscle.

The aforementioned ability of compounds of the invention to block Ca channels which are identical or closely related to those which regulate cardiovascular function, as demonstrated in the examples which follow, indicate that compounds of the invention will find utility in therapy of cardiovascular disorders. Among the disorders currently known to be treatable by inhibitors of L-type Ca channels such as verapamil, diltiazem, and nifedipine are hypertension, angina pectoris, cardiac arrhythmias. As

shown in Example 148, a subset of compounds of the invention show equal or greater potency for block of L-type Ca channels when compared with the ability of verapamil or diltiazem to block said channels using the same assay protocol.

5

10 The aforementioned ability of compounds of the invention to block Na channels which are closely related to those which regulate cardiovascular function, as shown in Example 149, *infra*, indicate that compounds of the invention will find utility in therapy of cardiovascular disorders treatable by blockers of Na channels. The invention thus includes methods for blocking voltage-activated sodium channels of mammalian cardiac cells comprising administration to such cells a blockage-effective amount of one or more compounds of the invention. A major indication for such Na channel blockers is cardiac arrhythmias, which are currently treated by blockers of Na channels, among them quinidine, procainamide, lidocaine, and 15 diphenylhydantoin (phenytoin). Among the cardiac arrhythmias successfully treatable by said Na channel blockers are ventricular tachycardia; ventricular premature depolarizations; digitalis-induced atrial tachycardia and atrial and ventricular arrhythmias; paroxysmal supraventricular tachycardia; atrial 20 fibrillation; and prophylaxis against the development of supraventricular arrhythmias (see Bigger, J.T. et al., *The Pharmacological Basis of Therapeutics, 7th Ed.*, eds., New York, MacMillan, pp. 748-783 (1985)). Accordingly, compounds of the invention should find utility in treatment of cardiac arrhythmias treatable by blockers of cardiac Na channels.

25 Compounds of the invention should find utility in treatment of hypertension and/or angina pectoris treatable by blockers of cardiac Na channels. For these and other indications treatable by blocking cardiac Na channel activity, compounds of the invention, particularly those which are charged and/or hydrophilic and otherwise do not cross the blood/brain barrier, are 30 believed to be clinically useful upon systemic and/or local administration.

Some cardiac arrhythmias, among them paroxysmal supraventricular tachycardia and other supraventricular arrhythmias, are treatable by both blockers of cardiovascular L-type channels and by blockers of cardiac Na channels. Compounds of the invention with dual actions against Na
5 channels and Ca channels (such as N-(5-acenaphthyl)-N-(4-iso-propylbenzyl) guanidine and N-(4-methoxynaphthyl)-N'-(2,3,4-trichlorophenyl) guanidine), should thus have particular utility in treatment of said arrhythmias.

As discussed above, compounds of the invention also will be useful
10 for treatment of chronic pain and as a local anesthetic.

The compounds of this invention can be administered to a subject such as a human intranasally, orally or by injection, e.g., intramuscular, intraperitoneal, subcutaneous or intravenous injection, or by transdermal,
15 intraocular or enteral means. The optimal dose can be determined by conventional means including the assays described in the examples which follow. Guanidines of the invention are suitably administered to a subject in the protonated and water-soluble form, e.g., as a pharmaceutically acceptable salt of an organic or inorganic acid, e.g., hydrochloride,
20 hydrobromide, sulfate, hemi-sulfate, mesylate, gluconate, phosphate, nitrate, acetate, oxalate, citrate, maleate, etc., prepared by procedures such as those disclosed in the examples which follow.

The compounds of this invention can be employed, either alone or in
25 combination with one or more other therapeutic agents as discussed above, as a pharmaceutical composition in mixture with conventional excipient, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral or intranasal application which do not deleteriously react with the active compounds and are not deleterious to the recipient
30 thereof. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohol, vegetable oils, polyethylene glycols,

gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously react with the active compounds.

10 For parenteral application, particularly suitable are solutions, preferably oily or aqueous solutions as well as suspensions, emulsions, or implants, including suppositories. Ampules are convenient unit dosages.

For enteral application, particularly suitable are tablets, dragees or capsules having talc and/or carbohydrate carrier binder or the like, the carrier preferably being lactose and/or corn starch and/or potato starch. A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including those wherein the active component is protected with differentially degradable coatings, e.g., by microencapsulation, multiple coatings, etc.

Intravenous or parenteral administration, e.g., sub-cutaneous, intraperitoneal or intramuscular administration are generally preferred.

25 Instances may arise in which site-specific drug-delivery methods would constitute a preferred method of delivering compounds of the invention to the tissue in need of therapy [see Tomlinson, E., *Advanced Drug Delivery Reviews*, 1:87-198 (1987)]. For example, in the case of disorders of muscle function originating from a pathophysiological condition of Na channels of skeletal muscle, among them hyperkalemic periodic paralysis, it may be desirable to microencapsulate compounds of the

invention within delivery vehicles such as liposomes [Yagi, K., *Medical Applications of Liposomes*, Japan Soc. Press, Tokyo (1986); Gregoriadis, G., ed. *Liposome Technology*, Vol. I-III, CRC press, Inc., Cleveland (1984)], said liposomes containing a monoclonal antibody targeted to specific
5 antigens on or near the surface of the diseased muscle cells. Said method of drug delivery should result in selective binding of the liposomes to the target tissue, and release of the compound of the invention near the abnormally functioning skeletal muscle Na channels, where said compound will inhibit the persistent activation of muscle Na channels which
10 constitutes the molecular abnormality underlying the disease.

A targeted delivery method of one or more compounds of the invention also may be preferred for treatment of pheochromocytoma or another abnormality which results in hypersecretion of catecholamines into
15 the blood. Because the Ca channels of chromaffin cells are closely related to those of nerve, cardiac cells, and muscle (Neher, E. et al., *Neuron*, 10:21-30 (1993); Bean, B.P. *Ann. Rev. Physiol.*, 51:367-384 (1989); Hess, P., *Ann. Rev. Neurosci.*, 13:337-56 (1990)), systemic administration of a compound of the invention at concentrations sufficient to block release of
20 catecholamines from chromaffin cells may produce certain side effects resulting from block of, for example, neuronal and cardiovascular Ca channels. Accordingly, delivery of compounds of the invention to chromaffin cells may be enhanced and said side effects reduced by their incorporation into liposomes containing a monoclonal antibody targeted to
25 specific antigens on or near the surface of the chromaffin cells of the hypersecreting adrenal medulla.

This method of liposome-mediated drug targeting has been reported for delivery of a variety of agents, to be used for indications such as cancer
30 chemotherapy and destruction of tumors [e.g., Bassett, J.B. et al. *J. Urol.*, 135:612-615 (1986)]. Suitable refinements of that method as well as new

methods for site-specific drug delivery will be useful for administration of compounds of the invention to a subject. In particular, appropriate methods for site-specific delivery of compounds of the invention may include incorporation of said compounds into polymer beads which afford slow site-specific release [Mathiowitz, E. et al., *J. Controlled Release*, 5:13-18 (1987)], and delivery to the target tissue by means of surgically implanted pumps.

It will be appreciated that the actual preferred amounts of active compounds used in a given therapy will vary according to the specific compound being utilized, the particular compositions formulated, the mode of application, the particular site of administration, etc. Optimal administration rates for a given protocol of administration can be readily ascertained by those skilled in the art using conventional dosage determination tests conducted with regard to the foregoing guidelines. In general, a suitable effective dose of one or more compounds of the invention, particularly when using the more potent compound(s) of the invention, will be in the range of from 0.5 to 500 milligrams per kilogram bodyweight of recipient per day, preferably in the range of 1 to 100 milligrams per kilogram bodyweight of recipient per day. The desired dose is suitably administered once daily, or in several sub-doses, e.g., 2 to 4 sub-doses, are administered at appropriate intervals through the day, or other appropriate schedule. Such sub-doses may be administered as unit dosage forms, e.g., containing from 0.25 to 25 milligrams of compound(s) of the invention per unit dosage, preferably from 0.5 to 5 milligrams per unit dosage.

Alternatively, compounds of the invention may be administered continuously for a period of time, for example by an intravenous infusion or by means of a suitably placed transdermal patch incorporating and releasing compounds of the invention.

As disclosed above, suitably labeled compounds of the invention can be used to detect ion channel (e.g., Ca or Na) activity, which will serve to diagnosis certain human diseases as discussed herein. A compound of the invention may be suitably radiolabeled, e.g. with ^{125}I such as on an aryl ring of the compound, and the labeled compound administered to a subject and the subject then scanned for binding of the compound to ion channels using an appropriate scanning tool. For example, single photon emission computed topography ("SPECT") may be suitably employed to detected such binding. Suitable radiolabeled compounds of the invention may be prepared by known procedures. For example, a compound of the invention having an aromatic group, such as phenyl, that has a bromo or chloro ring substituent can be employed in an exchange labeling reaction to provide the corresponding compound having an ^{125}I ring substituent.

As with prior guanidines such as those reported in U.S. Patent No. 1,411,713, the guanidines of the present invention should have utility as rubber accelerators.

All documents mentioned herein are incorporated herein by reference in their entirety.

The present invention will be further illustrated with reference to the following examples which aid in the understanding of the present invention, but which are not to be construed as limitations thereof.

GENERAL COMMENTS

In the following examples, all percentages reported herein, unless otherwise specified, are percent by weight. All temperatures are expressed in degrees Celsius.

Melting points were determined in open capillary tubes on a Thomas-Hoover apparatus and are uncorrected. Thin-layer chromatography was performed on Baker-flex 1B2-F silica gel plates. Guanidines were visualized on TLC with 254-nM UV light or as a blue spot with bromocresol spray reagent (Sigma Chemical Co.). Preparative TLC was performed on Analtech GF precoated silica gel (1000 μ m) glass-backed plates (20 x 20 cm). The IR, ^1H and ^{13}C NMR spectra of all compounds were consistent with their assigned structures. NMR spectra were recorded on Varian Gemini 300 and the chemical shifts were reported in ppm (δ) relative to the residual signal of the deuterated solvent (CHCl_3 , δ 7.26; CHD_2OD , δ 3.30). Infrared spectra were recorded in CHCl_3 (unless otherwise noted) on Perkin-Elmer model 1420. All new compounds were analyzed either for C, H, and N elemental analyses or for exact mass. Elemental analyses were performed by either Galbraith Laboratories (Knoxville, TN) or MHW Laboratories (Tucson, AZ). High Resolution Mass spectra (HRMS) were recorded on a Finnegan MAT 90. HPLC were performed on a C18 reverse phase column using 50:50 water:acetonitrile with 0.1% TFA as the mobile phase. BrCN was obtained from Aldrich Chemical Co., and was used as received. All starting amines were obtained from commercial sources and were purified by standard procedures before use, or they were prepared, where noted, by published procedures. Chlorobenzene, ether (Et_2O) and tetrahydrofuran (THF) were anhydrous quality solvents (Sure Seal) supplied by Aldrich. All other solvents were reagent grade. Alkyl- and arylcyanamides were prepared as described above and according to published procedures (e.g., PCT/US92/01050) by reaction of the amines with BrCN in ether.

Example 1: Preparation of N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

Part 1: Preparation of N-(4-*sec*-Butylphenyl)-4-*tert*-butylbenzylamine

5 A mixture of 4-*sec*-butylaniline (2.89 g, 20 mmol) and triethylamine (2.5 g, 25 mmol) in toluene (100 mL) was stirred at 4°C for 15 hours and precipitates formed. The precipitates (triethylamine•HBr) were filtered out; the filtrate was concentrated to dryness. Then the crude reaction mixture was purified by column chromatography (SiO₂, hexane/CH₂Cl₂ = 5/1). N-(4-*sec*-butylphenyl)-4-*tert*-butylbenzylamine (4.5 g, an oil) was obtained.

Part 2: Preparation of N-(4-*sec*-Butylphenyl)-4-*tert*-butylbenzylamine•HCl

15 To a solution of N-(4-*sec*-butylphenyl)-4-*tert*-butylbenzylamine (4.5 g) in diethylether (10 mL) was added ether at HCl solution at 4°C, then the reaction mixture was stirred at 25°C for 10 minutes. The resulting solution was the evaporated and dried under vacuum to afford 4.7 g of N-(4-*sec*-butylphenyl)-4-*tert*-butylbenzylamine•HCl.

Part 3: Preparation of N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

20 A mixture of N-(4-*sec*-butylphenyl)-4-*tert*-butylbenzylamine•HCl (0.8 g, 2.7 mmol) and cyanamide (2 g) in methanol was heated at 70°C for 40 hours. During the 40 hour period, another two portions of cyanamide (0.5 g, each time) were added. The reaction did not go to completion; a small percentage of N-(4-*sec*-butylphenyl)-4-*tert*-butylbenzylamine•HCl remained in the mixture. The crude product was purified by column chromatography (SiO₂, CH₂Cl₂/MeOH = 9/1). Then the purified compound (a mixture of the guanidine and some cyanamide) dissolved in water (20 mL) was basified to pH 14. The guanidine•free base was extracted with CH₂Cl₂ (20 mL, two times) and the combined extracts were concentrated. Finally, the pure N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine was converted into its HCl salt by methanolic HCl treatment. After drying under vacuum for 15

hours, the pure product (0.6 g) was obtained as a white solid, mp: 177-178°C; TLC: R_f = 0.4 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CD_3OD): δ ppm 7.39-7.12 (m, ArH, 8H), 4.90 (s, CH_2 , 1H), 4.88 (s, CH_2 , 1H), 2.63 (m, CH, 1H), 1.57 (m, CH_2 , 2H), 1.29 (s, CH_3 , 9H), 1.21 (d, CH_3 , 3H), 0.80 (t, CH_3 , 3H); HRMS: 337.2524 (337.2518 Calcd. for $\text{C}_{22}\text{H}_{31}\text{N}_3$); HPLC: 99% pure.

Example 2: Preparation of N-(5-Acenaphthyl)-N-(4-*tert*-butylbenzyl)guanidine

Part 1: Preparation of 5-Acenaphthylamine

A mixture of 5 and 3-nitroacenaphthene was reduced with Pd/C in ethyl acetate under hydrogen at 40 psi pressure and the resulting amines were separated by recrystallization from cyclohexane/ethyl acetate.

Part 2: Preparation of N-(5-Acenaphthyl)-4-*tert*-butylbenzylamine

A mixture of 5-acenaphthylamine (1.0 g, 6 mmol) and triethylamine (0.76 g, 7.5 mmol) in toluene (50 mL) was stirred at 4°C, and 4-*tert*-butylbenzyl-bromide (1.36 g, 6 mmol) was added in slowly. The reaction mixture was stirred at 23°C for 15 hours and precipitates formed. The precipitates (triethylamine•HBr) were filtered out; the filtrate was concentrated to dryness. Then the crude reaction mixture was purified by column chromatography (SiO_2 , hexane/ CH_2Cl_2 = 5/1). N-(5-acenaphthyl)-4-*tert*-butylbenzylamine (1.67 g, an oil) was obtained.

Part 3: Preparation of N-(5-Acenaphthyl)-4-*tert*-butylbenzylamine•mesylate

To a solution of N-(5-acenaphthyl)-4-*tert*-butylbenzylamine (1.1 g) in diethylether (10 mL) was added methanesulfonic acid (0.8 g) at 4°C, then the reaction mixture was stirred at 25°C for 10 minutes. The resulting solution was then evaporated and dried under vacuum to afford N-(5-acenaphthyl)-4-*tert*-butylbenzylamine•mesylate.

Part 4: Preparation of N-(5-Acenaphthyl)-N-(4-*tert*-butylbenzyl)guanidine

A mixture of N-(5-acenaphthyl)-4-*tert*-butylbenzylamine • mesylate (3.5 mmol) and cyanamide (0.4 g) in methanol was heated at 70°C for 40 hours. During the 40 hour period, another two portions of cyanamide (0.5 g, each time) were added. The reaction did not go to completion; a small percentage of N-(5-acenaphthyl)-4-*tert*-butylbenzylamine•HCl remained in the mixture. The crude product was purified by column chromatography (SiO₂, MeOH/CH₂Cl₂ = 0 to 10%). The purified compound (a mixture of the guanidine and some cyanamide) dissolved in water (20 mL) was basified to pH 14. Then the guanidine•free base was extracted with CH₂Cl₂ (20 mL, two times) and the combined extracts were concentrated. Finally, the pure N-(5-acenaphthyl)-N-(4-*tert*-butylbenzyl)guanidine was dried under vacuum for 15 hours to afford the pure product (0.6 g) as a white solid, mp: 85-86°C; TLC: R_f=0.4 (SiO₂, CH₂Cl₂/MeOH = 9/1); ¹H NMR (CD₃OD): δ ppm 7.60-7.20 (m, ArH, 9H), 4.90 (s, CH₂, 2H), 3.41-3.31 (m, CH₂, 4H), 1.26 (s, CH₃, 9H); MS(EI): m/e 357.2 (M⁺: C₂₄H₂₇N₃•1/4 H₂O); Calcd. (%): C, 79.62, H, 7.66, N, 11.60; Found (%): C, 79.47; H, 7.48, N, 11.79.

Example 3: Preparation of N-(5-Acenaphthyl)-N'-benzhydrylguanidine•HCl

Step 1. 5-Acenaphthyl cyanamide

5-aminoacenaphthene (7.0 g, 41.4 mmol) was dissolved in a mixture of ether (100 mL) and ethyl acetate (25 mL). To this solution was added 5.2 mL of a 5 M solution of cyanogen bromide in acetonitrile (25.6 mmol of cyanogen bromide). The solution was stirred overnight, with the gradual appearance of gray precipitate. The solid was removed by filtration (the hydrobromide of 5-aminoacenaphthene) and the resulting filtrate concentrated in vacuo to afford a semi-solid residue. Ether (60 mL) was added to the residue and the mixture was stirred overnight. The solid was removed (more hydrobromide of 5-aminoacenaphthene) and the filtrate concentrate to approximately 20 mL and then diluted with warm

cyclohexane (15 mL). Upon standing at room temperature, off-white crystals were deposited. They were collected, washed with cyclohexane-ether (1:1) and dried in vacuo to give 1.5 g of pure product, mp: 163-65°C.

5 **Step 2. Preparation of N-(5-Acenaphthyl)-N'-benzhydrylguanidine•HCl**

 A mixture of 5-acenaphthyl cyanamide (0.194 g, 1 mmol) and benzhydryl amine hydrochloride (0.209 g, 0.95 mmol; prepared from benzhydryl amine and 1.0 N HCl-ether) were heated at reflux in 10 mL of chlorobenzene. After 6 hours reflux, the mixture turned into a clear solution and the reflux continued another 12 hours. The mixture was cooled to 20°C and concentrated on a rotavapor to give a brown syrup. This syrup was treated with norite-A in a boiling ethanol for 20 minutes to give a colorless glass. Upon stirring this glass in anhydrous ether for 10 hours at room temperature resulted in a bright white solid. The solid was collected by filtration and washed with excess of ether and dried in vacuo at 40°C to give product (0.326 g, 79%) as a white solid, mp: 225-27°C; TLC: R_f =0.4 (CH₂Cl₂:MeOH; 9:1); ¹H NMR (CD₃OD): δ 7.47-7.33 (m, 15H, ArH), 6.14 (s, 1H), 3.41 (bs, 4H, 2 X -CH₂); MS(EI): m/e 378 (M⁺1).

20 **Example 4: Preparation of N,N'-bis(4-sec-Butylphenyl)guanidine•HCl**

 Cyanogen bromide (0.390 g, 3.7 mmol) was added in portions to a stirred 4-sec-butyl aniline (0.746 g, 5 mmol) at room temperature. Ethanol (absolute) was added (4 mL) and the resultant clear reaction mixture was heated to reflux on an oil bath (bath temperature 80-85°C) under argon for about 30 hours. The reaction mixture was allowed to cool to room temperature, diluted with ethanol (15 mL), treated with Norite-A (charcoal) and then 10% NaOH solution (4-5 mL) was added (pH > 10). The resultant white solid was filtered and crystallized from ethanol-water to give the title compound (0.204 g, 13%) as a bright white solid, mp: 113-15°C; TLC: R_f =0.4 (CH₂Cl₂:MeOH:NH₄OH; 9:1:2 drops); ¹H NMR (CDCl₃): δ 7.12 (d, 4H, J=8.48Hz, ArH), 7.03 (d, 4H, J=8.45Hz, ArH), 2.56 (m, 2H, 1 X

-CH-), 1.57 (m, 4H, 2 X -CH₂-), 1.21 (d, 6H, J = 6.99Hz, 2 X -CH₃), 0.82 (t, 6H, J = 7.23Hz, 2 X -CH₃); MS(EI): m/e 324 (M⁺1); Anal.: C₂₁H₂₉N₃ (323.46); Calcd. (%): C, 77.97, H, 9.04, N, 12.99; Found (%): C, 77.51, H, 9.08, N, 13.16.

5

Example 5: Preparation of N,N'-bis(4-sec-Butylphenyl)-N-methyl guanidine•HCl

10

Step 1. Preparation of N-Methyl-N-4-sec-butylphenyl cyanamide

A solution of 4-sec-butylphenyl cyanamide (2.3 g, 13.56 mmol) in THF (33 mL) was slowly added to a stirred suspension of sodium hydride (1.08 g, 27 mmol) in THF (12 mL) at room temperature. After 2 hours reflux, the reaction mixture was cooled to 20°C, methyl iodide (7.04 g, 49.6 mmol) was added and the mixture stirred the contents at 20°C. After 19 hours, the reaction was quenched by careful addition of methanol (45 mL) followed by water (100 mL). The aqueous mixture was extracted with methylene chloride (3 X 90 mL), dried over MgSO₄ and the residue was purified on flash chromatography to yield the product (2.0 g, 79%) as a orange syrup. TLC (CHCl₃:CH₃OH; 10:1); R_f=0.76.

20

Step 2. Preparation of N,N'-bis(4-sec-Butylphenyl)-N-methyl guanidine•HCl

Aluminum chloride (0.42 g, 3.1 mmol) was added to a stirred solution of N-4-sec-butylphenyl-N-methyl cyanamide (0.534 g, 2.84 mmol) in chlorobenzene (15 mL) at 145°C. After 10 minutes 4-sec-butylphenyl amine hydrochloride (0.474 g, 2.56 mmol, prepared from 4-sec-butyl aniline and 1.0 M HCl-ether) was added and continued reflux. After 2 hours, the reaction mixture was evaporated and the product was purified by flash chromatography to afford the title compound (0.45 g, 65%) as a syrup. TLC: R_f=0.22 (CH₂Cl₂:CH₃OH; 10:1); ¹H NMR (CD₃OD): δ 7.27 (d, 4H, J=8.18Hz, ArH), 7.20 (d, 4H, J=8.36Hz, ArH), 3.46 (s, 3H, -NCH₃), 2.69-2.58 (m, 2H, 2 X -CH), 1.68-1.57 (m, 4H, 2 X -CH₂), 1.25 (d, 6H, 2 X -

25

30

CH₃), 0.85 (t, 6H, 2 X -CH₃); HRMS: 337.2484 (337.2528 Calcd. for C₂₂H₃₁N₃); HPLC: (CH₃CN:H₂O; 50:50 with 0.1% TFA): 98% pure.

5 Example 6: Preparation of N,N'-Bis(4-sec-butylphenyl)-N,N'-bis-methylguanidine•HCl

Step 1. Preparation of N-Methyl-N-4-sec-butylphenyl cyanamide

10 Prepared as per Step 1 of Example 5 above.

Step 2. Preparation of N-Methyl-N-4-sec-butyl aniline

4-sec-butyl aniline (4.28 g, 28.7 mmol) was dissolved in formic acid (97%, 1.85 g, 40.2 mmol) and magnetically stirred at 100-105°C under argon. After 6 hours, the reaction mixture was cooled to 25°C and diluted with dichloromethane (40 mL). The mixture was washed with saturated sodium bicarbonate (3 X 30 mL), brine (3 X 30 mL) and the organic phase dried over MgSO₄ and then evaporated to afford the formamide (3.85 g, 76%) as an amber syrup, which was used in the next step without further purification.

LiAlH₄-THF solution (1.0 M, 1.0 g, 26 mmol) in THF (23 mL) at ice-bath temperature. After stirring the contents at 25°C for 20 hours, the reaction mixture was combined with saturated solution of sodium sulfate (150 mL) then filtered and washed with THF. The filtrate was evaporated and the residue was chromatographed on silica gel using hexane/ethyl acetate (8:2) as eluent to afford the title compound (1.76 g, 50%) as a liquid.

30 Step 3. Preparation of N,N'-Bis(4-sec-butylphenyl)-N,N'-bis-methylguanidine•HCl

Aluminum chloride (0.39 g, 2.93 mmol) was added to a stirred solution of 4-sec-butylphenyl-N-methyl cyanamide (0.5 g, 2.66 mmol) in

chlorobenzene (14 mL) at 145°C. After 10 minutes 4-sec- butylphenyl-N-methyl amine hydrochloride (0.466 g, 2.34 mmol; prepared from 4-sec-butylphenyl-N-methyl amine and 1.0 M HCl-ether) was added and continued reflux. After 5 hours, the reaction mixture was evaporated on rotavapor and the product was purified by flash chromatography using chloroform/methanol (10:1) to afford the title compound (0.34 g, 38%) as an extremely hygroscopic solid, mp: 65-66°C; TLC: R_f = 0.13 (CHCl_3 : CH_3OH ; 10:1); ^1H NMR (CD_3OD): δ 6.99 (d, 4H, J = 8.46Hz, ArH), 6.80 (d, 4H, J = 8.34Hz, ArH), 3.36 (s, 6H, 2 X -NCH₃), 2.50 (m, 2H, 2 X -CH), 1.58-1.53 (m, 4H, 2 X -CH₂), 1.16 (d, 6H, 2 X -CH₃), 0.79 (t, 6H, J = 7.35Hz, 2 X -CH₃); HRMS: 351.2660 (351.2674 Calcd. for $\text{C}_{23}\text{H}_{33}\text{N}_3$); Anal.: $\text{C}_{23}\text{H}_{33}\text{N}_3 \cdot \text{HCl}$; 1.75 H_2O (418.74); Calcd. (%): C, 66.19, H, 8.64, N, 10.07; Found (%): C, 66.56, H, 8.55, N, 11.27; HPLC: (CH_3CN : H_2O 50:50 with 0.1% TFA): 99% pure.

Example 7: Preparation of N-(5-Acenaphthyl)-N'-(1,2,3,4-tetrahydroquinolinyl)guanidine•mesylate

Part 1: Preparation of 5-Acenaphthyl cyanamide

A mixture of 5 and 3-nitroacenaphthene was reduced with Pd/C in ethyl acetate under hydrogen at 40 psi pressure and the resulting amines were separated by recrystallization from cyclohexane/ethyl acetate. 5-acenaphthyl amine further reacted with cyanogen bromide to yield 5-acenaphthyl cyanamide.

Part 2: Preparation of N-(5-Acenaphthyl)-N'-(1,2,3,4-tetrahydroquinolinyl)guanidine•mesylate

A mixture of 5-acenaphthyl cyanamide (582 mg, 3 mmol), 1,2,3,4-tetrahydroquinolinyl • mesylate (688 mg, 3 mmol), and chlorobenzene (2 mL) in a round bottom flask were heated at 150-160°C for 1 hour and a precipitate formed. The precipitate was collected by filtration, washed with

ether, and dried under vacuum to yield the pure product (1.25 g) as a white solid, mp: 203-204°C; TLC: R_f = 0.3 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CD_3OD): δ ppm 7.70-7.15 (m, ArH, 9H), 3.90 (t, CH_2 , 2H), 3.44-3.35 (m, CH_2 , 4H), 2.87 (t, CH_2 , 2H), 2.69 (s, CH_3 , 3H), 2.15 (m, CH_2 , 2H); MS(EI):
5 m/e 327.2 (M^+ : $\text{C}_{22}\text{H}_{21}\text{N}_3$); Anal.: ($\text{C}_{23}\text{H}_{25}\text{N}_3\text{O}_3\text{S}$); Calcd. (%): C, 65.23, H, 5.95, N, 9.92; Found (%): C, 64.81, H, 6.00, N, 9.74.

Example 8: Preparation of N,N'-Bis-(4-sec-butylphenyl)-2-iminopyrimidazolidine•HBr

10

Part 1: Preparation of N,N'-Bis-(4-sec-butylphenyl)melonyldiamide

Malonyldichloride (10 mmol) in methylene chloride (10 mL) was added dropwise to a solution of 4-sec-butylaniline (52 mmol) in methylene
15 chloride (30 mL) over a period of 10 minutes at 4°C. After the exotherm subsided, the solution was removed from ice bath and stirred at 25°C for 2 hours. The methylene chloride solution was concentrated down to dryness. The crude reaction mixture was purified by column chromatography (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{EtOAc}$ = 2/1) to yield the pure N,N'-bis-(4-sec-
20 butylphenyl)melonyldiamide (75% in yield).

Part 2: Preparation of N,N'-Bis-(4-sec-butylphenyl)-1,3-diaminopropane

Diborane (38.24 mmol, 38.24 mL, 1M) in THF was added dropwise to a dry THF solution of N,N'-bis-(4-sec-butylphenyl)melonyldiamide (9.56
25 mmol) over 10 minutes at 4°C. After 15 minutes, the reaction mixture was heated at refluxing temperature for 16 hours. The reaction mixture was then quenched by aqueous HCl (1M) at 0°C. Then the THF was evaporated, and the solution was basified to pH 14 and extracted with CH_2Cl_2 . The combined organic extracts were dried over MgSO_4 , filtered,
30 and concentrated to yield the pure N,N'-bis-(4-sec-butylphenyl)-1,3-diaminopropane as a yellow liquid (78% in yield).

Part 3: Preparation of N,N'-Bis-(4-sec-butylphenyl)-2-imino-pyrimidazolidine • HBr

N,N'-bis-(4-sec-butylphenyl)-1,3-diaminopropane (1.69 g, 5 mmol) in EtOH (15 mL) was added cyanogen bromide, then the solution was brought to reflux for 16 hours. After the reaction, the mixture was concentrated down and purified by column chromatography (SiO₂, MeOH/CH₂Cl₂ = 0% to 10%). The pure N,N'-bis-(4-sec-butylphenyl)-2-imino-pyrimidazolidine•HBr was obtained as a white solid (75% in yield), mp: >250°C; TLC: R_f=0.5 (SiO₂, CH₂Cl₂/MeOH = 9/1); MS(EI): m/e 363.2 (M⁺: C₂₄H₃₃N₃); Anal.: (C₂₄H₃₃N₃•HBr•1/4 H₂O); Calcd. (%): C, 64.17, H, 7.74, N, 9.40; Found (%): C, 64.18, H, 7.82, N, 9.19.

Example 9: Preparation of N-N'-Bis(3-fluoroanthenyl)guanidine hydrobromide

To a solution of 3-aminofluoroanthene (1.95g, 8.98 mmol) in ethanol (100.0 ml) was slowly added cyanogen bromide (0.922g, 8.7 mmol) while the flask was surrounded by an ice bath and under static Ar. The solution was refluxed for 9.5 hours then let stir at room temperature for 18 hours. The yellow mixture was suction filtered and the yellow solid washed with ether and a little ethyl acetate to yield the title compound as a yellow solid, 1.53 g, 63.0% yield.

Yellow solid; mp: >300°C; R_f=0.24 (10:1 CHCl₃/MeOH); ¹H NMR (300 MHz, CD₃OD) δ 8.10-8.55 (m, 6H, Ar-H), 7.94-8.00 (m, 4H, ArH), 7.73-7.85 (m, 4H, ArH), 7.39-7.42 (m, 4H, ArH); MS(EI): m/e 459 (M⁺ for free base); Anal. Calcd. for C₃₃H₂₁N₃•HBr: C, 73.34, H, 4.1, N, 7.77; Found: C, 73.34, H, 4.48, N, 6.94, 6.76.

Examples 10-145

By methods indicated above and using appropriately substituted reagents, the following compounds were prepared having the specified physical characteristics.

5

Example 10: **N-(5-Acenaphthyl)-N'-[(1-naphthyl)-methylene]guanidine•CH₃SO₃H**

White solid; mp: 262°C.

10 **Example 11:** **N-(2-Naphthyl)-N'-(4-*isopropyl*phenyl)guanidine•HCl**

White solid; mp: 208-10°C; ¹H NMR (CDCl₃): δ 7.82-7.15 (m, 11H, ArH), 2.90-2.75 (m, 1H, -CH), 1.16 (d, 6H, J=6.84Hz, 2 X -CH₃); HRMS: 303.1738 (303.1735 Calcd. for C₂₀H₂₁N₃).

15 **Example 12:** **N,N'-Bis(2-fluorenyl)guanidine•HBr**

White solid; mp: 281°C.

Example 13: **N,N'-Bis(4-*tert*-butylphenyl)guanidine**

20 White solid; mp: 161-62°C; TLC (CHCl₃:CH₃OH; 9:1): R_f=0.54; ¹H NMR (CDCl₃): δ 7.34 (d, 4H, J=8.57Hz, ArH), 7.05 (d, 4H, J=8.52Hz, ArH), 1.28 (s, 18H, 2 X -CH₃); HRMS: 323.2372 (323.2362 Calcd. for C₂₁H₂₉N₃).

25 **Example 14:** **N-(4-*tert*-Butylphenyl)-N'-(2,3,4-trichlorophenyl)guanidine•HCl**

30 White solid; mp: 227-29°C; TLC (CH₂Cl₂:CH₃OH; 9:1): R_f=0.43; ¹H NMR (CD₃OD): δ 7.63 (d, 1H, J=8.7Hz, ArH), 7.52 (d, 2H, J=8.67Hz, ArH), 7.44 (d, 1H, J=8.67Hz, ArH), 7.27 (d, 2H, J=8.66Hz, ArH), 1.33 (s, 9H, 3 X -CH₃); Anal. Calcd. for C₁₇H₁₈N₃Cl₃•HCl (414.17): C, 50.15; H, 4.7; N, 10.32. Found: C, 49.97; H, 4.57; N, 10.22.

Example 15: N-(4-Methoxy-1-naphthyl)-N'-(2,3,4-trichlorophenyl)guanidine•HCl

White solid; mp: 238-40°C; TLC (CH₂Cl₂:CH₃OH; 9:1): R_f=0.48; ¹H NMR (CD₃OD): δ 8.45 (d, 1H, J=8.43Hz, ArH), 8.08 (d, 1H, J=8.24Hz, ArH), 7.82-7.63 (m, 5H, ArH), 7.13 (d, 1H, J=8.27Hz, ArH), 4.19 (s, 3H, -OCH₃); Anal. Calcd. for C₁₈H₁₄N₃Cl₃•HCl (431.14): C, 50.14; H, 3.51; N, 9.75.; Found: C, 49.38; H, 3.59; N, 9.57.

Example 16: N-(2-Naphthyl)-N'-(2-adamantyl)guanidine•HCl

White solid; mp: 219-20°C; ¹H NMR (CDCl₃): δ 7.81-7.22 (m, 7H, ArH), 1.97-1.51 (m, 15H, ArH); Anal. Calcd. for C₂₁H₂₅N₃•HCl (355.45): C, 70.90; H, 7.31; N, 11.82; Found: C, 70.90, H, 7.34; N, 11.73.

Example 17: N-(4-sec-Butylphenyl)-N-benzylguanidine

mp: 65-67°C.; TLC: R_f=0.5 (SiO₂, CH₂Cl₂/MeOH = 9/1); ¹H NMR (CD₃OD): δ 7.30-6.90 (m, ArH, 9H), 4.98 (s, CH₂, 2H), 2.58 (m, CH, 1H), 1.55 (m, CH₂, 2H), 1.20 (d, CH₃, 3H), 0.79 (t, CH₃, 3H); HRMS: 281.1891 (281.1891 calculated for C₁₈H₂₃N₃).

Example 18: N-(5-Acenaphthyl)-N-benzylguanidine

mp: 138-140°C.; TLC: R_f=0.5 (SiO₂, CH₂Cl₂/MeOH = 9/1); ¹H NMR (CD₃OD): δ 7.45-6.93 (m, ArH, 10H), 5.54 (d, CH₂, J=15.8Hz, 1H), 4.63 (d, CH₂, J=15.6Hz, 1H), 3.45-3.40 (m, CH₂, 4H); HRMS: 301.1559 (301.1589 calculated for C₂₀H₁₉N₃).

Example 19: N-(5-Acenaphthyl)-N-(4-isopropylbenzyl)guanidine

mp: 153-155°C; TLC: R_f=0.3 (SiO₂, CH₂Cl₂/MeOH = 9/1); ¹H NMR (CD₃OD): δ ppm 7.60-7.10 (m, ArH, 9H), 5.17 (d, CH₂, J=16.2Hz, 1H), 4.57 (d, CH₂, J=16.1Hz, 1H), 3.45-3.40 (m, CH₂, 4H), 2.85 (m, CH, 1H), 1.20 (d, CH₃, 6H); HRMS: 343.2048 (343.2067 calculated for C₂₃H₂₅N₃); HPLC: 92% pure.

Example 20: N-(4-Cyclohexylphenyl)-N-(4-*isopropylbenzyl*)guanidine•HCl

mp: 130-131°C; TLC: R_f = 0.4 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CD_3OD): δ ppm 7.35-7.12 (m, ArH, 8H), 4.92 (s, CH_2 , 2H), 2.90 (m, CH, 1H), 2.53 (m, CH, 1H), 1.87-1.25 (m, CH_2 , 10H), 1.12 (d, CH_3 , 6H);
5 HRMS: 349.2512 (349.2518 calculated for $\text{C}_{23}\text{H}_{31}\text{N}_3$); HPLC: 99% pure.

Example 21: N-(4-Cyclohexylphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

10 mp: 219-220°C; TLC: R_f = 0.4 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CDCl_3): δ ppm 7.45-7.18 (m, ArH, 8H), 4.88 (s, CH_2 , 2H), 2.55 (s, CH, 1H), 1.90-1.30 (m, CH_2 , 10H), 1.32 (s, CH_3 , 9H); MS(EI): m/e 363.2 (M^+ : $\text{C}_{24}\text{H}_{33}\text{N}_3$); Anal.: ($\text{C}_{24}\text{H}_{33}\text{N}_3\cdot\text{HCl}$); Calcd. (%): C, 72.06, H, 8.57, N, 10.50; Found (%): C, 71.97, H, 8.32, N, 10.33.

Example 22: N-(2-Fluorenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

15 mp: 155-157°C; TLC: R_f = 0.6 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CDCl_3): δ 7.90-7.20 (m, ArH, 11H), 4.95 (s, CH_2 , 2H), 3.95 (s, CH_2 , 2H), 1.32 (s, CH_3 , 9H); MS(EI): m/e 369.2 (M^+ : $\text{C}_{25}\text{H}_{27}\text{N}_3$); Anal.: ($\text{C}_{25}\text{H}_{27}\text{N}_3\cdot\text{HCl}\cdot\text{H}_2\text{O}$); Calcd. (%): C, 70.89, H, 7.14, N, 9.93; Found (%): C, 71.14, H, 6.88, N, 9.77.

Example 23: N-(4-*sec*-Butylphenyl)-N-(*trans*-cinnamyl)guanidine•HCl

25 mp: 169-170°C; TLC: R_f = 0.2 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CDCl_3): δ ppm 7.40-7.20 (m, ArH, 9H), 6.68 (d, CH, J = 16Hz, 1H), 6.25 (dt, CH, J = 16Hz, 1H), 4.55 (d, CH_2 , 2H), 2.66 (m, CH, 1H), 1.60 (m, CH_2 , 2H), 1.25 (d, CH_3 , 3H), 0.83 (t, CH_3 , 3H); Anal.: ($\text{C}_{20}\text{H}_{25}\text{N}_3\cdot\text{HCl}$); Calcd. (%): C, 69.93, H, 7.64, N, 12.24; Found (%): C, 69.79, H, 7.45, N, 12.29.

Example 24: N-(4-*n*-Butoxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

mp: 188-189°C; TLC: R_f = 0.5 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CDCl_3): δ ppm 7.44-6.90 (m, ArH, 8H), 4.84 (s, CH_2 , 2H), 3.97 (t, CH_2 , 2H), 1.73 (p, CH_2 , 2H), 1.49 (p, CH_2 , 2H), 1.32 (s, CH_3 , 9H), 0.94 (t, CH_3 , 3H); MS(EI): m/e 353.3 (M^+ : $\text{C}_{22}\text{H}_{31}\text{N}_3\text{O}$); Anal.: ($\text{C}_{22}\text{H}_{31}\text{N}_3\text{O}\cdot\text{HCl}$); Calcd. (%): C, 67.83, H, 8.29, N, 10.79; Found (%): C, 68.00, H, 8.18, N, 11.04.

Example 25: N-(3-Biphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

mp: 255-256°C; TLC: R_f = 0.5 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CDCl_3): δ 7.70-7.15 (m, ArH, 13H), 4.93 (s, CH_2 , 2H), 1.30 (s, CH_3 , 9H); MS(EI): m/e 357.3 (M^+ : $\text{C}_{24}\text{H}_{27}\text{N}_3$); Anal.: ($\text{C}_{24}\text{H}_{27}\text{N}_3\cdot\text{HCl}$); Calcd. (%): C, 73.25, H, 7.18, N, 10.68; Found (%): C, 73.41, H, 7.18, N, 10.86.

Example 26: N-(5-Indanyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

TLC: R_f = 0.5 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CDCl_3): δ ppm 7.45-6.90 (m, ArH, 7H), 4.86 (s, CH_2 , 2H), 2.87 (m, CH_2 , 4H), 2.08 (m, CH_2 , 2H), 1.29 (s, CH_3 , 9H); MS(EI): m/e 321.2 (M^+ : $\text{C}_{21}\text{H}_{27}\text{N}_3$).

Example 27: N-(3-Trifluoromethoxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

TLC: R_f = 0.5 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CDCl_3): δ ppm 7.65-7.10 (m, ArH, 8H), 4.94 (s, CH_2 , 2H), 1.29 (s, CH_3 , 9H); MS(EI): m/e 365.1 (M^+ : $\text{C}_{19}\text{H}_{22}\text{F}_3\text{N}_3\text{O}$).

Example 28: N-(Methoxy-1-naphthyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

TLC: R_f = 0.5 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CDCl_3): δ ppm 8.35-6.80 (m, ArH, 10H), 5.25 (d, CH_2 , J = 15.6 Hz, 1H), 4.59 (d, CH_2 , J = 15.6 Hz, 1H), 1.29 (s, CH_3 , 9H); MS(EI): m/e 361.5 (M^+ : $\text{C}_{23}\text{H}_{27}\text{N}_3\text{O}$).

Example 29: N-(5-Acenaphthyl)-N'-(indoliny)guanidine•mesylate

mp: 249-250°C; TLC: R_f =0.3 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 9/1); ^1H NMR (CD_3OD): δ ppm 7.70-7.15 (m, ArH, 9H), 4.27 (t, CH_2 , 2H), 3.44-3.35 (m, CH_2 , 4H), 3.30 (t, CH_2 , 2H), 2.69 (s, CH_3 , 3H); MS(EI): m/e 313.2 (M^+ : $\text{C}_{21}\text{H}_{19}\text{N}_3$); Anal.: ($\text{C}_{22}\text{H}_{23}\text{N}_3\text{O}_3\text{S}\cdot 1/2 \text{H}_2\text{O}$); Calcd. (%): C, 63.13, H, 5.78, N, 10.04; Found (%): C, 63.13, H, 5.70, N, 10.06.

Example 30: N,N'-Bis(3-biphenyl)guanidine•HBr

White solid; mp: 176-178°C; TLC($\text{CH}_2\text{Cl}_2:\text{CH}_3\text{OH}$; 15:1): R_f =0.50; ^1H NMR (CD_3OD): 7.66-7.33 (m, ArH). Anal. Calcd. for $\text{C}_{25}\text{H}_{33}\text{N}_3\cdot\text{HBr}$ (444.37): C, 67.27; H, 5.42; N, 9.41.; Found: C, 67.01, H, 4.86, N, 9.29.

Example 31: N,N'-Di-(3-*tert*-butylphenyl)guanidine•HBr

White solid; mp: 206-208°C; TLC (CH_2Cl_2 : MeOH; 15:1): R_f =0.50; ^1H NMR (CD_3OD): 7.53-7.23 (m, 8H, ArH), 1.34 (s, 18H, $-\text{C}(\text{CH}_3)_3$); Anal. Calcd. for $\text{C}_{21}\text{H}_{29}\text{N}_3\cdot\text{HBr}$ (404.39): C, 62.37, H, 7.48, N, 10.39; Found: C, 62.49, H, 7.45, N, 10.52.

Example 32: N,N'-Bis-(4-methoxy-1-naphthyl)guanidine•HBr

Light green solid; mp: 236-238°C; TLC ($\text{CH}_2\text{Cl}_2:\text{MeOH}$; 15:1): R_f =0.48; ^1H NMR (CD_3OD): 8.32 (d, 2H, $J=7.72\text{Hz}$, ArH), 7.97 (d, 2H, $J=8.18\text{Hz}$, ArH), 7.69 (t, 2H, $J=7.52\text{Hz}$, ArH), 7.57 (q, 4H, $J=7.52\text{Hz}$, ArH), 7.00 (d, 2H, $J=8.18\text{Hz}$, ArH), 4.05 (s, 3H, $-\text{OCH}_3$); Anal. Calcd. for $\text{C}_{23}\text{H}_{21}\text{N}_3\text{O}_2\cdot\text{HBr}$ (452.35): C, 61.07, H, 4.90, N, 9.29; Found: C, 60.98, H, 4.99, N, 9.50.

Example 33: N,N'-Bis-(5-indanyl)guanidine•HBr

White solid; mp: 118-120°C; TLC ($\text{CH}_2\text{Cl}_2:\text{MeOH}$; 15:1): R_f =0.31; ^1H NMR (CD_3OD): 7.30 (d, $J=8.25\text{Hz}$, 2H, ArH), 7.18 (s, 2H, ArH), 7.07 (d, 2H, $J=7.27\text{Hz}$, ArH), 2.93 (q, $J=7.10\text{Hz}$, 8H, $-\text{CH}_2$), 2.11 (p, 4H, $-\text{CH}_2$).

$J = 7.42\text{Hz}$); Anal. Calcd. for $\text{C}_{19}\text{H}_{21}\text{N}_3 \cdot \text{HBr}$ (372.31): C, 61.30, H, 5.96, N, 11.29; Found: C, 61.12; H, 5.81; N, 11.11.

Example 34: N,N' -Bis-(3-*sec*-butylphenyl)guanidine•HBr

5 White solid; mp: 85-87°C; TLC(CH_2Cl_2 :MeOH; 15:1): $R_f = 0.31$; ^1H NMR (CD_3OD): 7.39 (t, $J = 7.57\text{Hz}$, 2H, ArH), 7.17 (t, $J = 7.15\text{Hz}$, 4H, ArH), 2.65 (m, 2H, -CH), 1.64 (m, 4H, $-\text{CH}_2$), 1.58 (d, $J = 0.82\text{Hz}$, 6H, $-\text{CH}_3$), 0.84 (t, $J = 1.90\text{Hz}$, 6H, $-\text{CH}_3$); Anal. Calcd. for $\text{C}_{21}\text{H}_{29}\text{N}_3 \cdot \text{HBr}$ (404.39): C, 62.37, H, 7.29, N, 10.39; Found: C, 62.58, H, 7.29, N, 10.61.

Example 35: N,N' -Bis(4-*tert*-butylphenyl)- N -methylguanidine•HBr

10 White solid; mp: 194-196°C; TLC (CH_2Cl_2 : MeOH; 10:1): $R_f = 0.35$; ^1H NMR (CD_3OD): 7.60-7.57 (d, $J = 8.39\text{Hz}$, 2H, ArH), 7.51-7.48 (d, $J = 8.24\text{Hz}$, 2H, ArH), 7.38-7.35 (d, $J = 8.52\text{Hz}$, 2H, ArH), 7.22-7.20 (d, $J = 8.24\text{Hz}$, 2H, ArH), 3.46 (s, 1H, $-\text{CH}_3$), 1.35 (s, 9H, $-\text{C}(\text{CH}_3)_3$), 1.33 (s, 9H, $-\text{C}(\text{CH}_3)_3$); Anal. Calcd. for $\text{C}_{22}\text{H}_{31}\text{N}_3 \cdot \text{HCl}$ (373.97): C, 70.66, H, 8.62, N, 11.29; Found: C, 70.69, H, 8.31, N, 11.39.

Example 36: N,N' -Bis(4-*tert*-butylphenyl)- N,N' -methylguanidine•HBr

20 White solid; mp: 175-177°C; TLC(CH_2Cl_2 :MEOH; 10:1): $R_f = 0.39$; ^1H NMR (CD_3OD): 7.21-7.18 (d, $J = 8.52\text{Hz}$, 4H, ArH), 6.80-6.78 (d, $J = 8.51\text{Hz}$, 4H, ArH), 3.35 (s, 6H, $-\text{CH}_3$), 1.26 (s, 18H, $-\text{C}(\text{CH}_3)_3$); Anal. Calcd. for $\text{C}_{23}\text{H}_{33}\text{N}_3 \cdot \text{HCl}$ (388.00): C, 71.20, H, 8.83, N, 10.83; Found: C, 70.88, H, 8.61, N, 10.75.

Example 37: N,N' -Bis(4-*n*-butylphenyl)guanidine•mesylate

30 White solid: M.P. 120-22°C; ^1H NMR (300 MHz, CD_3OD): δ 7.28 (dd, 4H, $J = 8.5\text{Hz}$ and $J = 2.2\text{Hz}$, ArH); 7.23 (dd, 4H, $J = 8.5\text{Hz}$ and $J = 2.2\text{Hz}$, ArH); 2.69 (s, 3H, $-\text{SO}_3\text{H}-\text{CH}_3$); 2.64 (t, 4H, $J = 7.69\text{Hz}$, 2 X ArCH₂-); 1.65-1.55 (m, 4H, 2 X $-\text{CH}_2-\text{CH}_2\text{CH}_3$); 1.43-1.30 (m, 4H, 2 X

-CH₂-CH₃); 0.94 (t, 6H, J = 7.35Hz, 2 X -CH₃); Anal Calcd. for C₂₁H₂₉N₃•CH₄SO₃ (419.56): C, 62.98, H, 7.93, N, 10.01; Found: C, 62.96, H, 7.69, N, 9.93.

5 **Example 38:** **N-(5-Acenaphthyl)-N'-(1-methyl-2-phenoxyethyl) guanidine•HCl**
 ((5-C₁₂H₉)NHC(=NH)NHCH(CH₃)CHOC₆H₅•HCl)

Bubbly white solid: mp: 107-110°C; R_f=0.24 (10:1 CHCl₃/MeOH);
 10 ¹H NMR (300 MHz, CD₃OD): δ 7.29-7.56 (m, 7H, ArH), 6.96-7.01 (t, 3H, J = 16Hz, ArH), 4.20-4.32 (m, 1H, CH), 4.11-4.89 (m, 1H, CH), 3.90-4.00 (m, 1H, CH), 3.40-3.50 (m, 4H, CH₂); MS(EI): m/e 345 (M⁺ for free base); Anal. Calcd. for C₂₂H₂₃N₃O•HCl: C, 69.19, H, 6.33, N, 11.00; Found: C, 69.07, H, 6.40, N, 10.87.

15 **Example 39:** **N-(5-Acenaphthyl)-N'-(piperonyl)guanidine•HCl**

Tan solid: mp: 142-144°C; R_f=0.20 (10:1 CHCl₃/MeOH); ¹H NMR (300 MHz, CD₃OD): δ 7.34-7.54 (m, 5H, ArH), 6.84 (s, 3H, ArH), 5.98 (s, 2H, CH₂), 4.41 (s, 2H, CH₂), 3.43-3.50 (m, 4H, 2-CH₂); MS(EI): m/e 345 (M⁺ for free base); Anal. Calcd. for C₂₁H₁₉N₃O₂•HCl•0.5C₄H₈O₂: C, 64.92, 20 H, 5.69, N, 9.88; Found: C, 64.37, H, 5.93, N, 10.25.

Example 40: **N-(5-Acenaphthyl)-N'-(1-methyl-2-(4-chlorophenyl)ethyl) guanidine•HCl** ((5-C₁₂H₉)NHC(=NH)NHCH(CH₃)CH(4-ClC₆H₄)•HCl)

Bubbly amber solid: M.P. 123-125°C; R_f=0.22 (10:1 CHCl₃/MeOH);
 25 ¹H NMR (300 MHz, CD₃OD): δ 7.51-7.54 (m, 1H, ArH), 7.18-7.37 (m, 8H, ArH), 4.00-4.12 (m, 1H, CH), 3.41-3.51 (m, 4H, 2-CH₂), 2.90-3.00 (dd, 1H, CH), 2.71-2.82 (dd, 1H, CH) 1.33-1.35 (d, 3H, J = 6.53Hz, CH₃); MS(EI): m/e 363 (M⁺ for free base); Anal. Calcd. for C₂₂H₂₂N₃Cl•HCl: C, 30 66.00, H, 5.79, N, 10.50; Found: C, 65.94, H, 5.91, N, 10.30.

Example 41: N-(5-Acenaphthyl)-N'-(1,2-diphenylethyl)guanidine•HCl

Cream colored solid: M.P. 167-170°C; R_f = 0.18 (10:1 $\text{CHCl}_3/\text{MeOH}$); ^1H NMR (300 MHz, CD_3OD): δ 7.16-7.44 (m, 15H, ArH), 5.1-5.2 (m, 1H, CH) 3.30-3.41 (m, 4H, 2- CH_2), 3.13-3.30 (m, 2H, CH_2); MS(EI): m/e 391 (M^+ for free base); Anal. Calcd. for $_{27}\text{H}_{25}\text{N}_3\bullet\text{HCl}$: C, 75.77, H, 6.12, N, 9.82; Found: C, 75.60, H, 6.00, N, 9.71.

Example 42: N-(3-Benzyloxyphenyl)-N'-(4-benzyloxyphenyl)guanidine•HCl

Bubbly pink solid: mp: 59-62°C; R_f = 0.19 (10:1 $\text{CHCl}_3/\text{MeOH}$); ^1H NMR (300 MHz, CD_3OD): δ 7.24-7.45 (m, 13H, ArH), 6.90-7.10 (m, 5H, ArH), 5.12 (brs, 4H, 2- CH_2); MS(EI): m/e 423 (M^+ for free base); Anal. Calcd. for $\text{C}_{27}\text{H}_{25}\text{N}_3\text{O}_2\bullet\text{HCl}\bullet 1.5\text{H}_2\text{O}$: C, 66.64, H, 6.01, N, 8.64; Found: C, 67.21, 66.95, H, 6.12, 6.01, N, 8.25, 7.98.

Example 43: N-(5-Acenaphthyl)-N'-(3-phenylpropyl)guanidine•HCl

White solid: mp: 84°C; R_f = 0.28 (1:5 $\text{MeOH}:\text{EtOAc}$); ^1H NMR (300 MHz, CD_3OD): δ 7.54-7.57 (m, 2H, ArH), 7.35-7.41 (m, 3H, ArH), 7.25-7.35 (m, 2H, ArH), 7.18-7.24 (m, 3H, ArH), 3.40-3.48 (brs, 4H, 2- CH_2), 3.29-3.34 (t, J = 7Hz, 2H, $\text{CH}_2\text{-N}$), 2.66-2.71 (t, J = 8Hz, 2H, CH_2), 1.91-2.00 (m, 2H, $\text{CH}_2\text{-Ar}$); MS(EI): m/e 329 (M^+ for free base); Anal. Calcd. for $\text{C}_{24}\text{H}_{21}\text{N}_3\bullet\text{HCl}\bullet 0.5\text{H}_2\text{O}$: C, 70.48, H, 6.72, N, 11.26; Found: C, 70.52, H, 6.65, N, 11.18.

Example 44: N-(5-Acenaphthyl)-N'-(2-methyl-2-phenylethyl)guanidine•HCl

White solid: mp: 110°C; R_f = 0.26 (1:5 $\text{MeOH}:\text{EtOAc}$); ^1H NMR (300 MHz, CD_3OD): δ 7.49-7.54 (t, J = 7Hz, 1H, ArH), 7.24-7.38 (m, 8H, ArH), 7.15-7.18 (m, 1H, ArH), 3.51-3.53 (d, J = 6Hz, 2H, $\text{CH}_2\text{-Ar}$), 3.38-3.48 (m, 4H, 2- CH_2), 3.03-3.10 (m, 1H, CH), 1.30-1.33 (d, J = 7Hz, 3H, CH_3);

MS(EI): m/e 329 (M^+ for free base); Anal. Calcd. for $C_{24}H_{21}N_3 \cdot HCl \cdot 0.5H_2O$: C, 70.48, H, 6.72, N, 11.26; Found: C, 70.62, H, 6.58, N, 11.17.

Example 45: N,N' -(*sec*-Butylphenyl)- N' -(2-phenoxyethyl)guanidine•HCl

5 Semisolid: $R_f = 0.20$ (1:5 MeOH:EtOAc); 1H NMR (300 MHz, CD_3OD): δ 7.22-7.34 (m, 6, ArH), 7.16-7.22 (d, $J = 7$ Hz, 2, ArH), 7.00-7.07 (d, $J = 7$ Hz, 2H, ArH), 6.87-6.98 (m, 3H, ArH), 4.20 (brs, 4H, 2- CH_2), 2.51-2.70 (m, 2H, 2-CH), 1.53-1.68 (m, 4H, 2- CH_2), 1.20-1.27 (2d, $J = 6$ Hz, 6H, 2- CH_3); MS(EI): m/e 444 (M^+ for free base); Anal. Calcd. for $C_{29}H_{37}N_3O \cdot 0.75HCl$: C, 73.96, H, 8.08, N, 8.92; Found: C, 74.21, H, 8.26, N, 9.21.

Example 46: N,N' -(*sec*-Butylphenyl)- N' -(*n*-pentyl)guanidine•HCl

15 White solid: $R_f = 0.17$ (1:5 MeOH:EtOAc); 1H NMR (300 MHz, CD_3OD): δ 7.31-7.38 (m, 4H, ArH), 7.25-7.28 (d, $J = 8.5$ Hz, 2H, ArH), 7.16-7.19 (d, $J = 8.5$ Hz, 2H, ArH), 3.75-3.80 (t, $J = 8$ Hz, 2H, N- CH_2), 2.59-2.71 (m, 2H, 2-CH), 1.56-1.68 (m, 6H, 3- CH_2), 1.30-1.34 (m, 4H, 2- CH_2), 1.22-1.26 (2d, $J = 7$ Hz, 6H, 2- CH_3), 0.80-0.91 (m, 9H, 3- CH_3); MS(EI): m/e 394 (M^+ for free base); Anal. Calcd. for $C_{26}H_{39}N_3 \cdot HCl \cdot H_2O$: C, 69.69, H, 9.45, N, 9.38; Found: C, 69.89, H, 8.88, N, 10.09.

Example 47: N -(1-Naphthyl)- N -(4-*tert*-butylbenzyl)guanidine•HCl

1H NMR ($CDCl_3$): δ ppm 8.35-7.10 (m, ArH, 11H), 5.25 (d, CH_2 , $J = 15.6$ Hz, 1H), 4.67 (d, CH_2 , $J = 15.6$ Hz, 1H), 1.29 (s, CH_3 , 9H). MS(EI): m/e 331 (M^+ : $C_{22}H_{25}N_3$); Anal. ($C_{22}H_{25}N_3 \cdot HCl$): Calcd. (%): C, 71.82, H, 7.12, N, 11.42; Found (%): C, 71.79, H, 7.12, N, 11.43; TLC: $R_f = 0.50$ (SiO_2 , CH_2Cl_2 /MeOH = 9/1); mp: 241-242°C.

Example 48: N-(3-Iodophenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

^1H NMR (CD_3OD): δ ppm 7.78-7.13 (m, ArH, 8H), 4.87 (s, ArCH_2 , 2H), 1.30 (s, CH_3 , 9H); MS(EI): m/e 407.3 (M^+ : $\text{C}_{18}\text{H}_{22}\text{N}_3\text{I}$); Anal.

5 ($\text{C}_{18}\text{H}_{22}\text{IN}_3\cdot\text{HCl}$): cal. (%): C, 48.72, H, 5.22, N, 9.47; found (%): C, 48.90, H, 5.46, N, 9.52; TLC: $R_f=0.42$ (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH} = 10/1$); mp: 271-272°C.

Example 49: N-(4-Chloro-1-naphthyl)-N-(4-*tert*-benzyl)guanidine•HCl

10 ^1H NMR (CD_3OD): δ ppm 8.37-7.13 (m, ArH, 10H), 5.25 (d, CH_2 , $J=15.5\text{Hz}$, 1H), 4.69 (d, CH_2 , $J=15.4\text{Hz}$, 1H), 1.26 (s, CH_3 , 9H); MS(EI): m/e 365.1 (M^+ : $\text{C}_{22}\text{H}_{24}\text{ClN}_3$); Calcd. ($\text{C}_{22}\text{H}_{24}\text{ClN}_3\cdot\text{HCl}\cdot\text{H}_2\text{O}$): Calcd. (%): C, 62.86, H, 6.47, N, 10.03; Found (%): C, 63.19, H, 6.27, N, 9.69; TLC: $R_f=0.52$ (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH} = 10/1$); mp: 115-117°C.

Example 50: N-(4-*tert*-Butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl


15 ^1H NMR (CDCl_3): δ ppm 7.51-7.18 (m, ArH, 8H), 4.93 (s, ArCH_2 , 2H), 1.33 (s, CH_3 , 9H), 1.31 (s, CH_3 , 9H); MS(EI): m/e 338 (M^+ : $\text{C}_{22}\text{H}_{31}\text{N}_3$); Anal. ($\text{C}_{22}\text{H}_{31}\text{N}_3\cdot\text{HCl}$): Calcd. (%): C, 70.66, H, 8.62, N, 11.24; Found (%): C, 70.43, H, 8.42, N, 11.17; TLC: $R_f=0.45$ (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH} = 10/1$); mp: 290-291°C.

Example 51: N-(4-Iodophenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

25 ^1H NMR (CD_3OD): δ ppm 7.81-7.02 (m, ArH, 8H), 4.89 (s, ArCH_2 , 2H), 1.30 (s, CH_3 , 9H); MS(EI): m/e 408 (M^+ : $\text{C}_{18}\text{H}_{22}\text{N}_3\text{I}$); Anal. ($\text{C}_{18}\text{H}_{22}\text{IN}_3\cdot\text{HCl}$): Calcd. (%): C, 48.72, H, 5.22, N, 9.47; Found (%): C, 48.72, H, 5.26, N, 9.28; TLC: $R_f=0.45$ (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH} = 10/1$); mp: 219-220°C.

30

Example 52: N-(1-Naphthylmethyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl



¹H NMR (CDCl₃): δ ppm 7.95-7.10 (m, ArH, 11H), 5.10 (s, CH₂, 2H),
 5 4.58 (s, CH₂, 2H), 1.32 (s, CH₃, 9H); MS(EI): m/e 346 (M⁺: C₂₃H₂₇N₃);
 Anal. (C₂₃H₂₇N₃•HCl•H₂O): Calcd. (%): C, 69.07, H, 7.56, N, 10.51; Found
 (%): C, 68.70, H, 7.71, N, 10.19; TLC: R_f=0.50 (SiO₂, CH₂Cl₂/MeOH =
 9/1); mp: 134-135°C.

Example 53: N-(5-Acenaphthyl)-N-(3-phenoxybenzyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.62-6.80 (m, ArH, 14H), 5.20 (d, CH₂,
 J = 15.7Hz, 1H), 4.70 (d, CH₂, J = 15.9Hz, 1H), 3.46-3.40 (m, CH₂, 4H);
 MS(EI): m/e 393.5 (M⁺: C₂₆H₂₃ON₃); Anal. (C₂₆H₂₃ON₃•HCl): Calcd. (%): C,
 72.63, H, 5.63, N, 9.77; Found (%): C, 72.77, H, 5.57, N, 9.67; TLC:
 15 R_f=0.45 (SiO₂, CH₂Cl₂/MeOH = 10/1); mp: 100-101°C.

**Example 54: N-(3-Trifluoromethylphenyl)-N-(4-*tert*-butyl
benzyl)guanidine•HCl**

¹H NMR (CD₃OD): δ ppm 7.63-7.13 (m, ArH, 8H), 4.92 (s, ArCH₂,
 20 2H), 1.29 (s, CH₃, 9H); MS(EI): m/e 349.1 (M⁺: C₁₉H₂₂F₃N₃); Anal.
 (C₁₉H₂₂F₃N₃•HCl): Calcd. (%): C, 59.14, H, 6.01, N, 10.89; Found (%): C,
 59.14, H, 6.12, N, 10.88; TLC: R_f=0.49 (SiO₂, CH₂Cl₂/MeOH = 10/1);
 mp: 281-282°C.

**Example 55: N-(3-Methylthiophenyl)-N-(4-*tert*-butyl
benzyl)guanidine•HCl**

¹H NMR (CD₃OD): δ ppm 7.42-7.15 (m, ArH, 8H), 4.87 (s, ArCH₂,
 2H), 2.35 (s, SCH₃, 3H), 1.30 (s, CH₃, 9H); MS(EI): m/e 327.3 (M⁺:
 C₁₉H₂₅SN₃); Anal. (C₁₉H₂₅SN₃•HCl): Calcd. (%): C, 62.70, H, 7.20, N,
 25 11.55; Found (%): C, 62.81, H, 7.27, N, 11.57; TLC: R_f=0.37 (SiO₂,
 30 CH₂Cl₂/MeOH = 10/1); mp: 247-248°C.

Example 56: N-(5-Acenaphthyl)-N-(3-iodobenzyl)guanidine•HCl

¹H NMR (CDCl₃): δ ppm 7.62-7.00 (m, ArH, 9H), 5.46 (d, CH₂, J=0.55Hz, 1H), 4.75 (d, CH₂, J=1.98Hz, 1H), 3.49-3.42 (m, CH₂, 4H); MS(EI): m/e 427.1 (M⁺: C₂₀H₁₈IN₃); Anal. (C₂₀H₁₈IN₃•HCl): Calcd. (%): C, 51.80, H, 4.13, N, 9.06; Found (%): C, 52.00, H, 4.14, N, 9.00; TLC: R_f=0.46 (SiO₂, CH₂Cl₂/MeOH = 10/1); mp: 249-250°C.

Example 57: N-(5-Acenaphthyl)-N-(cinnamyl)guanidine•HCl

¹H NMR (CDCl₃): δ ppm 7.62-7.18 (m, ArH, 10H), 6.65 (d, CH₂, J=15.66Hz, 1H), 6.30 (dd, CH₂, J=15.5Hz, 1H), 5.27 (m, =CH, 0.5H), 4.92 (m, =CH, 0.5H), 4.40 (m, =CH, 1H), 3.57-3.41 (m, CH₂, 4H); MS(EI): m/e 327.2 (M⁺: C₂₂H₂₁N₃); Anal. (C₂₂H₂₁N₃•HCl•3/4H₂O): Calcd. (%): C, 70.01, H, 6.27, N, 11.13; Found (%): C, 70.21, H, 6.31, N, 11.10; TLC: R_f=0.46 (SiO₂, CH₂Cl₂/MeOH = 10/1); mp: 205-206.5°C.

Example 58: N-(5-Acenaphthyl)-N-(4-iodobenzyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.65-7.00 (m, ArH, 9H), 5.20 (d, CH₂, J=15.7Hz, 1H), 4.70 (d, CH₂, J=15.9Hz, 1H), 3.46-3.40 (m, CH₂, 4H); MS(EI): m/e 427.2 (M⁺: C₂₀H₁₈IN₃); Anal. (C₂₀H₁₈IN₃•HCl•3/2H₂O): Calcd. (%): C, 48.95, H, 4.52, N, 8.56; Found (%): C, 48.62, H, 4.42, N, 8.39; TLC: R_f=0.43 (SiO₂, CH₂Cl₂/MeOH = 10/1); mp: 269-270°C.

Example 59: N-(5-Acenaphthyl)-N-(4-trifluoromethoxybenzyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.56-7.17 (m, ArH, 9H), 5.25 (d, CH₂, J=15.9Hz, 1H), 4.82 (d, CH₂, J=15.7Hz, 1H), 3.43-3.41 (m, CH₂, 4H); MS(EI): m/e 385.3 (M⁺: C₂₁H₁₈F₃ON₃); Anal. (C₂₁H₁₈F₃ON₃•HCl•2H₂O): Calcd. (%): C, 55.09, H, 5.06, N, 9.18; Found (%): C, 55.33, H, 4.78, N, 9.08; TLC: R_f=0.53 (SiO₂, CH₂Cl₂/MeOH = 10/1); mp: 120-122°C.

Example 60: N-(5-Acenaphthyl)-N'-((4-*tert*-butylphenyl)-(4-*sec*-butylphenyl)-N'-methylguanidine)•HCl

¹H NMR (CD₃OD): δ ppm 8.20-7.15 (m, ArH, 13H), 6.10 (d, CH, 1H), 3.43-3.41 (m, CH₂, 4H), 2.62 (m, CH, 1H), 1.59 (m, CH₂, 2H), 1.29 (s, CH₃, 9H), 1.21 (d, CH₃, J = 7.0Hz, 3H), 0.8 (t, CH₃, J = 7.0Hz, 3H); MS(EI): m/e 489.4 (M⁺: C₃₄H₃₉N₃); Anal. (C₃₄H₃₉N₃•HCl): Calcd. (%): C, 77.61, H, 7.66, N, 7.99; Found (%): C, 77.43, H, 7.45, N, 7.97; TLC: R_f = 0.5 (SiO₂, CH₂Cl₂/MeOH = 10/1); mp: 173-174°C.

Example 61: N-(4-Butoxyphenyl)-N,N'-bis(4-*tert*-butylbenzyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.50-6.60 (m, ArH, 12H), 4.90 (s, CH₂, 2H), 4.43 (s, CH₂, 2H), 3.97 (t, CH₂, J = 6.4Hz, 2H), 1.73 (m, CH₂, 2H), 1.49 (m, CH₂, 2H), 1.31 (s, CH₃, 18H), 0.94 (t, CH₃, J = 7.4Hz, 3H); MS(EI): m/e 499.3 (M⁺: C₃₃H₄₅N₃O₁); Anal. (C₃₃H₄₅N₃O₁•HCl): Calcd. (%): C, 73.92, H, 8.65, N, 7.84; Found (%): C, 73.87, H, 8.46, N, 7.91; TLC: R_f = 0.4 (SiO₂, CH₂Cl₂/MeOH = 10/1); mp: 112-113°C.

Example 62: N-(3-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.38-6.81 (m, ArH, 8H), 4.86 (s, ArCH₂, 2H), 2.53 (m, CH, 1H), 1.45 (m, CH₂, 2H), 1.29 (s, CH₃, 9H), 1.12 (d, CH₃, J = 6.93Hz, 3H), 0.71 (t, CH₃, J = 7.37Hz, 3H); MS(EI): m/e 337.4 (M⁺: C₂₂H₃₁N₃); Anal. (C, H, N; C₂₂H₃₁N₃•HCl•0.5H₂O): Calcd. (%): C, 68.79, H, 8.68, N, 10.97; Found (%): C, 69.23, H, 8.35, N, 10.92; TLC: R_f = 0.46 (SiO₂, CH₂Cl₂/MeOH = 10/1); mp: 190-191°C.

Example 63: N-(3-*tert*-Butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

¹H NMR (CDCl₃): δ ppm 7.51-7.19 (m, ArH, 8H), 4.92 (s, ArCH₂, 2H), 1.34 (s, CH₃, 9H), 1.31 (s, CH₃, 9H); MS(EI): m/e 337.4 (M⁺: C₂₂H₃₁N₃); Anal. (C, H, N; C₂₂H₃₁N₃•HCl): Calcd. (%): C, 70.66, H, 8.62, N,

11.24; Found (%): C, 70.50, H, 8.55, N, 11.29; TLC: R_f = 0.45 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 10/1); mp: 291-292°C.

Example 64: N-(3-Pentoxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

5 ^1H NMR (CD_3OD): δ ppm 7.41-7.20 (m, ArH, 8H), 4.93 (s, ArCH_2 , 2H), 3.90 (t, OCH_2 , J = 6.49Hz, 2H), 1.78-1.60 (m, CH_2 , 2H), 1.44-1.40 (m, 2CH_2 , 4H), 1.31 (s, CH_3 , 9H), 0.94 (t, CH_3 , J = 6.98Hz, 3H); MS(EI): m/e 367.3 (M^+ : $\text{C}_{23}\text{H}_{33}\text{N}_3\text{O}$); Anal. (C, H, N; $\text{C}_{23}\text{H}_{33}\text{N}_3\text{O}\cdot\text{HCl}\cdot 0.5\text{H}_2\text{O}$); Calcd. (%): C, 66.89, H, 8.54, N, 10.17; Found (%): C, 66.98, H, 8.33, N, 10.05; 10 TLC: R_f = 0.51 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 10/1); mp: 198-199°C.

Example 65: N-(5-Acenaphthyl)-N-(4-benzyloxybenzyl)guanidine•HCl
((5- C_{12}H_9)(4-($\text{C}_6\text{H}_5\text{CH}_2\text{O}$) $\text{C}_6\text{H}_4\text{CH}_2$) $\text{NC}(=\text{NH})\text{NH}_2$)•HCl)

15 ^1H NMR (CDCl_3): δ ppm 7.56-6.86 (m, ArH, 14H), 5.31 (d, CH_2 , J = 15.02Hz, 1H), 5.02 (s, CH_2 , 2H), 4.70 (d, CH_2 , J = 15.41Hz, 1H), 3.50-3.41 (m, 2CH_2 , 4H); MS(EI): m/e 407.3 (M^+ : $\text{C}_{27}\text{H}_{25}\text{N}_3\text{O}$); Anal. (C, H, N; $\text{C}_{27}\text{H}_{25}\text{N}_3\text{O}\cdot\text{HCl}$); Calcd. (%): C, 73.04, H, 5.90, N, 9.46; Found (%): C, 72.93, H, 5.68, N, 9.30; TLC: R_f = 0.51 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 10/1); mp: 118-119°C.

Example 66: N-(4-*sec*-Butylphenyl)-N-(4-benzyloxybenzyl)guanidine•HCl

25 ^1H NMR (CD_3OD): δ ppm 7.39-6.92 (m, ArH, 13H), 5.05 (s, OCH_2 , 2H), 4.82 (s, CH_2 , 2H), 2.62 (m, CH, 1H), 1.60 (m, CH_2 , 2H), 1.21 (d, CH_3 , J = 6.96Hz, 3H), 0.81 (t, CH_3 , J = 7.4Hz, 3H); MS(EI): m/e 387.3 (M^+ : $\text{C}_{25}\text{H}_{29}\text{N}_3\text{O}$); Anal. (C, H, N; $\text{C}_{25}\text{H}_{29}\text{N}_3\text{O}\cdot\text{HCl}\cdot 0.6\text{H}_2\text{O}$); Calcd. (%): C, 69.06, H, 7.23, N, 9.66; Found (%): C, 68.86, H, 6.83, N, 9.80; TLC: R_f = 0.48 (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$ = 10/1); mp: 84-85°C.

30

Example 67: N-(4-Benzyloxyphenyl)-N-(4-benzyloxybenzyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.40-6.93 (m, ArH, 18H), 5.08 (s, OCH₂, 2H), 5.06 (s, OCH₂, 2H), 4.79 (s, CH₂, 2H); MS(EI): m/e 437.2 (M⁺: C₂₈H₂₇N₃O₂); Anal. (C, H, N; C₂₈H₂₇N₃O₂•HCl); Calcd. (%): C, 70.95, H, 5.95, N, 8.86; Found (%): C, 70.81, H, 5.71, N, 8.71; TLC: R_f=0.44 (SiO₂, CH₂Cl₂/MeOH = 10/1); mp: 193-194°C.

Example 68: N-(5-Acenaphthyl)-N-(3-benzyloxybenzyl)guanidine•HCl

¹H NMR (CDCl₃): δ ppm 7.45-6.55 (m, ArH, 14H), 5.25-5.45 (m, CH₂, 1H), 4.86 (s, CH₂, 2H), 4.45-4.65 (d, CH₂, 1H), 3.25-3.50 (m, 2CH₂, 4H); MS(EI): m/e 407.4 (M⁺: C₂₇H₂₅N₃O); Anal. (C, H, N; C₂₇H₂₅N₃O•HCl•H₂O); Calcd. (%): C, 70.20, H, 6.11, N, 9.10; Found (%): C, 70.42, H, 6.00, N, 9.18; TLC: R_f=0.38 (SiO₂, CHCl₃/MeOH = 10/1); mp: 140-141°C.

Example 69: N-(4-Isopropylphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.12-7.41 (m, ArH, 8H), 4.88 (s, CH₂, 2H), 2.80-3.00 (m, CH, 1H), 1.30 (s, CMe₃, 9H), 1.23 (d, CH₃, J=6.9Hz, 3H); MS(EI): m/e 324 (M⁺: C₂₁H₂₉N₃); HRMS: 323.2366 (Calcd. 323.2361); Anal. (C, H, N; C₂₁H₂₉N₃•HCl); Calcd. (%): C, 70.08, H, 8.40, N, 11.67; Found (%): C, 69.85, H, 8.24, N, 11.90; TLC: R_f=0.64 (SiO₂, CHCl₃/MeOH = 10/1); HPLC: 100%; mp: 260-261°C.

Example 70: N-(4-Benzyloxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.00-7.50 (m, ArH, 13H), 5.08 (s, OCH₂, 2H), 4.84 (s, CH₂, 2H), 1.30 (s, CMe₃, 9H); MS(EI): m/e 388 (M⁺: C₂₅H₂₉N₃O); HRMS: 387.2315 (Calcd.: 387.2311); TLC: R_f=0.49 (SiO₂, CHCl₃/MeOH = 10/1); HPLC: 99.8%; mp: 178-179°C.

Example 71: N-(4-Hexylphenyl)-N-(4-hexylbenzyl)guanidine•mesylate

¹H NMR (CD₃OD): δ ppm 7.05-7.30 (m, ArH, 8H), 4.86 (s, CH₂, 2H), 2.69 (s, CH₃SO₃H, 3H), 2.55-2.64 (m, 2CH₂, 4H), 1.50-1.70 (m, 2CH₂, 4H), 1.25-1.40 (m, 6CH₂, 12H), 0.80-1.00 (m, CH₃, 6H); MS(EI): m/e
5 493.4 (M⁺: C₂₆H₃₉N₃); HRMS: 393.3166 (Calcd: 393.3160); TLC: R_f=0.52 (SiO₂, CHCl₃/MeOH = 10/1); HPLC: 99.7%; mp: oil.

Example 72: N-(4-sec-Butylphenyl)-N-(4-t-butylbenzyl)-N'-pyrrolidinyguanidine•HCl

10 ¹H NMR (CD₃OD): δ ppm 7.50-7.15 (m, ArH, 8H), 4.95 (s, CH₂, 2H), 3.25-3.15 (m, CH₂, 4H), 2.65-2.50 (m, CH, 1H), 1.80-1.90 (m, CH₂, 4H), 1.68-1.55 (m, CH₂, 2H), 1.31 (s, t-butyl, 9H), 1.23 (d, CH₃, J=7.0Hz, 3H), 0.79 (t, CH₃, J=7.4Hz, 3H); HRMS(EI): m/e 391.2991 (Calcd. 391.2987 for C₂₆H₃₇N₃); TLC: R_f=0.4 (SiO₂, CHCl₃/MeOH = 10/1); HPLC: 96% pure;
15 mp: 98-100°C.

Example 73: N-(4-sec-Butylphenyl)-N-(4-t-butylbenzyl)-N'-(4-thiomorpholinyl)guanidine•HCl

20 ¹H NMR (CD₃OD): δ ppm 7.50-7.18 (m, ArH, 8H), 4.97 (s, CH₂, 2H), 3.15-3.05 (m, CH₂, 4H), 2.70-2.60 (m, CH, 1H), 2.40-2.25 (m, CH₂, 4H), 1.68-1.56 (m, CH₂, 2H), 1.31 (s, t-butyl, 9H), 1.23 (d, CH₃, J=7.0Hz, 3H), 0.79 (t, CH₃, J=7.4Hz, 3H); HRMS(EI): m/e 423.2722 (Calcd. 423.2708 for C₂₆H₃₇S₁N₃); Anal. (C₂₆H₃₇S₁N₃•HCl); Calcd (%): C, 67.87, H, 8.32, N, 9.13; Found (%): C, 67.85, H, 8.11, N, 9.36; TLC: R_f=0.4 (SiO₂,
25 CHCl₃/MeOH = 10/1); HPLC: 98.6% pure; mp: 70-72°C.

Example 74: N-(4-sec-Butylphenyl)-N-(4-tert-butylbenzyl)-N'-piperidinyguanidine•HCl

30 ¹H NMR (CD₃OD): δ ppm 7.50-7.16 (m, ArH, 8H), 4.96 (s, CH₂, 2H), 2.62 (m, CH, 1H), 1.70-1.40 (m, 6CH₂, 12H), 1.31 (s, CH₃, 9H), 1.22 (d, CH₃, J=6.93Hz, 3H), 0.79 (t, CH₃, J=7.39Hz, 3H); HRMS: 405.3138

(Calcd: 405.3144 for $C_{27}H_{39}N_3$); HPLC: 99.80%; TLC: $R_f = 0.47$ (SiO_2 , $CH_2Cl_2/MeOH = 10/1$); mp: 217-218°C.

5 **Example 75:** **N-(4-sec-Butylphenyl)-N-(4-tert-butylbenzyl)-N'-(4-morpholinyl)guanidine•HCl**

1H NMR (CD_3OD): δ ppm 7.21-7.46 (m, ArH, 8H), 4.99 (s, CH_2 , 2H), 3.41-3.45 (m, $2OCH_2$, 4H), 3.30-3.31 (m, NCH_2 , 4H), 2.55-2.70 (m, CH, 1H), 1.59-1.62 (m, CH_2 , 2H), 1.31 (s, CMe_3 , 9H), 1.23 (d, CH_3 , $J = 7.0Hz$, 3H), 0.79 (t, CH_3 , $J = 7.4Hz$, 3H); MS(EI): m/e 408 (M^+ : $C_{26}H_{37}N_3O$); Anal. (C, H, N; $C_{26}H_{37}N_3O \cdot HCl \cdot 0.75 H_2O$); Calcd. (%): C, 68.24, H, 8.70, N, 9.18; Found (%): C, 67.76, H, 8.56, N, 9.08; TLC: $R_f = 0.43$ (SiO_2 , $CHCl_3/MeOH = 10/1$); HPLC: 98.57%; mp: 220-221°C.

15 **Example 76:** **N-(4-sec-Butylphenyl)-N-(4-tert-butylbenzyl)-N'-(4-propylpiperidiny)guanidine•HCl**

1H NMR (CD_3OD): δ ppm 7.16-7.46 (m, ArH, 8H), 4.94 (s, CH_2 , 2H), 3.70-3.90 (m, ring-H, 2H), 2.80-2.95 (m, ring-H, 3H), 2.55-2.70 (m, CH, 1H), 0.59-2.70 (m, ring-4H; CMe_3 , 9H; $3CH_3$, 9H; $3CH_2$, 6H); MS(EI): m/e 448 (M^+ : $C_{30}H_{45}N_3$); Anal. (C, H, N; $C_{30}H_{45}N_3 \cdot HCl \cdot 1.25H_2O$); Calcd. (%): C, 71.11, H, 9.65, N, 8.29; Found (%): C, 71.13, H, 9.50, N, 8.53; TLC: $R_f = 0.64$ (SiO_2 , $CHCl_3/MeOH = 10/1$); HPLC: 93.1%; mp: 109-110°C.

Example 77: **N-(4-Butoxyphenyl)-N-(4-tert-butylbenzyl)-N'-(4-piperidiny)guanidine•HCl**

25 1H NMR (CD_3OD): δ ppm 6.94-7.45 (m, ArH, 8H), 4.91 (s, CH_2 , 2H), 3.97 (t, OCH_2 , $J = 6.36Hz$, 2H), 1.70-1.85 (m, CH_2 , 2H), 1.25-1.60 (m, ring-H, 10H; CMe_3 , 9H; CH_2 , 2H), 0.98 (t, CH_3 , $J = 7.4Hz$, 3H); MS(EI): m/e 422.3 (M^+ : $C_{27}H_{39}N_3O$); HRMS: 421.3110 (Calcd.: 421.3093); TLC: $R_f = 0.28$ (SiO_2 , $CHCl_3/MeOH = 10/1$); HPLC: 93.7%; mp: 99-100°C.

Example 78: N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-benzylpiperidiny)guanidine•mesylate

¹H NMR (CD₃OD): δ ppm 7.03-7.46 (m, ArH, 13H), 4.94 (s, CH₂, 2H), 3.70-3.90 (m, ring-H, 3H), 2.77-2.95 (m, ring-H, 3H), 2.70 (s, CH₃SO₃H, 3H), 2.60-2.70 (m, CH, 1H), 2.40 (d, CH₂, J = 7.15Hz, 2H), 1.55-1.75 (m, CH₂, 2H), 1.40-1.55 (m, ring-H, 3H), 1.31 (m, CMe₃, 9H), 1.24 (d, CH₃, J = 6.93Hz, 3H), 0.82 (t, CH₃, J = 7.4Hz, 3H); MS(EI): m/e 495.3 (M⁺: C₃₄H₄₅N₃); HRMS: 495.3604 (Calcd.: 495.3613); TLC: R_f = 0.55 (SiO₂, CHCl₃/MeOH = 10/1); HPLC: 94.7%; mp: 85-86°C.

Example 79: N-(4-Benzyloxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-morpholinyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.00-7.50 (m, ArH, 13H), 5.10 (s, OCH₂, 2H), 4.94 (s, CH₂, 2H), 3.41-3.50 (m, 2OCH₂, 4H), 3.25-3.35 (m, NCH₂, 4H), 1.31 (s, CMe₃, 9H); MS(EI): m/e 457.4 (M⁺: C₂₉H₃₅N₃O₂): Anal. (C, H, N; C₂₉H₃₅N₃O₂•HCl); Calcd. (%): C, 70.50, H, 7.34, N, 8.50; Found (%): C, 70.29, H, 7.15, N, 8.36; TLC: R_f = 0.50 (SiO₂, CHCl₃/MeOH = 10/1); HPLC: 95%; mp: 75-77°C.

Example 80: N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(1,2,3,4-tetrahydroisoquinolinyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 6.95-7.50 (m, ArH, 12H), 5.01 (s, benzylic-CH₂, 2H), 4.46 (s, benzylic-CH₂N, 2H), 3.60 (t, CH₂, J = 6Hz, 2H), 2.55-2.70 (m, CH, 1H), 2.48 (t, CH₂, J = 6.0Hz, 2H), 1.50-1.65 (m, CH₂, 2H), 1.30 (s, CMe₃, 9H), 1.20 (d, CH₃, J = 6.93Hz, 3H), 0.77 (t, CH₃, J = 7.4Hz, 3H); MS(EI): m/e 454.4 (M⁺: C₃₁H₃₉N₃); HRMS: 453.3165 (Calcd.: 453.3144); TLC: R_f = 0.51 (SiO₂, CHCl₃/MeOH = 10/1); HPLC: 100%; mp: 217-218°C.

Example 81: N-(3-Butoxy-4-methoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-morpholinyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 6.55-7.50 (m, ArH, 7H), 4.93 (s, CH₂, 2H), 3.80-3.90 (m, OCH₂+OCH₃, 5H), 3.45-3.55 (m, 2OCH₂, 4H), 3.30-3.40 (m, NCH₂, 4H), 1.65-1.80 (m, CH₂, 2H), 1.40-1.55 (m, CH₂, 2H), 1.31 (s, CMe₃, 9H), 0.97 (t, CH₃, J = 7.4Hz, 3H); MS(EI): m/e 453.4 (M⁺: C₂₇H₃₉N₃O₃); HRMS: 453.3010 (Calcd.: 453.2991); TLC: R_f = 0.39 (SiO₂, CHCl₃/MeOH = 10/1); HPLC: 98%; mp: 197-199°C.

Example 82: N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(3,5-dimethyl-4-morpholinyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.20-7.50 (m, ArH, 8H), 4.99 (s, CH₂, 2H), 3.55-3.70 (m, 2OCH, 2H), 3.15-3.25 (m, NCH₂, 2H), 2.50-2.70 (m, CH+NCH₂, 3H), 1.55-1.70 (m, CH₂, 2H), 1.30 (s, CMe₃, 9H), 1.23 (d, CH₃, J = 6.93Hz, 3H), 0.99 (m, 2CH₃, 6H), 0.79 (m, CH₃, 3H); MS(EI): m/e 435.5 (M⁺: C₂₈H₄₁N₃O); TLC: R_f = 0.51 (SiO₂, CHCl₃/MeOH = 10/1); HPLC: 98.7%; mp: 98-99°C.

Example 83: N-(4-*tert*-Butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-*sec*-butylphenyl)-N'-(methyl)guanidine•HCl

¹H NMR (CDCl₃): δ ppm 7.42-6.60 (m, ArH, 12H), 4.92 (s, CH₂, 2H), 3.56 (s, CH₃, 3H), 2.52 (m, CH, 1H), 1.54 (m, CH₂, 2H), 1.33 (s, CH₃, 9H), 1.26 (s, CH₃, 9H), 1.17 (d, CH₃, J = 7.14Hz, 3H), 0.80 (t, CH₃, J = 7.28Hz, 3H); MS(EI): m/e 483.5 (M⁺: C₃₃H₄₅N₃); Anal. (C, H, N: C₃₃H₄₅N₃•HCl); Calcd. (%): C, 76.19, H, 8.91, N, 8.08; Found (%): C, 76.12, H, 8.87, N, 8.03; TLC: R_f = 0.53 (SiO₂, CH₂Cl₂/MeOH = 10/1); mp: 111-112°C.

Example 84: N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-*sec*-butylphenyl)-N'-(methyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.50-6.70 (m, ArH, 12H), 4.88 (s, CH₂, 2H), 3.35 (s, CH₃, 3H), 2.50 (m, 2CH, 2H), 1.54 (m, 2CH₂, 4H), 1.30 (s, CH₃, 9H), 1.15 (m, 2CH₃, 6H), 0.80 (m, 2CH₃, 6H); MS(EI): m/e 483.5

(M^+ : $C_{33}H_{45}N_3$); Anal. (C, H, N; $C_{33}H_{45}N_3 \cdot HCl$); Calcd. (%): C, 76.19, H, 8.91, N, 8.08; Found (%): C, 75.94, H, 9.07, N, 7.86; TLC: $R_f = 0.30$ (SiO_2 , $CH_2Cl_2/MeOH = 10/1$); mp: 106-107°C.

5 **Example 85:** **N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(phenyl)guanidine•HCl**

10 1H NMR (CD_3OD): δ ppm 7.50-7.23 (m, ArH, 13H), 5.03 (s, CH_2 , 2H), 2.62 (m, CH, 1H), 1.60 (m, CH_2 , 2H), 1.30 (s, CH_3 , 9H), 1.15 (d, CH_3 , $J = 6.96$ Hz, 3H), 0.80 (t, CH_3 , $J = 7.37$ Hz, 3H); HRMS: 413.2818 (Calcd: 413.2831 for $C_{28}H_{35}N_3$); HPLC: 96.80%; TLC: $R_f = 0.58$ (SiO_2 , $CH_2Cl_2/MeOH = 10/1$); mp: 204-205°C.

15 **Example 86:** **N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-chlorophenyl)guanidine•HCl**

20 1H NMR (CD_3OD): δ ppm 7.22-7.46 (m, ArH, 12H), 5.02 (s, CH_2 , 2H), 2.55-2.70 (m, CH, 1H), 1.59-1.62 (m, CH_2 , 2H), 1.30 (s, CMe_3 , 9H), 1.20 (d, CH_3 , $J = 6.9$ Hz, 3H), 0.80 (t, CH_3 , $J = 7.4$ Hz, 3H); MS(EI): m/e 447.7 (M^+ : $C_{28}H_{34}N_3Cl$); HRMS: 447.2412 (Calcd. 447.2441); Anal. (C, H, N; $C_{28}H_{34}N_3Cl \cdot HCl \cdot 0.5H_2O$); Calcd. (%): C, 68.15, H, 7.35, N, 8.51; Found (%): C, 68.15, H, 7.44, N, 8.50; TLC: $R_f = 0.40$ (SiO_2 , $CHCl_3/MeOH = 10/1$); mp: 98-99°C.

25 **Example 87:** **N-(4-Butoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(phenyl)guanidine•HCl**

30 1H NMR (CD_3OD): δ ppm 6.94-7.47 (m, ArH, 13H), 4.99 (s, CH_2 , 2H), 3.97 (t, OCH_2 , $J = 6.43$ Hz, 2H), 2.55-2.70 (m, CH, 1H), 1.65-1.80 (m, CH_2 , 2H), 1.40-1.55 (m, CH_2 , 2H), 1.31 (s, CMe_3 , 9H), 0.96 (t, CH_3 , $J = 7.4$ Hz, 3H); MS(EI): m/e 430 (M^+ : $C_{28}H_{35}N_3O$); Anal. (C, H, N; $C_{28}H_{35}N_3O \cdot HCl \cdot 0.75H_2O$); Calcd. (%): C, 70.13, H, 7.88, N, 8.76; Found (%): C, 70.29, H, 7.53, N, 9.12; TLC: $R_f = 0.26$ (SiO_2 , $CHCl_3/MeOH = 10/1$); HPLC: 97.1%; mp: 95-96°C.

Example 88: N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(phenyl)-N'-methylguanidine•HCl

¹H NMR (CD₃OD): δ ppm 6.79-7.44 (m, ArH, 13H), 4.88 (s, CH₂, 2H), 3.35 (s, CH₃, 3H), 2.45-2.60 (m, CH, 1H), 1.45-1.62 (m, CH₂, 2H),
5 1.30 (s, CMe₃, 9H), 1.15 (d, CH₃, J = 6.9Hz, 3H), 0.77 (t, CH₃, J = 7.4Hz, 3H); MS(EI): m/e 427.1 (M⁺: C₂₉H₃₇N₃); HRMS: 427.2987 (Calcd.: 427.2954); Anal. (C, H, N; C₂₉H₃₇N₃•HCl); Calcd. (%): C, 75.05, H, 8.25, N, 9.05; Found (%): C, 74.91, H, 8.12, N, 9.11; TLC: R_f = 0.56 (SiO₂, CHCl₃/MeOH = 10/1); mp: 80-82°C.

Example 89: N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(3,4-dichlorophenyl)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.21-7.57 (m, ArH, 11H), 5.04 (s, CH₂, 2H), 2.55-2.70 (m, CH, 1H), 1.50-1.65 (m, CH₂, 2H), 1.30 (s, CMe₃, 9H),
15 1.20 (d, CH₃, J = 6.9Hz, 3H), 0.79 (t, CH₃, J = 7.4Hz, 3H); MS(EI): m/e 482.1 (M⁺: C₂₈H₃₃N₃Cl₂); HRMS: 481.2025 (Calcd.: 481.2052); Anal. (C, H, N; C₂₈H₃₃N₃Cl₂•HCl); Calcd. (%): C, 64.80, H, 6.60, N, 8.10; Found (%): C, 64.78, H, 6.58, N, 8.14; TLC: R_f = 0.71 (SiO₂, CHCl₃/MeOH = 10/1); mp: 168-169°C.

Example 90: N-(4-Hexylphenyl)-N-(4-*tert*-hexylbenzyl)-N'-phenylguanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.15-7.70 (m, ArH, 13H), 5.01 (s, CH₂, 2H), 2.55-2.70 (m, 2CH₂, 4H), 1.50-1.70 (m, 2CH₂, 4H), 1.25-1.40 (m,
25 6CH₂, 12H), 0.80-1.00 (m, CH₃, 6H); MS(EI): m/e 469.4 (M⁺: C₃₂H₄₃N₃); HRMS: 469.3476 (Calcd.: 469.3457); TLC: R_f = 0.56 (SiO₂, CHCl₃/MeOH = 10/1); HPLC: 100%; mp: oil.

Example 91: N-(4-*sec*-Butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-benzyloxyphenyl)guanidine•mesylate

¹H NMR (CD₃OD): δ ppm 7.05-7.50 (m, ArH, 17H), 5.11 (s, OCH₂, 2H), 5.00 (s, CH₂, 2H), 2.69 (s, CH₃SO₃H, 3H), 2.55-2.70 (m, CH, 1H),

1.50-1.70 (m, CH₂, 2H), 1.30 (s, CMe₃, 9H), 1.20 (d, CH₃, J = 6.9Hz, 3H),
0.80 (t, CH₃, J = 7.4Hz, 3H); MS(EI): m/e 519.5 (M⁺: C₃₅H₄₁N₃O); Anal. (C,
H, N; C₃₅H₄₁N₃O•CH₃SO₃H); Calcd. (%): C, 70.21, H, 7.37, N, 6.82; Found
(%): C, 70.12, H, 7.16, N, 6.69; TLC: R_f = 0.53 (SiO₂, CHCl₃/MeOH =
5 10/1); HPLC: 96.9%; mp: 142-143°C.

Example 92: N,N'-Bis-(4-*tert*-butylphenyl)-N,N'-dimethylguanidine•HBr

White solid; mp: 175-177°C; TLC (CH₂Cl₂:MeOH; 10:1); R_f = 0.39;
¹H NMR (CD₃OD): 7.21-7.18 (d, J = 8.52Hz, 4H, ArH), 6.80-6.78 (d,
10 J = 8.51Hz, 4H, ArH), 3.35 (s, 6H, -CH₃), 1.26 (s, 18H,
-C(CH₃)₃); Anal. Calcd. for C₂₃H₃₃N₃•HCl (388.00); C, 71.20, H, 8.83, N,
10.83; Found: C, 70.88, H, 8.61, N, 10.75.

Example 93: N-(4-Benzyloxyphenyl)-N'-(4-*tert*-butylphenyl)guanidine•HCl

White solid; mp: 143-144°C; TLC (CH₂Cl₂:MeOH; 15:1); R_f = 0.35;
¹H NMR (CD₃OD): 7.547-7.089 (m, 13H, ArH), 5.135 (s, CH₂, 2H), 1.346
(s, *t*-butyl, 9H); Anal. Calcd. for C₂₄H₂₈N₃ClO (409.96): C, 70.32, H, 6.88,
N, 10.25; Found: C, 70.49, H, 6.94, N, 10.09.

Example 94: N,N'-Bis-(3-(1'-methyl-2'-phenyl)ethyl)guanidine•HCl

White solid; mp: 93-95°C; TLC (CH₂Cl₂:MeOH; 15:1); R_f = 0.41; ¹H
NMR (CD₃OD): 7.395-7.344 (t, J = 8.69Hz, 2H, ArH), 7.225-7.077 (m,
16H, ArH), 3.092-3.045 (m, CH, 2H), 2.952-2.827 (m, CH₂, 4H), 1.273-
1.250 (d, J = 6.87Hz, 6H); Anal. Calcd. for C₃₁H₃₄N₃Br•1/2H₂O (537.55): C,
25 69.27, H, 6.51, N, 7.81; Found: C, 69.61 H, 6.63, N, 7.87.

Example 95: N-Methyl-N-4-benzyloxyphenyl-N'-(4-*tert*-butylphenyl)guanidine•mesylate

White solid; mp: 230-232°C; TLC (AcOEt:MeOH; 10:1); R_f = 0.45; ¹H
NMR: 7.421-6.961 (m, ArH, 13H), 5.059 (s, OCH₂, 2H), 3.413 (s, N-CH₃,

3H), 2.770 (s, CH₃, 3H), 1.286 (s, t-butyl, 9H); Anal. Calcd. for C₂₅H₃₃N₃O₃S•1/2H₂O (492.63: C, 63.33, H, 6.90, N, 8.53; Found: C, 63.48, H, 6.40, N, 8.48.

5 **Example 96:** **N,N'-Bis-(4-Hexylphenyl)guanidine•mesylate**

White solid; mp: 92-94°C; TLC (AcOEt:MeOH; 10:1): R_f=0.49; ¹H NMR: 7.265-7.172 (m, ArH, 8H), 2.823 (s, CH₃, 3H), 2.627-2.575 (t, CH₂, J=7.695Hz, 4H), 1.583-1.558 (m, CH₂, 4H), 1.336-1.261 (m, CH₂CH₂, 8H), 0.905-0.861 (t, CH₃, J=6.59Hz, 6H); Anal. Calcd. for C₂₆H₄₁N₃O₃S (475.69): C, 65.65, H, 8.69, N, 8.83; Found: C, 65.17, H, 8.63, N, 8.70.

15 **Example 97:** **N-(3-(1-(4'-Ethoxy)benzyl)phenethyl)-N'-(4-tert-butylphenyl)guanidine•mesylate**
 (3-(4-(CH₃CH₂O)C₆H₄CH₂)C₆H₄)(CH₃)CHNHC(=NH)NH(4-(CH₃)₃C)C₆H₄ mesylate)

White solid; mp: 83-85°C; TLC (AcOEt:MeOH; 10:1): R_f=0.49; ¹H NMR: 7.465-6.680 (m, ArH, 12H), 3.957-3.888 (m, CH₂, 2H), 2.961 (m, CH, 1H), 2.836 (s, CH₃, 2H), 2.825-2.727 (m, CH₂, 2H), 1.322 (s, tert-butyl, 9H), 1.361-1.268 (m, CH₂, CH₃, 5H); HRMS: 429.2749 (429.2780 Calcd. for C₂₈H₃₅ON₃).

25 **Example 98:** **N-(4-Benzyloxyphenyl)-N'-methyl-N'-(4-tert-butylphenyl)guanidine•mesylate**

White solid; mp: 100-101°C; TLC (AcOEt:MeOH; 10:1): R_f=0.46; ¹H NMR: 7.457-7.371 (m, ArH, 7H), 7.199-7.170 (d, ArH, J=8.58Hz, 2H), 7.081-7.052 (d, ArH, J=8.85Hz, 2H), 6.924-6.894 (d, ArH, J=9.01Hz, 2H), 5.034 (s, CH₂, 2H), 3.456 (s, CH₃, 3H), 3.456 (s, CH₃, 3H), 1.323 (s, tert-butyl, 9H); HRMS: 387.2292 (387.2311 Calcd. for C₂₅H₂₉ON₃).

Example 99: N-(3-(4-*tert*-Butylbenzyloxy)phenyl)-N'-(4-*tert*-butylphenyl)guanidine•mesylate

White solid; mp: 88-92°C; TLC (AcOEt:MeOH; 10:1): R_f = 0.48; ^1H NMR (CDCl_3): 7.479-7.199 (m, ArH, 10H), 6.909-6.878 (d, ArH, J = 8.94Hz, 2H), 5.071 (s, CH_2 , 2H), 2.856 (s, $-\text{CH}_3$, integration is 1.5H instead of 3H), 1.329 (s, *t*-butyl, 9H), 1.292 (s, *t*-butyl, 9H); HRMS: 429.2783 (429.2780 Calcd. for $\text{C}_{30}\text{H}_{31}\text{N}_3$).

Example 100: N-(3-(1'-Benzylbutyl)phenyl)-N'-(4-*tert*-butylphenyl)guanidine•mesylate

White solid; mp: 70-73°C; TLC (AcOEt:MeOH; 10:1): R_f = 0.78; ^1H NMR (CDCl_3): 7.494-6.759 (m, ArH, 17H), 3.014-2.691 (m, $-\text{CH}_2$, $-\text{CH}$, $-\text{CH}_3$, 6H), 1.755-1.168 (m, CH_2CH_2 , 4H), 1.331 (s, *t*-butyl, 9H), 0.896-0.847 (t, CH_3 , J = 7.28Hz, 3H); HRMS: 413.2840 (413.2831 Calcd. for $\text{C}_{28}\text{H}_{35}\text{N}_3$).

Example 101: N,N'-Bis-(4-butylphenyl)-N-methylguanidine•HCl

White solid; mp: 137-138°C; TLC (CHCl_3 :MeOH; 10:1): R_f = 0.40; ^1H NMR (CDCl_3): 7.234-7.026 (m, ArH, 8H), 3.618 (s, CH_3 , 3H), 2.628-2.535 (m, CH_2 , 4H), 1.600-1.493 (m, CH_2 , 4H), 1.382-1.280 (m, CH_2 , 4H), 0.963-0.900 (m, CH_3 , 6H); Anal. Calcd. for $\text{C}_{22}\text{H}_{32}\text{N}_3\text{Cl}$: C, 70.66, H, 8.62, N, 11.24; Found: C, 70.47, H, 8.04, N, 11.31.

Example 102: N,N'-Bis-(4-*tert*-butylphenyl)-N,N'-dimethylguanidine•HCl

White solid; mp: 134°C; TLC (CHCl_3 :MeOH; 10:1): R_f = 0.33; ^1H NMR (CDCl_3): 9.460 (s, 2H), 6.923-6.895 (d, ArH, J = 6.37Hz, 4H), 6.668-6.640 (d, ArH, J = 8.45Hz, 4H), 3.557 (s, $-\text{CH}_3$, 3H), 2.529-2.479 (t, CH_2 , J = 7.525Hz), 1.570-1.470 (m, CH_2 , 4H), 1.358-1.283 (m, CH_2 , 4H), 0.972-0.924 (t, CH_3 , J = 7.265Hz, 6H); Anal. Calcd. for $\text{C}_{23}\text{H}_{34}\text{N}_3\text{Cl} \cdot 1/2\text{H}_2\text{O}$: C, 69.52, H, 8.82, N, 10.58; Found: C, 69.38, H, 8.52, N, 10.56.

Example 103: **N-(3-Naphthylmethylenedioxyphenyl)-N'-(4-*tert*-butylphenyl)guanidine•mesylate**
(3-(C₁₀H₇CH₂O)C₆H₄ NHC(=NH)NH(4-(CH₃)₃C)C₆H₅ mesylate)

5 White solid; mp: 138-146°C; TLC (CHCl₃:MeOH; 10:1): R_f=0.52; ¹H NMR (CDCl₃): 7.861-7.803 (m, ArH, 4H), 7.516-7.113 (m, ArH, 8H), 6.936-6.881 (m, ArH, 3H), 5.222 (s, CH₂, 2H), 2.814 (s, CH₃, 3H), 1.291 (s, *t*-butyl, 9H); Anal. Calcd. for C₂₉H₃₃N₃O₄S: C, 67.03, H, 6.40, N, 8.09; Found: C, 67.26, H, 6.64, N, 8.29.

10

Example 104: **N-(4-Benzyloxyphenyl)-N'-(4-butylphenyl)guanidine•HCl**

 White solid; mp: 112°C; TLC (CHCl₃:MeOH; 10:1): R_f=0.50; ¹H NMR (CDCl₃): 7.403-7.206 (m, ArH, 11H), 7.009-7.039 (d, J=8.91Hz, ArH, 2H), 5.068 (s, Ar-CH₂, 2H), 2.638-2.587 (t, J=7.42Hz, CH₂, 2H),
15 1.576-1.560 (m, CH₂, 2H), 1.371-1.321 (m, CH₂, 2H), 0.938-0.890 (t, CH₃, J=7.36Hz, 3H); Anal. Calcd. for C₂₄H₂₈ClN₃O: C, 70.32, H, 6.88, N, 10.25; Found: C, 70.42, H, 7.00, N, 10.07.

Example 105: **N,N'-Bis-(4-butylphenyl)-N-butylguanidine•HCl**

20 White solid; mp: 118-119°C; TLC (CHCl₃:MeOH; 10:1): R_f=0.43; ¹H NMR: 7.235, 7.207, 7.033, 7.005 (q, ArH, 4H), 7.127-7.095 (d, J=9.63Hz, ArH, 4H); Anal. Calcd. for C₂₇H₃₉N₃O₄ (469.62): C, 69.05, H, 8.37, N, 8.95; Found: C, 69.25, H, 8.38, N, 9.03.

25 **Example 106:** **N-3-(Benzyloxymethyl)phenyl-N'-(4-*tert*-butylphenyl)guanidine•mesylate**

 White solid; mp: 66-68°C; TLC (CHCl₃:MeOH; 10:1): R_f=0.56; ¹H NMR (CDCl₃): 7.456-7.170 (m, ArH, 13H), 4.580 (s, CH₂, 2H), 4.529 (s, CH₂, 2H), 2.821 (s, CH₃, 3H), 1.303 (s, C(CH₃)₃, 9H); HRMS: 387.2299
30 (387.2229 calculated for C₂₅H₂₉N₂O₂).

Example 107: N-(3,4-Bis-butyloxyphenyl)-N'-(4-*tert*-butylphenyl)guanidine•oxalate

White solid; mp: 100-101°C; TLC (CHCl₃:MeOH; 10:1): R_f=0.58; ¹H NMR (CD₃OD): 7.5251-7.5027 (m, ArH, 2H), 7.2692-7.2402 (m, ArH, 2H), 7.0079-6.8720 (m, ArH, 3H), 4.0306-3.9880 (t, OCH₂, 4H), 1.7990-1.7404 (m, CH₂, 4H), 1.5580-1.5296 (m, CH₂, 4H), 1.3275 (s, C(CH₃)₃, 9H), 1.0243-1.9558 (t, CH₃, 6H); HRMS: 411.2886 (411.2886 calculated for C₂₅H₃₇N₃O₂).

Example 108: N-(3-Benzoyloxy)phenyl-N'-(4-*tert*-butylphenyl)guanidine•mesylate

White solid; mp: 140°C; TLC (CHCl₃:MeOH; 10:1): R_f=0.50; ¹H NMR (CDCl₃): 7.483-7.203 (m, 9H, ArH), 6.920-6.910 (m, 4H, ArH), 5.080 (s, CH₂, 2H), 2.849 (s, CH₃, 3H), 1.585-1.511 (m, NHs), 1.320 (s, C(CH₃)₃, 9H); HRMS: 373.2158 (373.2154 calculated for C₂₄H₂₇N₃O).

Example 109: N,N'-Bis-((3-butoxy-4-methoxy)phenyl)guanidine•HBr

White solid; mp: 174°C; TLC (CHCl₃:MeOH; 10:1): R_f=0.43; ¹H NMR (CDCl₃): 6.924-6.852 (m, 6H, ArH), 4.028-3.984 (t, CH₂, J=6.59Hz, 4H), 3.892 (s, CH₃, 6H), 1.872-1.798 (m, CH₂, 4H), 1.594-1.472 (m, -CH₂, 4H), 1.021-0.972 (t, CH₃, J=7.28Hz, 6H); HRMS: 415.2470 (415.2471 calculated for C₂₃H₃₃N₃O₄).

Example 110: N-(4-Benzoyloxyphenyl)-N-methyl-N'-(4-butylphenyl)guanidine•mesylate

White solid; mp: 178.4-178.8°C; TLC (CHCl₃:MeOH; 10:1): R_f=0.34; ¹H NMR (CDCl₃): 7.390-6.919 (m, ArH, 13H), 5.029 (s, Ar-CH₂, 2H), 3.389 (s, CH₃, 3H), 2.717 (s, CH₃, 3H), 2.552-2.501 (t, J=7.48Hz, CH₂, 2H), 1.544-1.493 (m, CH₂, 2H), 1.328-1.254 (m, CH₂, 2H), 0.916-0.867 (t, CH₃, J=7.33Hz, 3H); HRMS: 387.2311 (387.2285 calculated for C₂₅H₂₉N₃O).

Example 111: N,N'-Bis-(6-tetralinyl)guanidine•HBr

White solid; mp: too fluffy to measure; TLC (CHCl₃:MeOH; 10:1):
R_f=0.50; ¹H NMR (CDCl₃): 7.273-7.002 (m, ArH, 6H), 2.765 (s, 2CH₂,
4H), 1.824-1.780 (p, 2CH₂, 4H); Anal. Calcd. for C₂₁H₂₈N₃Br (400.36): C,
5 63.00, H, 6.55, N, 10.50; Found: C, 62.85, H, 6.62, N, 10.53.

Example 112: N-(6-Tetralinyl)-N'-(4-*tert*-butylphenyl)guanidine•mesylate

Colorless crystal; mp: 53-55 °C; TLC (CHCl₃:MeOH; 10:1): R_f=0.45;
¹H NMR (CDCl₃): 7.240-6.967 (m, 7H, ArH), 2.798 (s, CH₃, 3H), 2.736 (s,
10 CH₂, 4H), 2.628-2.577 (t, J=7.47Hz, CH₂, 2H), 1.790-1.770 (t, 2CH₂,
4H), 1.597-1.546 (m, CH₂, 2H), 1.374-1.300 (m, CH₂, 2H), 0.949-0.901
(t, CH₃, J=7.30Hz, 3H); HRMS: 321.2184 (321.2205 calculated for
C₂₁H₂₇N₃).

Example 113: N-(5-Acenaphthyl)-N'-(6-benzothiazolyl)guanidine•HCl

Light green solid; mp: °C; TLC (CH₂Cl₂:MeOH; 15:1): R_f=0.25; ¹H
NMR (CD₃OD): 7.639-7.289 (m, ArH, 9H), 3.443-3.403 (m, CH₂CH₂, 4H),
2.086-1.756 (m, CH's, 13H); Anal. Calcd for C₂₉H₃₂N₃Cl (458.08): C,
15 76.04, H, 7.04, N, 9.17; Found: C, 75.97, H, 6.88, N, 9.06.

20

Example 114: N-(5-Acenaphthyl)-N'-(6-N-benzylindolinyl)guanidine•mesylate

White solid; mp: 115-125 °C; TLC (AcOEt:MeOH; 10:1): R_f=0.27; ¹H
NMR: 7.709-7.260 (m, ArH, 13H), 3.469-3.377 (s, CH₂CH₂, 4H), 2.842 (s,
25 CH₃, 3H); Anal. Calcd. for C₂₉H₃₀N₄O₃S (514.64): C, 67.68, H, 5.88, N,
10.89; Found: C, 67.51, H, 5.58, N, 10.88.

Example 115: N-(5-Acenaphthyl)-N'-(4-benzo-2,1,3-thiadiazole)guanidine•HCl

¹H NMR (CD₃OD): δ ppm 7.33-8.05 (m, ArH, 8H), 3.43-3.50 (m, 4H,
30 CH₂); MS(EI): m/e 345.1 (M⁺: C₁₉H₁₅N₅S); Anal. (C, H, N;

$C_{19}H_{15}N_5S \cdot HCl \cdot 1/2 H_2O$): Calcd. (%): C, 58.45, H, 4.39, N, 17.95; Found (%): C, 58.62, H, 4.29, N, 17.47; TLC: $R_f = 0.13$ (SiO_2 , $CHCl_3/MeOH$) = 10:1; mp: 173-174°C.

5 **Example 116:** **N-(5-Acenaphthyl)-N'-[4-(6-methyl-benzothiazole)phenylguanidine]•HCl**

1H NMR (CD_3OD): δ ppm 8.15-8.17 (m, ArH, 2H), 7.36-7.91 (m, ArH, 10H), 3.40-3.50 (m, CH_2 , 4H), 2.50 (s, CH_3 , 1H); MS(EI): m/e 434.1 (M^+ : $C_{27}H_{22}N_4S$); Anal. (C, H, N; $C_{27}H_{22}N_4S \cdot HCl$): Calcd. (%): C, 68.85, H, 4.92, N, 11.89; Found (%): C, 68.66, H, 4.91, N, 11.86; TLC: $R_f = 0.23$ (SiO_2 , $CHCl_3/MeOH$) = 10:1; mp: 244.5-246°C.

Example 117: **N-(5-Acenaphthyl)-N'-(1-benz[cd]indoliny)guanidine•HCl**

1H NMR (CD_3OD): δ ppm 7.90-7.40 (m, ArH, 11H), 5.66 (s, NCH_2 , 2H), 3.49 (m, $2CH_2$, 4H); MS(EI): m/e 349.2 (M^+ : $C_{24}H_{19}N_3$); Anal. (C, H, N; $C_{24}H_{19}N_3 \cdot HCl$): Calcd. (%): C, 74.70, H, 5.22, N, 10.89; Found (%): C, 74.61, H, 5.08, N, 10.63; TLC: $R_f = 0.51$ (SiO_2 , $CH_2Cl_2/MeOH$ = 10/1); mp: 245-246°C.

20 **Example 118:** **N-(5-Acenaphthyl)-N'-(6-benz[cd]indo-2[1H]-one)guanidine•HCl**

1H NMR (CD_3OD): δ ppm 8.25-7.04 (m, ArH, 10H), 3.45-3.43 (m, CH_2 , 4H); MS(EI): m/e 378.3 (M^+ : $C_{24}H_{18}ON_4$); Anal. (C, H, N; $C_{24}H_{18}ON_4 \cdot HCl$): Calcd. (%): C, 69.48, H, 4.62, N, 13.50; Found (%): C, 69.36, H, 4.72, N, 13.27; TLC: $R_f = 0.26$ (SiO_2 , $CH_2Cl_2/MeOH$ = 10/1); mp: 327-328°C.

Example 119: **N,N'-Bis(6-benz[cd]indoliny-2[1H]-one)guanidine•HBr**

1H NMR (DMSO): δ ppm 11.00-7.00 (m, ArH, 10H); MS(EI): m/e 393.2 (M^+ : $C_{23}H_{15}N_5O_2$); Anal. (C, H, N; $C_{23}H_{15}N_5O_2 \cdot HCl$): Calcd. (%): C,

58.24, H, 3.40, N, 14.77; Found (%): C, 58.25, H, 3.25, N, 14.80; TLC: $R_f = 0.25$ (SiO_2 , $\text{CH}_2\text{Cl}_2/\text{MeOH} = 10/1$); mp: 390-391 °C.

5 **Example 120:** **N-(4-Butoxyphenyl)-N'-(4-chlorophenylethyl)guanidine•HCl**
(4-($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{O}$) $\text{C}_6\text{H}_4\text{NHC}(=\text{NH})\text{NH}((4\text{-(Cl)}\text{C}_6\text{H}_4)\text{CH}_2\text{CH}_2)\text{•HCl}$)

Purple solid: mp: 163-164 °C; $R_f = 0.026$ (9:1 EtOAc:MeOH); ^1H NMR (300 MHz, CD_3OD): δ 7.23-7.38 (m, 4H, ArH), 6.92-7.10 (m, 4H, ArH),
10 3.95-4.15 (t, 2H, CH_2), 3.50-3.58 (t, 2H, CH_2), 2.88-2.93 (m, 2H, CH_2), 1.72-1.81 (m, 2H, CH_2), 1.71-1.81 (m, 2H, CH_2), 1.44-1.58 (m, 2H, CH_2), 0.95-1.20 (m, 2H, CH_3); MS(Cl): m/e 346 (M^+ for free base); Anal. Calcd. for $\text{C}_{19}\text{H}_{24}\text{N}_3\text{ClO•HCl•0.10H}_2\text{O}$: C, 59.40, H, 6.61, N, 10.93; Found: C, 59.15, H, 6.31, N, 10.84.

15 **Example 121:** **N-(4-Benzyloxyphenyl)-N,N'-diphenylguanidine•oxalate**
White solid: mp: 145 °C; ^1H NMR (300 MHz, CD_3OD): δ 7.30-7.50 (m, 15H, ArH), 7.20-7.25 (d, $J = 9\text{Hz}$, 2H, ArH), 7.02-7.07 (d, 2H, ArH), 5.10 (s, 2H, $\text{ArCH}_2\text{O-}$); HPLC: 97.6% pure; MS(EI): m/e 393 (M^+ for free
20 base); Anal. Calcd. for $\text{C}_{26}\text{H}_{23}\text{N}_3\text{O•0.8C}_2\text{H}_2\text{O}_4$: C, 71.21, H, 5.33, N, 9.03; Found: C, 70.87, H, 5.11, N, 9.05.

Example 122: **N-(4-Benzyloxyphenyl)-N'-benzyl-N'-phenylguanidine•oxalate**
25 White solid: mp: 180 °C; ^1H NMR (300 MHz, CD_3OD): δ 7.30-7.50 (m, 15H, ArH), 7.20-7.25 (d, $J = 9\text{Hz}$, 2H, ArH), 7.05-7.10 (d, $J = 9\text{Hz}$, 2H, ArH), 5.13 (s, 2H, $\text{Ph-CH}_2\text{O-}$), 5.08 (s, 2H, $\text{Ph-CH}_2\text{N-}$); HPLC: 99.4% pure; MS(EI): m/e 407 (M^+ for free base); Anal. Calcd. for
30 $\text{C}_{27}\text{H}_{25}\text{N}_3\text{O•C}_2\text{H}_2\text{O}_4\text{•0.5H}_2\text{O}$: C, 68.76, H, 5.57, N, 8.30; Found: C, 68.96, H, 5.36, N, 8.47.

Example 123: **N-(3-Benzyloxyphenyl)-N'-(4-benzylthiophenyl)**
guanidine•mesylate
((3-(C₆H₄CH₂O)C₆H₄)NHC(=NH)NH(4-(C₆H₄CH₂S)C₆H₄)
mesylate)

5

White solid: mp: 142-143°C; ¹H NMR (300 MHz, CD₃OD): δ 7.19-7.45 (m, 15H, ArH), 6.88-7.00 (m, 18H, ArH), 5.12 (s, 2H, S-CH₂), 4.20 (s, 2H, O-CH₂) 2.70 (s, 3H, Mesylate CH₃); HPLC: 96.6% pure; MS(EI): m/e 440 (M+ for free base); Anal. Calcd. for C₂₇H₂₅N₃OS•CH₃SO₃H: C, 62.78,
10 H, 5.46, N, 7.84; Found: C, 62.83, H, 5.49, N, 7.83.

Example 124: **N,N'-Bis(4-(pentylthio)phenyl)guanidine•HBr**

White solid: mp: 153°C; ¹H NMR (300 MHz, CD₃OD): δ 7.41-7.90
15 (m, 18H, ArH); MS(EI): m/e 427 (M+ for free base); Anal. Calcd. for C₂₅H₂₁N₃S₂•HBr•0.5H₂O: C, 58.02, H, 4.48, N, 8.12; Found: C, 57.94, H, 4.27, N, 8.09.

Example 125: **N,N'-Bis(3-(phenylthio)phenyl)guanidine•oxalate**

20

White solid: mp: 108°C; MS(EI): m/e 415 (M+ for free base); Anal. Calcd. for C₂₅H₃₅N₃O₄S₂•0.75H₂O: C, 57.83, H, 7.09, N, 8.09; Found: C, 57.94, H, 6.71, N, 8.34.

Example 126: **N-(5-Acenaphthyl)-N'-(2-phenylethyl)guanidine•HCl**

25

Yellow solid: mp: 98-101°C; R_f = 0.13 (10:1; CHCl₃/MeOH); ¹H NMR (300 MHz, CD₃OD): δ 7.25-7.57 (m, 10H, ArH), 3.56-3.59 (Brs, 2H, CH₂), 3.40-3.49 (m, 4H, CH₂), 2.90-2.94 (t, 2H, J = 14Hz, CH₂); MS(EI): m/e 315
30 (M+ for free base); Anal. Calcd. for C₂₁H₂₁N₃•HCl: C, 71.68, H, 6.30, N, 11.94; Found: C, 71.44, H, 6.26, N, 11.77.

Example 127: N-(5-Acenaphthyl)-N'-(3-butoxypropyl)guanidine•HCl

Brown semicrystalline solid: R_f = 0.19 (10:1; $\text{CHCl}_3/\text{MeOH}$): ^1H NMR (300 MHz, CD_3OD): δ 6.62-7.57 (m, 5H, ArH), 3.31-3.49 (m, 10H, CH_2), 1.61-1.65 (m, 2H, CH_2), 1.31-1.44 (m, 2H, CH_2), 1.25-1.31 (m, 2H, CH_2)
5 1.20-1.25 (m, 3H, CH_3); MS(EI): m/e 325 (M^+ for free base); Anal. Calcd. for $\text{C}_{20}\text{H}_{27}\text{N}_3\text{O}\cdot\text{HCl}$: C, 66.37, H, 7.80, N, 11.61; Found: C, 71.80, H, 7.27, N, 8.71.

10 **Example 128: N,N'-Bis(2,2-diphenylethyl)guanidine•HBr**

White solid: mp: 179-180°C, R_f = 0.20 (10:1; $\text{CHCl}_3/\text{MeOH}$): ^1H NMR (300MHz, CD_3OD): δ 7.21-7.36 (m, 20H, ArH), 3.76-3.79 (m, 4H, CH_2);
15 MS(EI): m/e 420 (M^+ for free base); Anal. Calcd. for $\text{C}_{29}\text{H}_{29}\text{N}_3\cdot\text{HBr}$: C, 69.6, H, 6.04, N, 8.4; Found: C, 69.43, H, 5.96, N, 8.27.

Example 129: N-(4-Butoxyphenyl)-N-(4-chlorobenzhydryl)guanidine•HCl

Purple solid: mp: 81-82°C; R_f = 0.53 (10:2; $\text{CHCl}_3/\text{MeOH}$); ^1H NMR (300MHz, CD_3OD): δ 6.15-7.40 (m, 13H, ArH), 3.94-3.99 (m, 2H, CH_2),
20 1.69-1.81 (m, 2H, CH_2), 1.42-1.55 (m, 2H, CH_2), 0.95-1.0 (m, 3H, CH_3); MS(Cl): m/e 408 (M^+ for free base); Anal. Calcd. for $\text{C}_{24}\text{H}_{26}\text{N}_3\text{OCl}\cdot\text{HCl}\cdot 1/3\text{H}_2\text{O}$: C, 63.85, H, 6.61, N, 4.73; Found: C, 63.80, H, 6.33, N, 9.59.

25 **Example 130: (5-Acenaphthyl)-N'-(phenethyl)-N'-benzylguanidine•maleate**

White solid: mp: 160°C; ^1H NMR (300 MHz, CD_3OD): δ 7.30-7.52 (m, 14H, ArH), 7.13-7.19 (m, 2H, ArH), 6.23 (s, 2H, maleate H), 4.83 (s, 2H, N-benzyl), 3.84-3.91 (t, 2H, N- CH_2), 3.40-3.47 (Brs, 4H, Acenaphthyl
30 CH_2), 3.06-3.12 (t, $J=7\text{Hz}$, 2H, Ph- CH_2); MS(EI): m/e 406 (M^+ for free base); Anal. Calcd. for $\text{C}_{28}\text{H}_{27}\text{N}_3\cdot\text{C}_4\text{O}_4$: C, 73.68, H, 5.99, N, 8.06; Found: C, 73.80, H, 6.09, N, 8.10.

Example 131: N-4-Benzyloxyphenyl)-N'-(3-benzyloxyphenyl)-N'-(4-chlorobenzyl)guanidine•mesylate

White solid: mp: 145-147°C; R_f = 0.30 (10:1; CHCl₃/MeOH); ¹H NMR (300MHz, CD₃OD): δ 6.91-7.45 (m, 22H, ArH), 5.12 (s, 2H, CH₂), 5.06 (s, 2H, CH₂), 5.01 (s, 2H, CH₂), 2.70 (s, 3H, CH₃); MS(EI): m/e 548 (M⁺ for free base); Anal. Calcd. for C₃₄H₃₀N₃O₂Cl•CH₃SO₃H: C, 62.67, H, 5.56, N, 6.27; Found: C, 62.82, H, 5.42, N, 6.73.

Example 132: N,N'-Bis(4-benzyloxyphenyl)-N'-methylguanidine•mesylate

White solid: mp: 168-171°C; R_f = 0.14 (10:1; CHCl₃/MeOH); ¹H NMR (300MHz, CD₃OD): δ 7.32-7.45 (m, 12H, ArH), 7.06-7.22 (m, 6H, ArH), 5.11-5.14 (d, 4H, J = 8Hz, CH₂), 3.42 (s, 3H, CH₃), 2.69 (s, 3H, CH₃); MS(EI): m/e 437 (M⁺ for free base); Anal. Calcd. for C₂₈H₂₇N₃O₂•CH₃SO₃H•1/4H₂O: C, 64.72, H, 5.90, N, 7.81; Found: C, 64.56, H, 5.87, N, 7.80.

Example 133: N-(3-Benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-phenylguanidine•HCl

White, grey solid: mp: 72-74°C; R_f = 0.20; (10:1; CHCl₃/MeOH); ¹H NMR (300MHz, CD₃OD): δ 7.21-7.43 (m, 18H, ArH), 7.00-7.03 (d, 2H, J = 9Hz, ArH), 6.70-6.79 (m, 3H, ArH), 5.08 (s, 4H, CH₂); MS(Cl): m/e 500 (M⁺ for free base); Anal. Calcd. for C₃₃H₂₉N₃O₂•HCl: C, 73.94, H, 5.64, N, 7.84; Found: C, 74.09, H, 5.41, N, 7.96.

Example 134: N-(4-sec-Butylphenyl)-N'-(4-isopropoxyphenyl)-N'-phenylguanidine•HCl

White solid: mp: 190-192°C; R_f = 0.59 (10:2; CHCl₃/MeOH); ¹H NMR (300MHz, CD₃OD): δ 6.97-7.51 (m, 13H, ArH), 4.59-4.89 (m, 1H, CH), 2.58-2.68 (m, 1H, CH), 1.53-1.66 (m, 2H, CH₂) 1.28-1.32 (m, 6H, CH₃), 1.20-1.25 (m, 3H, CH₃), 0.79-0.84 (m, 3H, CH₃); MS(Cl): m/e 402 (M⁺ for

free base); Anal. Calcd. for $C_{26}H_{31}N_3O \cdot HCl \cdot 1/2H_2O$: C, 68.48, H, 7.51, N, 9.21; Found: C, 69.81, H, 7.30, N, 9.89.

5 **Example 135: N-(4-Benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-phenylguanidine•mesylate**

White solid: mp: 168-170°C; R_f = 0.59 (9:3; EtOAc/MeOH); 1H NMR (300MHz, CD_3OD): δ 7.21-7.46 (m, 19H, ArH), 7.04-7.11 (m, 4H, ArH), 5.10-5.11 (m, 4H, CH_2), 2.69 (s, 3H, CH_3); MS(EI): m/e 499 (M^+ for free base); Anal. Calcd. for $C_{33}H_{29}N_3O_2 \cdot 1.5CH_3SO_3H$: C, 63.97, H, 5.24, N, 6.51; Found: C, 63.97, H, 5.09, N, 6.57.

10 **Example 136: N,N'-Bis(3-octyloxyphenyl)guanidine•HCl**

Light purple solid: mp: 107-108°C; R_f = 0.196 (10:1; $CHCl_3$ /MeOH); 1H NMR (300MHz, CD_3OD): δ 7.32-7.38 (t, 2H, J = 10Hz, ArH), 6.88-6.91 (m, 6H, ArH), 4.01-4.89 (m, 4H, CH_2), 1.75-1.80 (m, 4H, CH_2), 1.40-1.47 (m, 4H, CH_2), 1.31-1.40 (m, 16H, CH_2), 0.88-0.93 (t, 6H, J = 8Hz, CH_3); MS(EI): m/e 467 (M^+ for free base); Anal. Calcd. for $C_{29}H_{45}N_3O_2 \cdot HBr \cdot 1/2H_2O$: C, 62.56, H, 8.52, N, 7.55; Found: C, 62.80, H, 8.18, N, 7.63.

20 **Example 137: N,N'-Bis(4-butoxyphenyl)guanidine•HBr**

Cream solid: mp: 88°C; R_f = 0.11 (10:1; $CHCl_3$ /MeOH); 1H NMR (300MHz, CD_3OD): δ 7.23-7.26 (d, 4H, ArH), 6.99-7.02 (d, 4H, J = 9Hz, ArH), 3.98-4.02 (t, 4H, J = 13Hz, CH_2), 1.74-1.79 (m, 4H, CH_2), 1.47-1.54 (m, 4H, CH_2), 0.96-1.01 (t, 6H, J = 13Hz, CH_3); MS(EI): m/e 355 (M^+ for free base); Anal. Calcd. for $C_{21}H_{29}N_3O_2 \cdot HBr \cdot 1.0H_2O$: C, 57.80, H, 6.93, N, 9.63; Found: C, 55.17, H, 6.77, N, 10.36.

30 **Example 138: N,N'-Bis(4-phenoxyphenyl)guanidine•HBr**

White solid: mp: 127-128°C; R_f = 0.18 (10:1; $CHCl_3$ /MeOH); 1H NMR (300MHz, CD_3OD): δ 7.31-7.41 (m, 8H, ArH), 7.02-7.18 (m, 10H, ArH);

MS(EI): m/e 395 (M^+ for free base); Anal. Calcd. for $C_{25}H_{21}N_3O_2 \cdot HBr$: C, 63.03, H, 4.65, N, 8.82; Found: C, 62.77, H, 4.66, N, 8.84.

5 **Example 139: N-(3-Benzyloxyphenyl)-N'-(4-phenoxyphenyl)guanidine•mesylate**

White solid: mp: 89-90°C; R_f =0.18 (10:1; $CHCl_3$:MeOH); 1H NMR (300 MHz, CD_3OD): δ 7.30-7.45 (m, 10H, ArH), 7.15-7.30 (t, 1H, $J=10Hz$, ArH), 6.91-7.07 (m, 7H, ArH), 5.12 (s, 2H, CH_2), 2.69 (s, 3H, CH_3); MS(EI): m/e 409 (M^+ for free base); Anal. Calcd. for $C_{26}H_{32}N_3O_2 \cdot CH_3SO_3H$: C, 64.14, H, 5.38, N, 8.31; Found: C, 53.85, H, 5.38, N, 8.30.

15 **Example 140: N-(3-Benzyloxyphenyl)-N'-(4-phenylazophenyl)guanidine•mesylate**

Yellow orange solid: mp: 206-208°C; R_f =0.12 (10:1; $CHCl_3$:MeOH); 1H NMR (300 MHz, CD_3OD): δ 8.00-8.03 (m, 2H, ArH), 7.91-7.94 (m, 2H, ArH), 7.33-7.56 (m, 11H, ArH), 6.94-7.01 (m, 3H, ArH), 5.13 (s, 2H, CH_2), 2.69 (s, 3H, CH_3); MS(EI): m/e 421 (M^+ for free base); Anal. Calcd. for $C_{28}H_{23}N_5O \cdot CH_3SO_3H \cdot 0.25H_2O$: C, 62.11, H, 5.31, N, 13.42; Found: C, 61.86, H, 5.34, N, 13.25.

25 **Example 141: N,N'-Bis(3-benzyloxyphenyl)-N'-methylguanidine•HCl**

White solid: mp: 41-42°C; R_f =0.44 (10:2; $CHCl_3$:MeOH); 1H NMR (300 MHz, CD_3OD): δ 7.29-7.45 (m, 12H, ArH), 6.80-7.08 (m, 6H, ArH), 5.10-5.12 (d, 4H, $J=8Hz$, CH_2), 3.42 (s, 3H, CH_3); MS(EI): m/e 437 (M^+ for free base); Anal. Calcd. for $C_{28}H_{27}N_3O_2 \cdot HCl \cdot CH_3OH$: C, 68.83, H, 6.37, N, 8.30; Found: C, 69.13, H, 5.95, N, 8.09.

30 **Example 142: N-(4-Benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-methylguanidine•mesylate**

Purple-white solid: mp: 138-140°C; R_f =0.09 (10:1; $CHCl_3$:MeOH); 1H NMR (300 MHz, CD_3OD): δ 7.31-7.45 (m, 13H, ArH), 7.11-7.14 (m, 2H,

ArH), 6.86-7.00 (m, 3H, ArH), 5.10-5.14 (d, 4H, $J = 10\text{Hz}$, CH_2), 3.43 (s, 3H, CH_3), 2.68 (s, 3H, CH_3); MS(EI): m/e 437 (M^+ for free base); Anal. Calcd. for $\text{C}_{28}\text{H}_{27}\text{N}_3\text{O}_2 \cdot \text{CH}_3\text{SO}_3\text{H} \cdot 0.25\text{H}_2\text{O}$: C, 64.72, H, 5.90, N, 7.81; Found: C, 64.73, H, 5.96, N, 7.74.

5

Example 143: **N-(4-Butoxyphenyl)-N'-(4-isopropoxyphenyl)-N'-phenylguanidine•HCl**
 $(4-(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{O})\text{C}_6\text{H}_4)\text{NHC}(=\text{NH})\text{N}[(4-((\text{CH}_3)_2\text{CHO})\text{C}_6\text{H}_4)][\text{C}_6\text{H}_5] \cdot \text{HCl}$

10 Purple solid: mp: $161-164^\circ\text{C}$; ^1H NMR (300 MHz, CD_3OD): δ 6.93-7.47 (m, 13H, ArH), 4.58-4.62 (m, 1H, CH), 3.94-3.98 (t, 2H, $J = 13\text{Hz}$, CH_2), 1.68-1.79 (m, 2H, CH_2), 1.42-1.57 (m, 2H, CH_2), 0.93-1.00 (t, $J = 15\text{Hz}$, 3H, CH_3); MS(EI): m/e 418 (M^+ for free base); Anal. Calcd. for $\text{C}_{26}\text{H}_{31}\text{N}_3\text{O}_2 \cdot \text{HCl}$: C, 68.78, H, 7.10, N, 9.26; Found: C, 68.79, H, 7.22, N, 9.36.

15

Example 144: **N-N'-Bis(4-(4-hydroxybutyl)phenyl)guanidine•HBr**

Light yellow solid: mp: $143-44^\circ\text{C}$; $R_f = 0.16$ (10:2; $\text{CHCl}_3:\text{MeOH}$); ^1H NMR (300 MHz, CD_3OD): δ 7.23-7.33 (m, 8H, ArH), 3.55-3.59 (m, 4H, CH_2), 2.65-2.69 (m, 4H, CH_2), 1.62-1.70 (m, 4H, CH_2), 1.56-1.62 (m, 4H, CH_2); MS(Cl): m/e 356 (M^+ for free base); Anal. Calcd. for $\text{C}_{21}\text{H}_{29}\text{N}_3\text{O}_2 \cdot \text{HBr} \cdot 0.5\text{H}_2\text{O}$: C, 56.62, H, 7.01, N, 9.43; Found: C, 56.39, H, 6.67, N, 9.34.

20

25 **Example 145:** **N-(4-Butoxyphenyl)-N'-(3-methoxyphenyl)-N'-phenylguanidine•HCl**

Tan solid: mp: $79-81^\circ\text{C}$; $R_f = 0.043$ (10:1; $\text{CHCl}_3:\text{MeOH}$); ^1H NMR (300 MHz, CD_3OD): δ 7.23-7.51 (m, 6H, ArH), 7.23-7.26 (m, 2H, ArH), 6.96-7.02 (m, 5H, ArH), 3.95-4.20 (t, 2H, CH_2), 3.80 (s, 3H, CH_3), 1.70-1.80 (m, 2H, CH_2), 1.45-1.56 (m, 2H, CH_2), 0.94-1.10 (m, 3H, CH_3); MS(Cl): m/e 390 (M^+ for free base); Anal. Calcd. for

30

$C_{24}H_{27}N_3O_2 \cdot HCl \cdot 0.5H_2O$: C, 66.27, H, 6.72, N, 9.60; Found: C, 65.97, H, 6.69, N, 9.76.

Example 146: Inhibition of Glutamate Release

5 Compounds were tested for the inhibition of glutamate release. As shown by the data below, compounds of the invention are effective blockers of glutamate release. The assay involves adaptation of a rapid superfusion system (said system disclosed in S. Goldin, U.S. Patent No. 4,891,185 (1990); Turner, T.J. et al., *Anal. Biochem.*, 178:8-16 (1989)) to
10 measure depolarization-induced 3H -glutamate release from brain nerve terminals. The depolarizing stimulus opens presynaptic voltage-activated ion channels as the key step required to initiate Ca-dependent exocytosis of glutamatergic synaptic vesicles. The method involves preloading rat brain synaptosomes with 3H -glutamate via the Na-dependent glutamate uptake
15 system. The preloaded nerve terminals are retained in a superfusion chamber accessed by high-speed, solenoid-driven valves. Microcomputer-operated circuitry controls the timing of valve operation; the valves control the delivery under nitrogen pressure of pulses of depolarizing buffer, Ca, and/or drugs to the synaptosomes. The 3H -glutamate-containing effluent is
20 continuously collected in a high speed fraction collector on a subsecond timescale as short as 30 msec (300 msec fractions were employed herein). The high solution flow rate and minimal dead volume of the superfusion chamber, afford rapid solution changes and precise control of the chemical microenvironment of the nerve terminal preparation.

25

More specifically, the assay method employed was as described in Goldin et al., PCT/US92/01050, with the following modifications. Introduction of a buffer containing high $[K^+]$ was the means employed to produce the depolarization. This mode of depolarization is the preferred
30 method of opening presynaptic voltage-activated Ca channels to trigger glutamate release. An additional method of depolarization was also

employed, namely introduction of veratridine, and a parallel set of such veratridine-based experiments was performed. Veratridine is known to stimulate neurotransmitter release by opening voltage-activated Na channels, which results in depolarization of the nerve terminal plasma membrane and in turn, secondarily, opens presynaptic Ca channels to directly trigger ^3H -glutamate release via Ca-dependent exocytosis. The use of veratridine-induced glutamate release was employed to detect compounds of the invention which may block neurotransmitter release by blocking voltage-activated presynaptic Na channels. It has been previously reported that tetrodotoxin, a highly specific blocker of Type I and Type II neuronal Na channels, blocks veratridine-induced ^3H -glutamate release with no effect on high $[\text{K}^+]$ -induced ^3H -glutamate release (Katragadda et al. *Abs. Soc. for Neurosci.* 19:1750 (1993)). In experiments measuring veratridine-induced glutamate release, 50 μM veratridine in "basal" buffer was substituted for the "high-K buffer" employed as described in PCT/US92/01050; the protocol was otherwise identical to that described therein. In superfusion solutions containing compounds to be tested, compounds were made as stock solutions in methanol and diluted so that the final concentration of methanol never exceeded 0.3% (v/v). All solutions including compound free controls contained the same solvent [methanol]. Results are shown in the below tables (1a-1f) identified together as Table I. In that Table and tables of other examples which follow, the designation "FB" refers to the free base form of the specified compound.

TABLE 1

Inhibition of ^3H -glutamate release in brain nerve terminal preparations

Table 1a: Compounds of Formula IA

5

10

15

Example No.	Name	% Inhibition of ³ H-Glu Rel		Salt
		@ 3μM	@ 1μM	
1	N-(4-sec-butylphenyl)-N-(4-tert-butylbenzyl)guanidine	61	67*	HCl
2	N-(5-acenaphthyl)-N-(4-tert-butylbenzyl)guanidine	61	52	FB
19	N-(5-acenaphthyl)-N-(4-isopropylbenzyl)guanidine	81*		HCl
23	N-(4-sec-butylphenyl)-N-(transcinnamyl)guanidine	33*		HCl
24	N-(4-n-butoxyphenyl)-N-(4-tert-butylbenzyl)guanidine		62*	HCl
27	N-(3-trifluoromethoxyphenyl)-N-(4-tert-butylbenzyl)guanidine		24	HCl
47	N-(1-naphthyl)-N-(4-tert-butylbenzyl)guanidine		40	HCl
48	N-(3-iodophenyl)-N-(4-tert-butylbenzyl)guanidine		23*	HCl
49	N-(4-chloronaphthyl)-N-(4-tert-butylbenzyl)guanidine		32*	HCl
52	N-(1-naphthylmethyl)-N'-(4-tert-butylbenzyl)guanidine	55*		HCl
53	N-(5-acenaphthyl)-N-(3-phenoxybenzyl)guanidine		37	HCl
58	N-(5-acenaphthyl)-N-(4-iodobenzyl)guanidine		43	HCl• 3/2H ₂ O
59	N-(5-acenaphthyl)-N-(4-trifluoromethoxybenzyl)guanidine		44	HCl• 2H ₂ O

20

Table 1b: Compounds of Formula II

25

Example No.	Name	% Inhibition of ³ H-Glu Rel		Salt
		@ 3μM	@ 1μM	
9	N,N'-bis(fluoranthyl)guanidine	17*		HBr

Table 1c: Compounds of Formula IIIA

Example No.	Name	% Inhibition of ³ H-Glu Rel			Salt
		@ 3 μ M	@ 1 μ M	@ 0.3 μ M	
3	N-(5-acenaphthyl)-N'-benzhydrylguanidine	53			HCl
10	N-(5-acenaphthyl)-N'-(1-naphthylmethylene)guanidine	71*	24*		mesylate
40	N-(5-acenaphthyl)-N'-(1-methyl-2-(4-chlorophenyl)ethyl)guanidine		64	46	HCl
41	N-(5-acenaphthyl)-N'-(1,2-diphenylethyl)guanidine		52		HCl
42	N-(5-acenaphthyl)-N'-(3-phenylpropyl)guanidine	60*			HCl
43	N-(5-acenaphthyl)-N'-(2-methyl-2-phenylethyl)guanidine	60	43*		HCl
44	N-(5-acenaphthyl)-N'-(2-methyl-2-phenylethyl)guanidine	71*	43*		HCl
60	N-(5-acenaphthyl)-N'-((4-tert-butylphenyl)-(4-sec-butylphenyl)-methyl)guanidine	32			HCl

Table 1d: Compounds of Formula IV

Example No.	Name	% Inhibition of ³ H-Glu Rel			Salt
		@ 3 μ M	@ 1 μ M	@ 0.3 μ M	
3	N,N'-bis(4-sec-butylphenyl)guanidine	85	52		FB
5	N,N'-bis(4-sec-butylphenyl)-N-methyl guanidine		68*		HCl
6	N,N'-bis(4-sec-butylphenyl)-N,N'-bismethyl guanidine	92	71		HCl
8	N,N'-bis(4-sec-butylphenyl)-2-iminopyrimidazolidine	50			HBr
13	N,N'-bis(4-tert-butylphenyl)guanidine	70	62*		FB

Example No.	Name	% Inhibition of ³ H-Glu Rel			Salt
		@ 3 μ M	@ 1 μ M	@ 0.3 μ M	
14	N-(4- <i>tert</i> -butylphenyl)-N'-(2,3,4-trichloro phenyl)guanidine	63			HCl
15	N-(4-methoxynaphthyl)-N'-(2,3,4-trichlorophenyl)guanidine	88			HCl
30	N,N'-bis(3-biphenyl)guanidine		25		
31	N,N'-di-(3- <i>tert</i> -butylphenyl)guanidine		24		HBr
34	N,N'-bis-(3- <i>sec</i> -butylphenyl)guanidine		63		HBr
35	N,N'-bis(4- <i>tert</i> -butylphenyl)-N-methylguanidine		61		HCl
36	N,N'-bis(4- <i>tert</i> -butylphenyl)-N,N'-methylguanidine		67		FB
37	N,N'-bis(4-n-butylphenyl)guanidine		100	57	mesylate
42	N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl) guanidine		90	54	HCl
45	N,N'-bis(<i>sec</i> -butylphenyl)-N'-(2-phenoxyethyl)guanidine		57		HCl
46	N,N'-bis(<i>sec</i> -butylphenyl)-N'-(<i>n</i> -pentyl)guanidine		100	65	HCl

Table 1e: Compounds of Formula V

Example No.	Name	% Inhibition of ³ H-Glu Rel		Salt
		@ 3 μ M	@ 1 μ M	
7	N-(5-acenaphthyl)-N'-(1,2,3,4-tetrahydroquinoliny)guanidine	16	20	Mesylate
29	N-(5-acenaphthyl)-N-(indolynyl)guanidine	22		FB

* Results of veratridine-induced glutamate release assay as specified above.

Table 1f: Additional Compounds of the Invention
(Including Compounds of Formulas I-V)

Ex. No.	Name	% Inhibition of ³ H-Glu Rel				Salt
		@ 3 μ M	@ 1 μ M	@ 0.3 μ M	@ 0.1 μ M	
62	N-(3- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)guanidine		22			HCl
63	N-(3- <i>tert</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)guanidine		40			HCl
64	N-(3-pentoxyphenyl)-N-(4- <i>tert</i> -butylbenzyl)guanidine		41			HCl
65	N-(5-acenaphthyl)-N-(4-benzyloxybenzyl)guanidine		42			HCl
66	N-(4- <i>sec</i> -butylphenyl)-N-(4-benzyloxybenzyl)guanidine		54			HCl
67	N-(4-benzyloxyphenyl)-N-(4-benzyloxybenzyl)guanidine		36			HCl
68	N-(5-acenaphthyl)-N-(3-benzyloxybenzyl)guanidine		76			HCl
72	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>t</i> -butylbenzyl)-N'-pyrrolidinylguanidine			48		HCl
73	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>t</i> -butylbenzyl)-N'-(4-thiomorpholinyl)guanidine		59			HCl
74	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-piperidinylguanidine		100	21		HCl
75	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(4-morpholinyl)guanidine			42		HCl
76	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(4-propylpiperidinyl)guanidine			45		HCl
77	N-(4-butoxyphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(4-piperidinyl)guanidine			44		HCl
78	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(4-benzylpiperidinyl)guanidine			22		mesylate
79	N-(4-benzyloxyphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(4-morpholinyl)guanidine			77		HCl
80	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(1,2,3,4-tetrahydroisoquinolinyl)guanidine			68		HCl
82	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(3,5-dimethyl-4-morpholinyl)guanidine			60		HCl
83	N-(4- <i>tert</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(4- <i>sec</i> -butylphenyl)-N'-(methyl)guanidine		63	31		HCl
84	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(4- <i>sec</i> -butylphenyl)-N'-(methyl)guanidine		37			HCl
85	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(phenyl)guanidine		100	55		HCl
86	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(4-chlorophenyl)guanidine			45	16	HCl

Ex. No.	Name	% Inhibition of ³ H-Glu Rel				Salt
		@ 3μM	@ 1μM	@ 0.3μM	@ 0.1μM	
5	87 N-(4-butoxyphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(phenyl)guanidine			36		HCl
	88 N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(phenyl)-N'-methylguanidine			77		HCl
	89 N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(3,4-dichlorophenyl)guanidine			21		HCl
	90 N-(4-hexylphenyl)-N-(4-hexylbenzyl)-N'-phenylguanidine			42		HCl
	91 N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)-N'-(4-benzyloxyphenyl)guanidine			49		mesylate
10	92 N,N'-bis-(4- <i>tert</i> -butylphenyl)-N,N'-dimethylguanidine		45			HBr
	93 N-(4-benzyloxyphenyl)-N'-(4- <i>tert</i> -butylphenyl)guanidine		91	48		HCl
	94 N,N'-bis-(3-(1'-methyl-2'-phenyl)ethyl)guanidine		66			HCl
	95 N-methyl-N-4-benzyloxyphenyl-N'-(4- <i>tert</i> -butylphenyl)guanidine		100	44		mesylate
	96 N,N'-bis-(4-hexylphenyl)guanidine			54	54	mesylate
15	97 N-(3-(1-(4'-ethoxy)benzyl)phenethyl)-N'-(4- <i>tert</i> -butylphenyl)guanidine			31		mesylate
	98 N-(4-benzyloxyphenyl)-N'-methyl-N-(4- <i>tert</i> -butylphenyl)guanidine			76		mesylate
	99 N-(3-(4- <i>tert</i> -butylbenzyloxy)phenyl)-N'-(4- <i>tert</i> -butylphenyl)guanidine			20		mesylate
	100 N-(3-(1'-benzylbutyl)phenyl)-N'-(4- <i>tert</i> -butylphenyl)guanidine			20		mesylate
	101 N,N'-bis-(4-butylphenyl)-N-methylguanidine			35		HCl
20	102 N,N'-bis-(4- <i>tert</i> -butylphenyl)-N,N'-dimethylguanidine			73	18	HCl
	103 N-(3-naphthylmethylenoxyphenyl)-N'-(4- <i>tert</i> -butylphenyl)guanidine			50		mesylate
	104 N-(4-benzyloxyphenyl)-N'-(4-butylphenyl)guanidine			76		HCl
	105 N,N'-bis-(4-butylphenyl)-N-butylguanidine			66	24	HCl
	106 N-3-(benzyloxymethyl)phenyl-N'-(4- <i>tert</i> -butylphenyl)guanidine			48		mesylate
	107 N-(3,4-bis-butylloxyphenyl)-N'-(4- <i>tert</i> -butylphenyl)guanidine			34		oxalate
	108 N-(3-benzyloxy)phenyl-N'-(4- <i>tert</i> -butylphenyl)guanidine			49		mesylate
	109 N,N'-bis-(3-butoxy-4-methoxy)phenylguanidine			86		HBr
	110 N-(4-benzyloxyphenyl), methyl-N'-(4-butylphenyl)guanidine			77		mesylate

Ex. No.	Name	% Inhibition of ³ H-Glu Rel				Salt
		@ 3μM	@ 1μM	@ 0.3μM	@ 0.1μM	
111	N,N'-bis-(6-tetralinyl)guanidine			61		HBr
117	N-(5-acenaphthyl)-N'-(1-benz[cd]indolyl)guanidine		39			HCl
138	N-(bis(4-phenoxyphenyl)guanidine			57		HBr
---	N,N'-bis(benzyloxyphenyl)guanidine			46		mesylate
---	N-(4-benzyloxyphenyl)-N'-(4-benzylthiophenyl)guanidine			81		mesylate
140	N-(3-benzyloxyphenyl)-N'-(4-(azophenyl)phenyl)guanidine			44		mesylate
---	N,N'-bis(3-benzyloxyphenyl)-N'-methylguanidine			67		HCl
---	N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-methylguanidine			55		mesylate
---	N,N'-bis(4-benzyloxyphenyl)-N'-methylguanidine			22		mesylate

Example 147: Inhibition of ⁴⁵Ca uptake through Presynaptic Ca Channels

Compounds of the invention were tested to determine their ability to inhibit voltage-activated calcium channels in nerve terminals of mammalian brain. Said voltage-activated calcium channels directly control neurotransmitter release (see Nachsen, D.A. et al., *J. Gen Physiol.*, **79**:1065-1087 (1982)). The uptake of ⁴⁵Ca into brain synaptosomes was performed by an adaptation of the method of Nachsen and Blaustein (*J. Physiol.*, **361**:251-258 (1985)), as previously described [Goldin et al., PCT/US92/01050]. The principle of the method involves opening ion permeation through synaptosomal calcium channels by high K⁺- induced depolarization of the synaptosomal preparation. The rapid component of ⁴⁵Ca uptake measured by this procedure is mediated by presynaptic calcium channels.

Briefly, synaptosomes are prepared by the method of Hajos (*Brain Res.*, **93**:485-489 (1975)). Freshly prepared synaptosomes (8 μl) were suspended in low potassium "LK" buffer (containing 3 mM KCl). Test

compounds in 8 μ l LK were added to synaptosomes to final concentrations ranging from 0.3 μ M to 100 μ M, and the mixture was preincubated for 5 minutes at room temperature. 45 Ca uptake was then initiated by adding isotope in either LK or in buffer ("HK") containing high [potassium] (150 mM KCl). After 5 seconds, the 45 Ca uptake was stopped by adding 0.9 ml quench buffer (LK + 10 mM EGTA). This solution was then filtered under vacuum and the filters washed with 15 ml of quench buffer.

Washed filters were subjected to scintillation spectrophotometry to determine the extent of 45 Ca uptake. Net depolarization-induced 45 Ca uptake was determined for each concentration of each compound tested, as the difference between 45 Ca uptake in HK and LK buffers. Results were plotted as % inhibition of depolarization-induced 45 Ca uptake vs. [compound] for each compound tested. Representative IC_{50} for inhibition of depolarization-induced 45 Ca uptake are presented below in the tables (2a-2d) below which together are identified as Table 2.

TABLE 2

Inhibition of 45 Ca uptake through presynaptic Ca channels

Table 2a: Compounds of Formula IA

Example No.	Name	IC_{50} , block of 45 Ca uptake, μ M	Salt
1	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)guanidine	6.1	HCl
2	N-(5-acenaphthyl)-N-(4- <i>tert</i> -butylbenzyl)guanidine	8.1	FB

Table 2b: Compounds of Formula IIIA

Example No.	Name	IC ₅₀ , block of ⁴⁵ Ca uptake, μ M	Salt
5 9	N-(5-acenaphthyl)-N'-(1-naphthyl-methylene)guanidine	7.8	Mesylate

Table 2c: Compounds of Formula IV

Example No.	Name	IC ₅₀ , block of ⁴⁵ Ca uptake, μ M	Salt
10 4	N,N'-bis(4-sec-butylphenyl)guanidine	11.2	HCl
6	N,N'-bis(4-sec-butylphenyl)-N,N'-bismethyl guanidine	5.9	HCl
13	N,N'-bis(4-tert-butylphenyl)guanidine	1.5	FB
15 15	N-(4-methoxynaphthyl)-N'-(2,3,4-trichlorophenyl)guanidine	1.7	HCl

Example 148: Inhibition of ⁴⁵Ca uptake through L-type (dihydropyridine-sensitive) Calcium Channels

Compounds of the invention representative of each of the major classes of agents claimed herein were tested to determine their ability to inhibit voltage-activated, dihydropyridine-sensitive L-type calcium channels in clonal GH4C1 pituitary cells. Said voltage-activated L-type calcium channels are found in cardiac muscle, vascular smooth muscle, and the cardiac Purkije cell conduction system. They are the sites of action of the major classes of Ca antagonists employed to treat hypertension, angina, cardiac arrhythmias, and related disorders. L-type Ca channels are also the sites of action of certain neuroprotective dihydropyridine Ca antagonists such as nimodipine.

The uptake of ⁴⁵Ca into GH4C1 cells was performed by an adaptation of the method of Tan, K. et al. (*J. Biol. Chem.*, 259:418-426 (1984)). The

principle of the method involves activating ion permeation through synaptosomal calcium channels by high K^+ - induced depolarization of the synaptosomal preparation. The uptake of ^{45}Ca measured by this procedure is mediated by presynaptic L-type calcium channels, and is sensitive to

5 dihydropyridine, phenylalkylamine, and benzothiazipine Ca antagonists at therapeutically relevant concentrations [Tashjian et al., *ibid.*]. The adaptation of the aforementioned method involves growing GH4C1 cells in 96-well culture plates, and is designed to provide a rapid and quantitative determination of the potency of various compounds in inhibiting ^{45}Ca uptake

10 through L-type Ca channels.

Details of method:

GH4 cells, stored in liquid nitrogen, are suspended in 15 ml growth medium (Ham's F-10 medium plus 15% heat-inactivated horse serum and

15 2.5% heat-inactivated fetal bovine serum). The cells are centrifuged, resuspended, and then added to T-75 flasks containing 12-15 mls Growth Medium, and incubated at 37°C for approximately 1 week. The cells are then removed from the T75 flask after dissociation from the walls of the flask by treatment for 5 minutes at 37°C with 1 mg/ml Viocase. The

20 Viocase is decanted, and the cells are resuspended in ~ 200 ml of Growth Medium. The cells are then aliquoted (200 μ l/well) into each well of several 96 well plates. The cells are then grown under the aforementioned conditions for 3-4 weeks, with replacement of Growth Medium occurring twice per week. Cells are fed growth medium 24 hours before they are

25 employed for ^{45}Ca uptake determinations.

At the time of the assay, media is aspirated from each 96-well plate using a manifold designed to allow 50 μ L of liquid to remain in each well. Each plate is washed and aspirated twice with a low K^+ buffer solution

30 "LKHBBBS" (in mM 5 KCl, 145 NaCl, 10 Hepes, 1 $MgCl_2$, 0.5 $MgCl_2$, 10 glucose, pH 7.4), 200 μ l/well. Each plate is incubated for 10 minutes at

37°C, and aspirated as above. To each well of each plate, 50 μ l of HBBS containing the drug to be tested in twice the final concentration is added. The plates are incubated for 10 minutes at room temperature. To each well of each plate, 50 μ l of either of two solutions are added:

- 5 (a) LKHBBS containing 1 μ Ci of carrier-free ^{45}Ca , or
- (b) HKHBBS (a high K^+ buffer containing 150 mM KCl and no NaCl, but otherwise identical to LKHBBS).

Each plate is then incubated for 5 minutes at room temperature,
10 aspirated as above, and quenched with 200 μ l/well of Quench Buffer (Ca-free LKHBBS containing 10 mM Tris-EGTA). Each plate is aspirated and rinsed with Quench Buffer a second time, then carefully aspirated to dryness. To each well of each plate 100 μ l of High Safe II scintillation fluid is added. The plates are sealed, shaken, and subjected to scintillation
15 spectrophotometry on a Microbeta 96-well Scintillation Counter (Wallac, Gaithersburg, MD, USA).

Net depolarization-induced ^{45}Ca uptake was determined for each concentration of each compound tested, as the difference between ^{45}Ca
20 uptake in HKBBS and LK buffers. Results were plotted as % inhibition of depolarization-induced ^{45}Ca uptake vs. [compound] for each compound tested. Representative IC_{50} for inhibition of depolarization-induced ^{45}Ca uptake are presented below in the tables (3a-3c) below which together are identified as Table 3. The known compounds of verapamil and diltiazem
25 were also tested pursuant to the same protocol as specified above and the following activity against L-type Ca channels was observed: verapamil: IC_{50} (block of ^{45}Ca uptake, μM) 8.0 ± 4 ($n=3$); diltiazem: IC_{50} (block of ^{45}Ca uptake, μM) 19.7 ± 6 ($n=3$).

TABLE 3

Inhibition of ⁴⁵Ca uptake through L-type Ca channels

Table 3a: Compounds of Formula IA

Example No.	Name	IC ₅₀ , block of ⁴⁵ Ca uptake, μM	Salt
1	N-(4- <i>sec</i> -butylphenyl)-N-(4- <i>tert</i> -butylbenzyl)guanidine	1.7	FB
2	N-(5-acenaphthyl)-N-(4- <i>tert</i> -butylbenzyl)guanidine	2.1	FB
19	N-(5-acenaphthyl)-N-(4- <i>iso</i> -propylbenzyl)guanidine	2.1	HCl

Table 3b: Compounds of Formula IIIA

Example No.	Name	IC ₅₀ , block of ⁴⁵ Ca uptake, μM	Salt
3	N-(5-acenaphthyl)-N'-(benzhydryl)guanidine	2.7	HCl

Table 3c: Compounds of Formula IV

Example No.	Name	IC ₅₀ , block of ⁴⁵ Ca uptake, μM	Salt
4	N,N'-bis(4- <i>sec</i> -butylphenyl)guanidine	2.1	FB
6	N,N'-bis(4- <i>sec</i> -butylphenyl)-N,N'-bismethyl guanidine	4.3	HCl
13	N,N'-bis(4- <i>tert</i> -butylphenyl)guanidine	4.1	FB
15	N-(4-methoxynaphthyl)-N'-(2,3,5-trichlorophenyl)guanidine	1.7	HCl

Example 149: Inhibition of ^{14}C -guanidine uptake through Type II Neuronal Voltage-activated Sodium Channels

Antagonists of the neuron-specific type II subclass of voltage-gated Na channels are neuroprotective (Stys, P. K. et al., *J. Neurosci.*, **12**:430-439 (1992)). The ability of compounds of the invention to block voltage-activated Type II Na channels was determined in a functional assay employing a Chinese Hamster Ovary ("CHO") cell line expressing cloned Type II Na channels derived from rat brain (West, J.W. et al., *Neuron* **8**:59-70 (1992)). The assay is based on the observation that veratridine, an alkaloid neurotoxin, causes persistent activation of sodium channels, and tetrodotoxin, a heterocyclic agent derived from puffer fish, is a potent and highly specific blocker of several major subclasses of voltage-sensitive sodium channels, including the said Type II subclass. It further takes advantage of the finding that guanidinium cation will permeate through tetrodotoxin-sensitive Na channels when said channels are opened, either by membrane depolarization (Hille, B., *Ionic Channels of Excitable Membranes 2nd Edition*, Sinauer Associates, Sunderland, MA, pp. 349-353 (1992)) or by exposure to veratridine (Reith, M.E.A., *Eur. J. Pharmacol.*, **188**:33-41(1990)). Accordingly, the assay entails measuring veratridine-stimulated, tetrodotoxin-sensitive influx of [^{14}C]-guanidinium ion through cloned Type II Na channels expressed in CHO cells. The protocol of the assay is as follows.

Assay protocol:

The aforementioned CHO cell line is grown by standard cell culture techniques in RPMI 1640 medium (Media Tech), supplemented with 5% fetal calf serum (Hyclone), 200 $\mu\text{g/ml}$ G418 (Sigma) and 5.75 mg/ml proline (Sigma). Cells are routinely allowed to grow for 3-4 days *in vitro*.

Cultures are rinsed 3 times with 200 μl of "preincubation buffer" (5.4 mM KCl, 0.8mM MgSO_4 , 50 mM Hepes, 130 mM choline chloride, 0.1

mg/ml BSA, 1 mM guanidine HCl, 5.5 mM D-glucose, pH 7.4) and incubated with 200 μ l preincubation buffer at 37°C for 10 minutes. A 96-channel manifold connector is used to vacuum-aspirate the buffer from the wells between rinses.

5

Different concentrations of the tested compounds are prepared by dilution into "uptake buffer" (preincubation buffer plus ~ 2.5 mCi/ml [14 C]-guanidinium HCl, ~ 40 mCi/mmol) containing veratridine (100 μ M).

10 Aliquots (50 μ l) of these working stocks is added to the 96-well plates and incubated at room temperature for 1 hour. The veratridine-induced [14 C]-guanidinium uptake was linear with time and a good signal (4-8 fold basal uptake) was obtained following a 1 hour incubation. The following controls are also conducted in each 96-well plate: basal uptake (obtained in the absence of CNS compound and veratridine), uptake evoked by veratridine
15 alone, and veratridine evoked uptake in the presence of 10 μ M tetrodotoxin (the latter is a measure of non-specific uptake independent of Na channel activation).

20 The flux assay is terminated at the end of the incubation period by rinsing the 96-well plates 3 times with 200 μ l/well of ice cold "wash buffer" (163 mM choline chloride, 0.8 mM Mg SO₄, 1.8 mM Ca Cl₂, 5 mM Hepes, 1 mg/ml BSA). The remaining 50 μ l of wash buffer in the wells (following rinsing) is removed by vacuum aspiration with an 8-channel Drummond aspirator. 100 μ l of High Safe II scintillation fluid is added to each well.
25 The plates are sealed before shaking for 15 minutes. The plates are then allowed to sit for 45 minutes before counting in a 96-well scintillation counter.

30 Net veratridine-induced [14 C]-guanidinium uptake was determined for each concentration of each compound tested, as the difference between [14 C]-guanidinium uptake in the presence and absence of tetrodotoxin.

Results were plotted as % inhibition of veratridine-induced [¹⁴C]-guanidinium uptake vs. [compound] for each compound tested.

Representative IC₅₀'s for inhibition of veratridine-induced [¹⁴C]-guanidinium uptake are presented below in the following tables (4a-4d) which together are identified as Table 4.

TABLE 4

Inhibition of [¹⁴C]-guanidinium uptake through type II sodium channels

Table 4a: Compounds of Formula IA

Example No.	Name	IC ₅₀ , block of ¹⁴ C guanidinium uptake, μM	Salt
1	N-(4-sec-butylphenyl)-N-(4-tert-butylphenyl)guanidine	0.8	HCl
2	N-(5-acenaphthyl)-N-(4-tert-butylphenyl)guanidine	1	FB
16	N-(4-sec-butylphenyl)-N-(benzyl)guanidine	0.5	FB
18	N-(5-acenaphthyl)-N-(4-isopropylbenzyl)guanidine	2.1	HCl
20	N-(4-cyclohexyl)-N-(4-tert-butylbenzyl)guanidine	0.8	HCl
21	N-(fluorenyl)-N-(4-tert-butylbenzyl)guanidine	1.9	HCl
22	N-(4-sec-butylphenyl)-N-(transcinnamyl)guanidine	0.8	HCl
24	N-(3-biphenyl)-N-(4-tert-butylbenzyl)guanidine	1.8	HCl
26	N-(3-trifluoromethoxyphenyl)-N-(4-tert-butylbenzyl)guanidine	0.7	HCl
27	N-(4-methoxynaphthyl)-N-(4-tert-butylbenzyl)guanidine	1.2	HCl
48	N-(3-iodophenyl)-N-(4-tert-butylbenzyl)guanidine	0.4	HCl
49	N-(4-chloronaphthyl)-N-(4-tert-butylbenzyl)guanidine	2.2	HCl

Table 4b: Compounds of Formula II

Example No.	Name	IC ₅₀ , block of ¹⁴ C guanidinium uptake, μ M	Salt
11	N,N'-bis(fluorenyl)guanidine	0.9	HBr

Table 4c: Compounds of Formula IIIA

Example No.	Name	IC ₅₀ , block of ¹⁴ C guanidinium uptake, μ M	Salt
9	N-(5-acenaphthyl)-N'-(1-naphthyl-methylene)guanidine	0.7	HCl
42	N-(5-acenaphthyl)-N'-(2-methyl-3-phenylpropyl)guanidine	0.7	HCl

Table 4d: Compounds of Formula IV

Example No.	Name	IC ₅₀ , block of ¹⁴ C guanidinium uptake, μ M	Salt
4	N,N'-bis(4-sec-butylphenyl)guanidine	0.8	HCl
6	N,N'-bis(4-sec-butylphenyl)-N,N'-bismethyl guanidine	0.6	HCl
8	N,N'-bis(4-sec-butylphenyl)-2-iminopyrimidazolidine	0.2	HBr
12	N,N'-bis(4-tert-butylphenyl)guanidine	1.5	FB
13	N-(4-tert-butylphenyl)-N'-(2,3,4-trichlorophenyl)guanidine	0.9	HCl
14	N-(4-methoxynaphthyl)-N'-(2,3,4-trichlorophenyl)guanidine	1.8	HCl

Example 150: In vivo Anticonvulsant activity in the DBA/2 mouse model

The in vivo potency of compounds of the present invention is exemplified by data summarized in the Table V below and obtained pursuant to the following protocol.

Compounds were tested for their effectiveness in preventing seizures in DBA/2 mice which have a unique sensitivity to auditory stimulation. Exposure to loud high-frequency sounds can trigger seizure activity in these animals. This sensitivity develops from postnatal day 12 and peaks around day 21 and slowly diminishes as the animals mature. The unusual response to auditory stimulation in this strain of mouse is believed to be due to a combination of early myelination (causing an unusually low excitatory threshold) and delayed development of inhibitory mechanisms. Glutamate, the predominant excitatory neurotransmitter, has been implicated in this response. Blockade of glutamate receptors of the NMDA and AMPA subtypes, prevents audiogenic seizures in these mice. Compounds that block the release of glutamate should similarly act to prevent seizure activity, and may be therapeutic in other neurologic disorders such as stroke, which also involves glutamate-mediated damage.

Mice were injected intraperitoneally with the compound specified in Table V below or with vehicle control, 30 minutes prior to being placed in a bell jar and turning on the auditory stimulus (12 KHz sine wave at 110-120 db). Administered doses are specified in Table V as milligram of compound per kilogram bodyweight of mouse. The auditory stimulus was left on for 60 seconds and mice reactions were timed and recorded.


Percentage inhibition was determined with reference to vehicle controls. Results are shown in the Table 5 below.

Table 5

Example No.	Compound Name	Audiogenic Response		Salt
		Dose (mg/kg)	% Inhib.	
8	N,N'-bis-(4-sec-butyl-2-iminopyrimidazolidine	2	41	HBr
		4	75	
29	N-(5-acenaphthyl)-N'-(indoliny)guanidine	20	42	mesylate
		40	81	
1	N-(4-sec-butylphenyl)-N-(4-tert-butylbenzyl)guanidine	40	91	HCl
		10	54	
		20	80.5	
37	N,N'-bis(4-n-butylphenyl)guanidine	25	82	mesylate
		40	92	
72	N-(4-sec-butylphenyl)-N-(4-t-butylbenzyl)-N'-pyrrolidinyguanidine	40	81	HCl
		20	52	
---	N,N'-bis(4-n-butoxyphenyl)guanidine	20	60	
		40	82	
73	N-(4-sec-butylphenyl)-N-(4-t-butylbenzyl)-N'-(4-thiomorpholiny)guanidine	20	23	HCl
		40	89	
---	N-(4-benzyloxyphenyl)-N'-(4-phenoxyphenyl)guanidine	25	16	
		40	93	
104	N-(4-benzyloxyphenyl)-N'-(4-butylphenyl)guanidine	20	40	HCl
		40	90	
106	N-3-benzyloxymethyl)phenyl-N'-(4-tert-butylphenyl)guanidine	40	93	mesylate
108	N-(3-benzyloxy)phenyl-N'-(4-tert-butylphenyl)guanidine	40	80	mesylate
79	N-(4-benzyloxyphenyl)-N-(4-tert-butylbenzyl)-N'-(4-morpholiny)guanidine	20	41	HCl
		40	68	
110	N-(4-benzyloxyphenyl)-N-methyl-N'-(4-butylphenyl)guanidine	20	42.3	mesylate
		40	92.5	

What is claimed is:

1. A compound of the following formula I:




wherein:

R and R¹ are each independently substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least about 5 ring atoms, substituted or unsubstituted aralkyl having at least about 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having from 1 to 3 rings, 3 to 8 ring members in each ring and from 1 to 3 hetero atoms, with at least one of R and R¹ being carbocyclic aryl, aralkyl, or a heteroaromatic group or heteroalicyclic group;

R² and R³ each being independently selected from the group consisting of hydrogen, substituted and unsubstituted alkyl, substituted and unsubstituted alkoxy, substituted and unsubstituted thioalkyl, and substituted and unsubstituted aminoalkyl; and wherein said compound exhibits high inhibition of glutamate release; and pharmaceutically acceptable salts thereof.

2. A compound having the following Formula IA:




wherein R and R¹ are each independently substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least about 5 ring atoms, substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted aralkyl having at least about 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having from 1 to 3 rings, 3 to 8 ring members in each ring and from 1 to 3 hetero atoms; and pharmaceutically acceptable salts thereof.

3. A compound of claim 2 wherein at least one of R and R¹ is substituted or unsubstituted carbocyclic aryl or substituted or unsubstituted aralkyl.

4. A compound of claim 2 selected from the group of:
N-(4-*sec*-butylphenyl)-N-benzylguanidine;
N-(5-acenaphthyl)-N-benzylguanidine;
N-(3-acenaphthyl)-N-benzylguanidine;
N-(5-acenaphthyl)-N-(4-*isopropylbenzyl*)guanidine;
N-(3-acenaphthyl)-N-(4-*isopropylbenzyl*)guanidine;

N-(4-cyclohexylphenyl)-N-(4-*isopropyl*benzyl)guanidine;
N-(4-cyclohexylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(2-fluorenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(cinnamylmethylene)guanidine;
N-(4-*n*-butoxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-biphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(5-indanyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-trifluoromethoxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(5-acenaphthyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-acenaphthyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(methoxy-1-naphthyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(1-naphthyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-iodophenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(4-chloro-1-naphthyl)-N-(4-*tert*-benzyl)guanidine;
N-(4-*tert*-butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(4-iodophenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(1-naphthylmethyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(5-acenaphthyl)-N-(3-phenoxybenzyl)guanidine;
N-(3-trifluoromethylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-methylthiophenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(5-acenaphthyl)-N-(3-iodobenzyl)guanidine;
N-(5-acenaphthyl)-N-(cinnamyl)guanidine;
N-(5-acenaphthyl)-N-(4-iodobenzyl)guanidine;
N-(5-acenaphthyl)-N-(4-trifluoromethoxybenzyl)guanidine;
and pharmaceutically acceptable salts thereof.

5. A compound of the following Formula II:



wherein

R is selected from the group of fluorenyl, phenanthracenyl, anthracenyl and fluoranthenyl;

R¹ is substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfonyl having from 1 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least about 5 ring atoms, substituted or unsubstituted aralkyl having at least about 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having 1 to 3 rings, 3 to 8 ring members in each ring and 1 to 3 heteroatoms;

R² and R³ are each independently hydrogen or a group as defined for R¹ above; and pharmaceutically acceptable salts thereof.


6. A compound of claim 5 wherein R¹ is substituted or unsubstituted carbocyclic aryl.

7. A compound of claim 5 selected from the group of:
N,N'-bis(2-fluorenyl)guanidine;
N,N'-bis(2-fluorenyl)-N-methylguanidine;

N,N'-bis(2-fluorenyl)-N,N'-dimethylguanidine;
N,N'-bis(anthracenyl)guanidine;
N,N'-bis(anthracenyl)-N-methylguanidine;
N,N'-bis(anthracenyl)-N,N'-dimethylguanidine;
N,N'-bis(phenanthracenyl)guanidine;
N,N'-bis(phenanthracenyl)-N-methylguanidine;
N,N'-bis(phenanthracenyl)-N,N'-dimethylguanidine;
N,N'-bis(fluoranthenyl)guanidine;
N,N'-bis(fluoroanthenyl)-N-methylguanidine;
N,N'-bis(fluoroanthenyl)-N,N'-dimethylguanidine;
N-(anthracenyl)-N'-(1-adamantyl)guanidine;
N-(anthracenyl)-N'-(1-adamantyl)-N-methylguanidine;
N-(anthracenyl)-N'-(1-adamantyl)-N'-methylguanidine;
N-(anthracenyl)-N'-(1-adamantyl)-N,N'-dimethylguanidine;
N-(anthracenyl)-N'-(2-adamantyl)guanidine;
N-(anthracenyl)-N'-(2-adamantyl)-N-methylguanidine;
N-(anthracenyl)-N'-(2-adamantyl)-N'-methylguanidine;
N-(anthracenyl)-N'-(2-adamantyl)-N,N'-dimethylguanidine;
N-(phenanthracenyl)-N'-(1-adamantyl)guanidine;
N-(phenanthracenyl)-N'-(1-adamantyl)-N-methylguanidine;
N-(phenanthracenyl)-N'-(1-adamantyl)-N'-methylguanidine;
N-(phenanthracenyl)-N'-(1-adamantyl)-N,N'-dimethylguanidine;
N-(phenanthracenyl)-N'-(2-adamantyl)guanidine;
N-(phenanthracenyl)-N'-(2-adamantyl)-N-methylguanidine;
N-(phenanthracenyl)-N'-(2-adamantyl)-N'-methylguanidine;
N-(phenanthracenyl)-N'-(2-adamantyl)-N,N'-dimethylguanidine;
N-(fluorenyl)-N'-(1-adamantyl)guanidine;
N-(fluorenyl)-N'-(1-adamantyl)-N-methylguanidine;
N-(fluorenyl)-N'-(1-adamantyl)-N'-methylguanidine;
N-(fluorenyl)-N'-(1-adamantyl)-N,N'-dimethylguanidine;
N-(fluorenyl)-N'-(2-adamantyl)guanidine;

N-(fluorenyl)-N'-(2-adamantyl)-N-methylguanidine;
 N-(fluorenyl)-N'-(2-adamantyl)-N'-methylguanidine;
 N-(fluorenyl)-N'-(2-adamantyl)-N,N'-dimethylguanidine;
 and pharmaceutically acceptable salts of said compounds.

8. A compound of the following Formula III:



wherein

R and R¹ are each independently substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aryloxy having from 6 to about 20 carbon atoms, substituted or unsubstituted aralkoxy having from 6 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfonyl having from 1 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least 5 ring atoms, substituted or unsubstituted aralkyl having at least 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having 1 to 3 rings, 3 to 8 ring members in each ring and 1 to 3 heteroatoms;

R² and R³ are each independently hydrogen or a group as defined for R and R¹ above; or R² and R³ together form a ring having 5 or more ring members;

to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least 5 ring atoms, substituted or unsubstituted aralkyl having at least 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having 1 to 3 rings, 3 to 8 ring members in each ring and 1 to 3 heteroatoms;

R^2 and R^3 are each independently hydrogen or a group as defined for R and R^1 above; or R^1 and R^3 together form a ring having 5 or more ring members;

n is equal to 1, 2 or 3;

X is substituted or unsubstituted alkylene having from 1 to about 6 carbon atoms; and pharmaceutically acceptable salts thereof.

10. A compound of claim 8 selected from the group of:

N-5-acenaphthyl-N'-benzhydrylguanidine;

N-5-acenaphthyl-N'-benzhydryl-N-methylguanidine;

N-5-acenaphthyl-N'-benzhydryl-N'-methylguanidine;

N-5-acenaphthyl-N'-benzhydryl-N,N'-dimethylguanidine;

N-3-acenaphthyl-N'-benzhydrylguanidine;

N-3-acenaphthyl-N'-benzhydryl-N-methylguanidine;

N-3-acenaphthyl-N'-benzhydryl-N'-methylguanidine;

N-3-acenaphthyl-N'-benzhydryl-N,N'-dimethylguanidine;

N-(5-acenaphthyl)-N'-[(1-naphthyl)-methyl]guanidine;

N-(5-acenaphthyl)-N'-[(1-naphthyl)-methyl]-N-methylguanidine;

N-(5-acenaphthyl)-N'-[(1-naphthyl)-methyl]-N'-methylguanidine;

N-(5-acenaphthyl)-N'-[(1-naphthyl)-methyl]-N,N'-dimethylguanidine;

N-(5-acenaphthyl)-N'-(1-methyl-2-phenoxyethyl)guanidine;

N-(5-acenaphthyl)-N'-(1-methyl-2-phenoxyethyl)-N-methylguanidine;

N-(5-acenaphthyl)-N'-(1-methyl-2-phenoxyethyl)-N'-methylguanidine;


N-(5-acenaphthyl)-N'-(1-methyl-2-phenoxyethyl)-N,N'-dimethylguanidine;

N-(5-acenaphthyl)-N'-(1-methyl-2-(4-chlorophenyl)ethyl)guanidine;

N-(5-acenaphthyl)-N'-(1-methyl-2-(4-chlorophenyl)ethyl)-N-methylguanidine;

N-(5-acenaphthyl)-N'-(1-methyl-2-(4-chlorophenyl)ethyl)-N'-methylguanidine;
N-(5-acenaphthyl)-N'-(1-methyl-2-(4-chlorophenyl)ethyl)-N,N'-
dimethylguanidine;
N-(5-acenaphthyl)-N'-(1,2-diphenylethyl)guanidine;
N-(5-acenaphthyl)-N'-(1,2-diphenylethyl)-N-methylguanidine;
N-(5-acenaphthyl)-N'-(1,2-diphenylethyl)-N'-methylguanidine;
N-(5-acenaphthyl)-N'-(1,2-diphenylethyl)-N,N'-dimethylguanidine;
N-(5-acenaphthyl)-N'-(3-phenylpropyl)guanidine;
N-(5-acenaphthyl)-N'-(3-phenylpropyl)-N-methylguanidine;
N-(5-acenaphthyl)-N'-(2-methyl-2-phenylethyl)-N'-methylguanidine;
N,N'-(*sec*-butylphenyl)-N'-(2-phenoxyethyl)guanidine;
N,N'-(*sec*-butylphenyl)-N'-(2-phenoxyethyl)-N-methylguanidine;
N,N'-(*sec*-butylphenyl)-N'-(2-phenoxyethyl)-N'-methylguanidine;
N,N'-(*sec*-butylphenyl)-N'-(2-phenoxyethyl)-N,N'-dimethylguanidine;
N-(5-acenaphthyl)-N'-((4-*tert*-butylphenyl)-(4-*sec*-butylphenyl)-
methyl)guanidine;
N-(5-acenaphthyl)-N'-((4-*tert*-butylphenyl)-(4-*sec*-butylphenyl)-methyl)-N-
methylguanidine;
N-(5-acenaphthyl)-N'-((4-*tert*-butylphenyl)-(4-*sec*-butylphenyl)-methyl)-N'-
methylguanidine;
N-(5-acenaphthyl)-N'-((4-*tert*-butylphenyl)-(4-*sec*-butylphenyl)-methyl)-N,N'-
dimethylguanidine;
N-(4-butoxyphenyl)-N,N'-bis(4-*tert*-butylbenzyl)guanidine;
N-(4-butoxyphenyl)-N,N'-bis(4-*tert*-butylbenzyl)-N-methylguanidine;
N-(4-butoxyphenyl)-N,N'-bis(4-*tert*-butylbenzyl)-N'-methylguanidine;
N-(4-butoxyphenyl)-N,N'-bis(4-*tert*-butylbenzyl)-N,N'-dimethylguanidine;
and pharmaceutically acceptable salts of said compounds.

11. A compound having the following Formula IV:



wherein

each R is independently halo, hydroxy, amino, nitro, substituted or unsubstituted alkyl having from 3 to about 10 carbon atoms, substituted or unsubstituted alkoxy, substituted or unsubstituted aryloxy, substituted or unsubstituted aralkoxy, substituted or unsubstituted aminoalkyl, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfinyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted alkenyl having 3 to about 10 carbon atoms, or substituted or unsubstituted alkynyl having 3 to about 10 carbon atoms, at least one R group is a meta or para substituent;

n is an integer of from 1 to 5;

R¹ is substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having 1 to about 20 carbon atoms, substituted or unsubstituted aryloxy having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfonyl having 1 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least 5 ring atoms, substituted or unsubstituted aralkyl having at least 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having 1 to 3 rings, 3 to 8 ring members in each ring and 1 to 3 heteroatoms;


R² and R³ are each independently hydrogen or a group as defined for R¹ above; or R² and R³ are taken together to form a substituted or unsubstituted alkylene linkage of from 2 to about 6 carbon atoms; and wherein said compound exhibits high inhibition of glutamate release; and pharmaceutically acceptable salts thereof.

12. A compound of claim 11 selected from the group of:

N,N'-di-(4-*sec*-butylphenyl)guanidine;
N,N'-di-(4-*sec*-butylphenyl)-N-methylguanidine;
N,N'-di-(4-*sec*-butylphenyl)-N,N'-dimethylguanidine;
N-(2-naphthyl)-N'-(4-*isopropyl*phenyl)guanidine;
N-(2-naphthyl)-N'-(4-*isopropyl*phenyl)-N-methylguanidine;
N-(2-naphthyl)-N'-(4-*isopropyl*phenyl)-N'-methylguanidine;
N-(2-naphthyl)-N'-(4-*isopropyl*phenyl)-N,N'-dimethylguanidine;
N,N'-bis(4-*tert*-butylphenyl)guanidine;
N,N'-bis(4-*tert*-butylphenyl)-N-methylguanidine;
N,N'-bis(4-*tert*-butylphenyl)-N'-methylguanidine;
N,N'-bis(4-*tert*-butylphenyl)-N,N'-dimethylguanidine;
N-(4-*sec*-butylphenyl)-N'-(2,3,4-trichlorophenyl)guanidine;
N-(4-*sec*-butylphenyl)-N'-(2,3,4-trichlorophenyl)-N-methylguanidine;
N-(4-*sec*-butylphenyl)-N'-(2,3,4-trichlorophenyl)-N'-methylguanidine;
N-(4-*sec*-butylphenyl)-N'-(2,3,4-trichlorophenyl)-N,N'-dimethylguanidine;
N-(4-methoxy-1-naphthyl)-N'-(2,3,4-trichlorophenyl)guanidine;
N-(4-methoxy-1-naphthyl)-N'-(2,3,4-trichlorophenyl)-N-methylguanidine;
N-(4-methoxy-1-naphthyl)-N'-(2,3,4-trichlorophenyl)-N'-methylguanidine;
N-(4-methoxy-1-naphthyl)-N'-(2,3,4-trichlorophenyl)-N,N'-dimethylguanidine;
N,N'-bis-(4-*sec*-butylphenyl)-2-iminopyrimidazolidine;
N,N'-bis(3-biphenyl)guanidine;
N,N'-bis(3-biphenyl)-N-methylguanidine;
N,N'-bis(3-biphenyl)-N'-methylguanidine;
N,N'-bis(3-biphenyl)-N,N'-dimethylguanidine;

N,N'-di-(3-*tert*-butylphenyl)guanidine;
N,N'-di-(3-*tert*-butylphenyl)-N-methylguanidine;
N,N'-di-(3-*tert*-butylphenyl)-N'-methylguanidine;
N,N'-di-(3-*tert*-butylphenyl)-N,N'-dimethylguanidine;
N,N'-bis-(4-methoxy-1-naphthyl)guanidine;
N,N'-bis-(4-methoxy-1-naphthyl)-N-methylguanidine;
N,N'-bis-(4-methoxy-1-naphthyl)-N'-methylguanidine;
N,N'-bis-(4-methoxy-1-naphthyl)-N,N'-dimethylguanidine;
N,N'-bis-(3-*sec*-butylphenyl)guanidine;
N,N'-bis-(3-*sec*-butylphenyl)-N-methylguanidine;
N,N'-bis-(3-*sec*-butylphenyl)-N'-methylguanidine;
N,N'-bis-(3-*sec*-butylphenyl)-N,N'-methylguanidine;
N,N'-bis(4-*n*-butylphenyl)-N-methylguanidine;
N,N'-bis(4-*n*-butylphenyl)-N'-methylguanidine;
N,N'-bis(4-*n*-butylphenyl)-N,N'-dimethylguanidine;
N,N'-(*sec*-butylphenyl)-N'-(*n*-pentyl)guanidine;
N,N'-bis(3-benzyloxyphenyl)guanidine;
N,N'-bis(3-benzyloxyphenyl)-N-methylguanidine;
N,N'-bis(3-benzyloxyphenyl)-N,N'-dimethylguanidine;
N,N'-bis(4-benzyloxyphenyl)guanidine;
N,N'-bis(4-benzyloxyphenyl)-N-methylguanidine;
N,N'-bis(4-benzyloxyphenyl)-N,N'-dimethylguanidine;
N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)guanidine;
N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N-methylguanidine;
N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-methylguanidine;
N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N,N'-dimethylguanidine;
and pharmaceutically acceptable salts of said compounds.

13. A compound of the following Formula V:



wherein:

R is substituted or unsubstituted tetrahydroquinoliny, indoliny or piperonyl;

R¹ and R² are each independently hydrogen, substituted or unsubstituted alkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkenyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkynyl having from 2 to about 20 carbon atoms, substituted or unsubstituted alkoxy having from 1 to about 20 carbon atoms, substituted or unsubstituted aminoalkyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylthio having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfinyl having from 1 to about 20 carbon atoms, substituted or unsubstituted alkylsulfonyl having from 1 to about 20 carbon atoms, substituted or unsubstituted carbocyclic aryl having at least 5 ring atoms, substituted or unsubstituted aralkyl having at least 5 ring atoms, or a substituted or unsubstituted heteroaromatic or heteroalicyclic group having 1 to 3 rings, 3 to 8 ring members in each ring and 1 to 3 heteroatoms;

and pharmaceutically acceptable salts thereof.

14. A compound of claim 13 selected from the group of:

N-(5-acenaphthyl)-N'-(1,2,3,4-tetrahydroquinoliny)guanidine;

N-(5-acenaphthyl)-N'-(1,2,3,4-tetrahydroquinoliny)-N-methylguanidine;

N-(5-acenaphthyl)-N'-(1,2,3,4-tetrahydroquinoliny)-N'-methylguanidine;

N-(5-acenaphthyl)-N'-(1,2,3,4-tetrahydroquinoliny)-N,N'-dimethylguanidine;
N-(3-acenaphthyl)-N'-(indoliny)guanidine;
N-(3-acenaphthyl)-N'-(indoliny)-N-methylguanidine;
N-(3-acenaphthyl)-N'-(indoliny)-N'-methylguanidine;
N-(3-acenaphthyl)-N'-(indoliny)-N,N'-methylguanidine;
N-(5-acenaphthyl)-N'-(piperonyl)guanidine;
N-(5-acenaphthyl)-N'-(piperonyl)-N-methylguanidine;
N-(5-acenaphthyl)-N'-(piperonyl)-N'-methylguanidine;
N-(5-acenaphthyl)-N'-(piperonyl)-N,N'-dimethylguanidine;
and pharmaceutically acceptable salts of such compounds.

15. A compound of claims selected from the group of:

N-(3-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-*tert*-butylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(3-pentoxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(5-acenaphthyl)-N-(4-benzyloxybenzyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-benzyloxybenzyl)guanidine;
N-(4-benzyloxyphenyl)-N-(4-benzyloxybenzyl)guanidine;
N-(5-acenaphthyl)-N-(3-benzyloxybenzyl)guanidine;
N-(4-isopropylphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(4-benzyloxyphenyl)-N-(4-*tert*-butylbenzyl)guanidine;
N-(4-hexylphenyl)-N-(4-hexylbenzyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*t*-butylbenzyl)-N'-pyrrolidinylguanidine;
N-(4-*sec*-butylphenyl)-N-(4-*t*-butylbenzyl)-N'-(4-thiomorpholiny)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-piperidinylguanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-morpholiny)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-propylpiperidinyl)guanidine;
N-(4-butoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-piperidinyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-benzylpiperidinyl)guanidine;

N-(4-benzyloxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-morpholinyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(1,2,3,4-tetrahydroisoquinoliny)guanidine;
N-(3-butoxy-4-methoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-morpholinyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(3,5-dimethyl-4-morpholinyl)guanidine;
N-(4-*tert*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-*sec*-butylphenyl)-N'-(methyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-*sec*-butylphenyl)-N'-(methyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(phenyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-chlorophenyl)guanidine;
N-(4-butoxyphenyl)-N-(4-*tert*-butylbenzyl)-N'-(phenyl)guanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(phenyl)-N'-methylguanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(3,4-dichlorophenyl)guanidine;
N-(4-hexylphenyl)-N-(4-hexylbenzyl)-N'-phenylguanidine;
N-(4-*sec*-butylphenyl)-N-(4-*tert*-butylbenzyl)-N'-(4-benzyloxyphenyl)guanidine;
N,N'-bis-(4-*tert*-butylphenyl)-N,N'-dimethylguanidine
N-(4-benzyloxyphenyl)-N'-(4-*tert*-butylphenyl)guanidine;
N,N'-bis-(3-(1'-methyl-2'-phenyl)ethyl)guanidine;
N-methyl-N-4-benzyloxyphenyl-N'-(4-*tert*-butylphenyl)guanidine;
N,N'-bis-(4-hexylphenyl)guanidine;
N-(3-(1-(4'-ethoxy)benzyl)phenethyl)-N'-(4-*tert*-butylphenyl)guanidine;
N-(4-benzyloxyphenyl)-N'-methyl-N'-(4-*tert*-butylphenyl)guanidine;
N-(3-(4-*tert*-butylbenzyloxy)phenyl)-N'-(4-*tert*-butylphenyl)guanidine;
N-(3-(1'-benzylbutyl)phenyl)-N'-(4-*tert*-butylphenyl)guanidine;
N,N'-bis-(4-butylphenyl)-N-methylguanidine;
N,N'-bis-(4-*tert*-butylphenyl)-N,N'-dimethylguanidine;

N-(3-naphthylmethylenedioxyphenyl)-N'-(4-*tert*-butylphenyl)guanidine;
N-(4-benzyloxyphenyl)-N'-(4-butylphenyl)guanidine;
N,N'-bis-(4-butylphenyl)-N-butylguanidine;
N-3-(benzyloxymethyl)phenyl-N'-(4-*tert*-butylphenyl)guanidine;
N-(3,4-bis-butylphenoxyphenyl)-N'-(4-*tert*-butylphenyl)guanidine;
N-(3-benzyloxyphenyl)-N'-(4-*tert*-butylphenyl)guanidine;
N,N'-bis-(3-butoxy-4-methoxy)phenylguanidine;
N-(4-benzyloxyphenyl)-N-methyl-N'-(4-butylphenyl)guanidine;
N,N'-bis-(6-tetralinyl)guanidine;
N-(6-tetralinyl)-N'-(4-*tert*-butylphenyl)guanidine;
N-(5-acenaphthyl)-N'-(6-benzothiazolyl)guanidine;
N-(5-acenaphthyl)-N'-(6-N-benzylindolyl)guanidine;
N-(5-acenaphthyl)-N'-(4-benzo-2,1,3-thiadiazole)guanidine;
N-(5-acenaphthyl)-N'-[4-(6-methyl-benzothiazole)phenyl]guanidine;
N-(5-acenaphthyl)-N'-(1-benz[cd]indolyl)guanidine;
N-(5-acenaphthyl)-N'-(6-benz[cd]indo-2[1H]-one)guanidine;
N,N'-bis(6-benz[cd]indolyl-2[1H]-one)guanidine;
N-(4-butoxyphenyl)-N'-(4-chlorophenylethyl)guanidine;
N-(4-benzyloxyphenyl)-N,N'-diphenylguanidine;
N-(4-benzyloxyphenyl)-N'-benzyl-N'-phenylguanidine;
N-(3-benzyloxyphenyl)-N'-(4-thiobenzylphenyl)guanidine;
N,N'-bis(4-(phenylthio)phenyl)guanidine;
N,N'-bis(3-(phenylthio)phenyl)guanidine;
N-(5-acenaphthyl)-N'-(2-phenylethyl)guanidine;
N-(5-acenaphthyl)-N'-(3-butoxypropyl)guanidine;
N,N'-bis(2,2-diphenylethyl)guanidine;
N-(4-butoxyphenyl)-N'-(4-chlorophenylethyl)guanidine;
N-(4-butoxyphenyl)-N-(4-chlorobenzhydryl)guanidine;
(5-acenaphthyl)-N'-(phenethyl)-N'-benzylguanidine;
N-4-benzyloxyphenyl)-N'-(3-benzyloxyphenyl)-N'-(4-chlorobenzyl)guanidine;
N,N'-bis(4-benzyloxyphenyl)-N'-methylguanidine;

N-(4-benzyloxyphenyl)-N'-(3-benzyloxyphenyl)-N'-(4-chlorobenzyl)guanidine;
N-(3-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-phenylguanidine;
N-(4-sec-butylphenyl)-N'-(4-isopropoxyphenyl)-N'-phenylguanidine;
N-(4-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-phenylguanidine;
N,N'-bis(3-octyloxyphenyl)guanidine;
N,N'-bis(4-butoxyphenyl)guanidine;
N,N'-bis(4-phenoxyphenyl)guanidine;
N-(3-benzyloxyphenyl)-N'-(4-phenoxyphenyl)guanidine;
N-(3-benzyloxyphenyl)-N'-(4-phenylazophenyl)guanidine;
N,N'-bis(3-benzyloxyphenyl)-N'-methylguanidine;
N-(4-benzyloxyphenyl)-N'-(4-benzyloxyphenyl)-N'-methylguanidine;
N-(4-butoxyphenyl)-N'-(4-isopropoxyphenyl)guanidine;
N,N'-bis(4-(1-hydroxybutyl)phenyl)guanidine;
N-(4-butoxyphenyl)-N'-(3-methoxyphenyl)-N'-phenylguanidine;
N-(4-secbutylphenyl)-N'-phenyl-N'-(4-(2-isopropoxy)phenyl)guanidine;
N-(4-n-butoxyphenyl)-N'-(2-(4-chlorophenyl)ethyl)guanidine;
and pharmaceutically acceptable salts thereof.

16. A method for treating a disorder of the nervous system in which the pathophysiology of the disorder involves excessive or inappropriate release of a neurotransmitter from neuronal cells comprising administering to a mammal exhibiting symptoms of said disorder or susceptible to said disorder an effective amount of a compound of any of the preceding claims.

17. A method for treating nerve cell death comprising administering to a mammal exhibiting symptoms of nerve cell death or susceptible to nerve cell death an effective amount of a compound of any one of claims 1-15.

18. A method for treating a person who has suffered or is suffering from a stroke, or who is susceptible to a stroke, comprising administering to the person an effective amount of a compound of any one of claims 1-15.

19. A method of treating a disease of the cardiovascular system comprising administering to a mammal suffering from or susceptible to a cardiovascular disease an effective amount of a compound of any one of claims 1-15.

20. A method of modulating the release of excess endogenous neurotransmitters from a subject comprising administering to the subject an effective amount of a compound of any one of claims 1-15.

21. A method for treating a disease in which the pathophysiology of the disease involves inappropriate cellular secretion comprising administering to a mammal suffering from or susceptible to said disease an effective amount of a compound of any one of claims 1-15.

22. A method for 1) blocking voltage-activated calcium channels of mammalian neuronal cells; 2) blocking voltage-activated sodium channels of mammalian neuronal cells; 3) blocking voltage sensitive calcium channels of mammalian cardiac cells; 4) blocking voltage-activated sodium channels of mammalian cardiac cells; 5) blocking voltage-activated calcium channels of mammalian smooth or skeletal muscle cells; or 6) blocking voltage-activated sodium channels of mammalian smooth or skeletal muscle cells, comprising administering to said cells an effective amount of a compound of any one of claims 1-15.

23. A method for diagnosing a selected disease the pathophysiology of which involves ion-channel excitation or activity comprising administering to a mammal suffering from the disease a

detection effective amount of a labeled compound of any one of claims 1-15.

24. A pharmaceutical composition comprising a therapeutically effective amount of one or more compounds of any one of claims 1-15 and a pharmaceutically acceptable carrier.

DIKE, BRONSTEIN, ROBERTS & CUSHMAN
130 Water Street
Boston, Massachusetts 02109

DECLARATION AND POWER OF ATTORNEY
Original Application

COPY

As a below named inventor, I hereby declare that: My residence, post office address and citizenship are as stated below next to my name. I believe I am the original, first and sole inventor (if only one name is listed at 201) below or an original, first and joint inventor (if plural names are listed at 201-208 below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

THERAPEUTIC GUANIDINES

which is described and claimed in: ☒ the specification in application Serial Number 08/482,984, filed on June 7, 1995.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a). I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

I hereby claim the benefit under Title 35, United States code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)	(Filing Date)	(Status) (Patented, pending, abandoned)
PCT/US95/01536	03 February 1995	Pending
08/191,793	03 February 1994	Abandoned

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) with full powers of association, substitution and revocation to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (list name and registration number)

Sewell P. Bronstein (Reg. No. 16,919)	Ernest V. Linck (Reg. No. 29,822)	David S. Resnick (Reg. No. 34,235)
Dorinda Brown (Reg. No. 20,845)	Linda M. Buckley (Reg. No. 31,003)	Peter F. Corless (Reg. No. 33,860)
Robert L. Goldberg (Reg. No. 22,456)	Ronald I. Eisenstein (Reg. No. 30,628)	Kevin J. Fournier (Reg. No. 34,333)
David G. Conlin (Reg. No. 27,026)	Henry D. Pahl, Jr. (Reg. No. 20,438)	Brian L. Michaelis (Reg. No. 34,221)
George W. Neuner (Reg. No. 26,964)	Peter J. Manus (Reg. No. 26,766)	

SEND CORRESPONDENCE TO:

David G. Conlin
Dike, Bronstein, Roberts & Cushman
130 Water Street
Boston, Massachusetts 02109

DIRECT TELEPHONE CALLS TO:

(name and telephone number)

Peter F. Corless
(617) 523-3400

201	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE	
		Goldin	Stanley	M.
		Lexington	Massachusetts	United States
		10 Russell Road	Lexington	Massachusetts 02173

202	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE	
		Fischer	James	B.
		Cambridge	Massachusetts	United States
		1524 Cambridge Street	Cambridge	Massachusetts 02139

203	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE	
		Knapp	Andrew	Gannett
		Salem	Massachusetts	United States
		15 Warren Street	Salem	Massachusetts 01970

204	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE	
		Reddy	N. Laxma	
		Malden	Massachusetts	India
		24 Grover St., Apt. #4	Malden	Massachusetts 02148

(10)P/W

2 0 5	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Berlove	David	
		Cambridge	Massachusetts	United States
		1200 Massachusetts Avenue	Cambridge	Massachusetts 02138

2 0 6	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Durant	Graham	J.
		Marshfield	Massachusetts	United Kingdom
		55 Captain Luther Little Way	Marshfield	Massachusetts 02050

2 0 7	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Katragadda	Subbarao	
		Belmont	Massachusetts	United States
		65 Hawthorne Street	Belmont	Massachusetts 02178

2 0 8	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Hu	Lain-Yen	
		Bedford	Massachusetts	R.O.C. Taiwan
		#16 Old Stagecoach Road	Bedford	Massachusetts 01730

2 0 9	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Magar	Sharad	
		Somerville	Massachusetts	India
		303 Lowell Street, #14	Somerville	Massachusetts 02145

2 1 0	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Fan	Wenhong	
		Boston	Massachusetts	P.R. of China
		91 Westland Avenue, P.O. B. 10	Boston	Massachusetts 02115

2 1 1	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Yost	Elizabeth	
		Waltham	Massachusetts	United States
		125 Overland Road, Apt. 4	Waltham	Massachusetts 02154

2 1 2	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Guo	Jun Qing	
		Waltham	Massachusetts	P.R. of China
		16 Garden Lane, Apt. 4	Waltham	Massachusetts 02154

COPY

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Signature of Inventor 201	Signature of Inventor 202 <i>James B. Fink</i>	Signature of Inventor 203 <i>Andrew G. Knapp</i>
Date	Date <i>10/12/95</i>	Date <i>10/12/95</i>

Signature of Inventor 204 <i>N. Reddy</i>	Signature of Inventor 205 <i>Daniel Berlow</i>	Signature of Inventor 206 <i>Guahan Duxant</i>
Date <i>10/13/95</i>	Date <i>10/13/95</i>	Date <i>10/12/95</i>

Signature of Inventor 207 <i>S. K. Raghadda</i>	Signature of Inventor 208 <i>Lainjen Hu</i>	Signature of Inventor 209 <i>Shaga</i>
Date <i>10/12/1995</i>	Date <i>10/30/95</i>	Date <i>10/13/95</i>

Signature of Inventor 210	Signature of Inventor 211 <i>Elizabeth M. Mat</i>	Signature of Inventor 212 <i>Ingjeng Gao</i>
Date	Date <i>Oct. 13, 1995</i>	Date <i>10/13/95</i>

DIKE, BRONSTEIN, ROBERTS & CUSHMAN
130 Water Street
Boston, Massachusetts 02109

DECLARATION AND POWER OF ATTORNEY Original Application

As a below named inventor, I hereby declare that: My residence, post office address and citizenship are as stated below next to my name. I believe I am the original, first and sole inventor (if only one name is listed at 201) below or an original, first and joint inventor (if plural names are listed at 201-208 below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

THERAPEUTIC GUANIDINES

which is described and claimed in: ☒ the specification in application Serial Number 08/482,984,
filed on June 7, 1995.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a). I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

I hereby claim the benefit under Title 35, United States code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)	(Filing Date)	(Status) (Patented, pending, abandoned)
PCT/US95/01536	03 February 1995	Pending
08/191,793	03 February 1994	Abandoned

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) with full powers of association, substitution and revocation to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (list name and registration number)

Swall P. Bronstein (Reg. No. 16,919)	Ernest V. Linck (Reg. No. 29,822)	David S. Rennie (Reg. No. 34,235)
Donald Brown (Reg. No. 20,845)	Linda M. Buckley (Reg. No. 31,003)	Peter F. Corless (Reg. No. 33,860)
Robert L. Goldberg (Reg. No. 22,456)	Ronald I. Eisenstein (Reg. No. 30,628)	Kevin J. Fournier (Reg. No. 34,333)
David G. Conlin (Reg. No. 27,026)	Henry D. Pahl, Jr. (Reg. No. 20,438)	Brian L. Michaelis (Reg. No. 34,221)
George W. Neuner (Reg. No. 26,964)	Peter J. Manus (Reg. No. 26,766)	

SEND CORRESPONDENCE TO: David G. Conlin Dike, Bronstein, Roberts & Cushman 130 Water Street Boston, Massachusetts 02109	DIRECT TELEPHONE CALLS TO: (name and telephone number) Peter F. Corless (617) 523-3400
--	--

201	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
		Goldin	Stanley	M.
202	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
		Lexington	Massachusetts	United States
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		10 Russell Road	Lexington	Massachusetts 02173

202	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
		Fischer	James	B.
203	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
		Cambridge	Massachusetts	United States
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		1524 Cambridge Street	Cambridge	Massachusetts 02139

203	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
		Knapp	Andrew	Gannett
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
		Salem	Massachusetts	United States
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		15 Warren Street	Salem	Massachusetts 01970

204	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
		Reddy	N. Laxma	
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
		Malden	Massachusetts	India
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		24 Grover St., Apt. #4	Malden	Massachusetts 02148

2 9 5	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Berlova	David	
		Cambridge	Massachusetts	United States
		1200 Massachusetts Avenue	Cambridge	Massachusetts 02138

2 9 6	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Durant	Graham	J.
		Marshfield	Massachusetts	United Kingdom
		55 Captain Luther Little Way	Marshfield	Massachusetts 02050

2 9 7	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Katragadda	Subbarao	
		Belmont	Massachusetts	United States
		65 Hawthorne Street	Belmont	Massachusetts 02178

2 9 8	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Hu	Lain-Yen	
		Bedford	Massachusetts	R.O.C. Taiwan
		#16 Old Stagecoach Road	Bedford	Massachusetts 01730

2 9 9	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Magar	Sharad	
		Somerville	Massachusetts	India
		303 Lowell Street, #14	Somerville	Massachusetts 02145

2 1 0	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Fan	Wenhong	
		Boston	Massachusetts	P.R. of China
		91 Westland Avenue, P.O. B. 10	Boston	Massachusetts 02115

2 1 1	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Yost	Elizabeth	
		Waltham	Massachusetts	United States
		125 Overland Road, Apt. 4	Waltham	Massachusetts 02154

2 1 2	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Guo	Jun Qing	
		Waltham	Massachusetts	P.R. of China
		16 Garden Lane, Apt. 4	Waltham	Massachusetts 02154

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Signature of Inventor 201 <i>Stanley M. Goldin</i>	Signature of Inventor 202	Signature of Inventor 203
Date <i>11/20/95</i>	Date	Date

Signature of Inventor 204	Signature of Inventor 205	Signature of Inventor 206
Date	Date	Date

Signature of Inventor 207	Signature of Inventor 208	Signature of Inventor 209
Date	Date	Date

Signature of Inventor 210	Signature of Inventor 211	Signature of Inventor 212
Date	Date	Date

DIKE, BRONSTEIN, ROBERTS & CUSHMAN
130 Water Street
Boston, Massachusetts 02109

DECLARATION AND POWER OF ATTORNEY Original Application

As a below named inventor, I hereby declare that: My residence, post office address and citizenship are as stated below next to my name. I believe I am the original, first and sole inventor (if only one name is listed at 201) below or an original, first and joint inventor (if plural names are listed at 201-208 below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

THERAPEUTIC GUANIDINES

which is described and claimed in: ☒ the specification in application Serial Number 08/482,984,
filed on June 5, 1995.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a). I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

I hereby claim the benefit under Title 35, United States code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)	(Filing Date)	(Status) (Patented, pending, abandoned)
PCT/US95/01536	03 February 1995	Pending
08/191,793	03 February 1994	Abandoned

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) with full powers of association, substitution and revocation to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (list name and registration number)

Sewall P. Bronstein (Reg. No. 16,919)	Ernest V. Linek (Reg. No. 29,822)	David S. Resnick (Reg. No. 34,235)
Donald Brown (Reg. No. 20,845)	Linda M. Buckley (Reg. No. 31,003)	Peter F. Corless (Reg. No. 33,860)
Robert L. Goldberg (Reg. No. 22,456)	Ronald I. Eisenstein (Reg. No. 30,628)	Kevin J. Fournier (Reg. No. 34,333)
David G. Conlin (Reg. No. 27,026)	Henry D. Pahl, Jr. (Reg. No. 20,438)	Brian L. Michaelis (Reg. No. 34,221)
George W. Neuner (Reg. No. 26,964)	Peter J. Manus (Reg. No. 26,766)	

SEND CORRESPONDENCE TO:

David G. Conlin
Dike, Bronstein, Roberts & Cushman
130 Water Street
Boston, Massachusetts 02109

DIRECT TELEPHONE CALLS TO:

(name and telephone number)

Peter F. Corless
(617) 523-3400

201	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE	
		Goldin	Stanley	M.
		Lexington	Massachusetts	United States
		10 Russell Road	Lexington	Massachusetts 02173

202	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE	
		Fischer	James	B.
		Cambridge	Massachusetts	United States
		1524 Cambridge Street	Cambridge	Massachusetts 02139

203	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE	
		Knapp	Andrew	Gannett
		Salem	Massachusetts	United States
		15 Warren Street	Salem	Massachusetts 01970

204	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE	
		Reddy	N. Laxma	
		Malden	Massachusetts	India
		24 Grover St., Apt. #4	Malden	Massachusetts 02148

2 0 5	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Berlove	David	
		Cambridge	Massachusetts	United States
		1200 Massachusetts Ave	Cambridge	Massachusetts 02138

2 0 6	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Durant	Graham	J.
		Marshfield	Massachusetts	United Kingdom
		55 Captain Luther Little Way	Marshfield	Massachusetts 02050

2 0 7	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Katragadda	Subbarao	
		Belmont	Massachusetts	United States
		65 Hawthorne Street	Belmont	Massachusetts 02178

2 0 8	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Hu	Lain-Hu	
		Bedford	Massachusetts	R.O.C. Taiwan
		#16 Old Stagecoach Road	Bedford	Massachusetts 01730

2 0 9	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Magar	Sharad	
		Somerville	Massachusetts	India
		303 Lowell Street, #14	Somerville	Massachusetts 02145

2 1 0	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Fan	Wenhong	
		Rockey Hill	Connecticut	P.R. of China
		4-D Kangsley Court	Rockey Hill	Connecticut 06067

2 1 1	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Yost	Elizabeth	
		Waltham	Massachusetts	United States
		125 Overland Road, Apt. 4	Waltham	Massachusetts 02154

2 1 2	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY AND ZIP CODE
		Guo	Jun Qing	
		Waltham	Massachusetts	P.R. of China
		16 Garden Lane, Apt. 4	Waltham	Massachusetts 02154

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Signature of Inventor 201	Signature of Inventor 202	Signature of Inventor 203
Date	Date	Date

Signature of Inventor 204	Signature of Inventor 205	Signature of Inventor 206
Date	Date	Date

Signature of Inventor 207	Signature of Inventor 208	Signature of Inventor 209
Date	Date	Date

Signature of Inventor 210 <i>Wenhong Fan</i>	Signature of Inventor 211	Signature of Inventor 212
Date <i>12/4/95</i>	Date	Date